A simple one-step procedure for the separation of calpain I, calpain II and calpastatin

Jan-Olof KARLSSON, Staffan GUSTAVSSON, Christina HALL and Elisabet NILSSON

Institute of Neurobiology, University of Göteborg, P.O. Box 33 031, S-400 33 Göteborg, Sweden

The soluble fraction from rabbit brain was adsorbed on a column of phenyl-Sepharose. By applying a linear gradient with decreasing salt concentration and increasing pH, it was possible to separate calpain I and calpain II from each other and from the endogenous inhibitor calpastatin. Both enzymes were capable of degrading endogenously labelled neuronal proteins, including slowly axonally transported soluble proteins and rapidly transported membrane-bound proteins, as well as casein.

INTRODUCTION

A neutral Ca^{2+} -activated proteinase that required cysteine for activity was described in rat brain by Guroff (1964). Since then, a family of Ca^{2+} -dependent cysteine proteinases has been found in almost all mammalian and avian cells, together with an endogenous specific inhibitor (for review see Murachi, 1983). This group of proteinases has been called calpains, and the endogenous inhibitor termed calpastatin.

Calpains (EC 3.4.22.17) with a high Ca^{2+} -sensitivity (calpain I) and with a low Ca^{2+} -sensitivity (calpain II) have usually been separated from each other via $(\text{NH}_4)_2\text{SO}_4$ precipitation and DEAE-cellulose ion-exchange chromatography (Mellgren, 1980; Dayton *et al.*, 1981; Murachi *et al.*, 1981; Malik *et al.*, 1983; Yoshimura *et al.*, 1983; Inomata *et al.*, 1983; Croall & De Martino, 1984; Zimmermann & Schlaepfer, 1984).

However, DEAE-cellulose ion-exchange chromatography does not separate calpain I from calpastatin (see, e.g., Inomata *et al.*, 1983; Yoshimura *et al.*, 1983), and additional procedures are therefore required to separate these proteins.

Most of the work cited above used tissue that was not perfused, and that was subjected to handling for some period before homogenization. Separation of calpains I and II, and calpastatin, from each other usually took several days, with numerous purification steps.

The procedure now presented includes rapid perfusion of the tissue with EDTA before homogenization and a one-step procedure to separate calpains I and II from each other, and from the endogenous inhibitor, calpastatin. The entire procedure takes only 1 day and uses only one chromatographic step.

EXPERIMENTAL

Materials

Casein (nach Hammersten) was from Merck. Dithiothreitol was from Sigma Chemical Co. Phenyl-Sepharose 4B was supplied from Pharmacia. Other chemicals were of analytical grade.

General procedure

An albino rabbit, weighing approx. 2 kg, was anaesthetized with an intravenous injection of sodium pentobarbital (50 mg/kg) and perfused with 1 litre of

ice-cold 5 mM-EDTA (adjusted to a pH of 8 with NaOH) through the left heart ventricle at a pressure of 150 cmH₂O. The brain (wet weight approx. 7 g) was immediately dissected out and homogenized in 30 ml of ice-cold 10 mM-sodium borate buffer, pH 8, containing 5 mM-EDTA, 3 mM- NaN_3 and 1 mM-dithiothreitol. The homogenate was homogenized twice in a Teflon/glass homogenizer with six up-and-down strokes at 1500 rev./min. After dilution with buffer to 70 ml, the sample was centrifuged at 90000 g for 2 h at 4 °C. The supernatant obtained was decanted, and solid NaCl was added to a final concentration of 200 mM. The sample was then allowed to reach room temperature (21 °C) and adjusted to pH 7.00 with NaOH or HCl. All subsequent work was carried out at room temperature.

Phenyl-Sepharose chromatography

The sample was applied to a column (15 cm × 0.8 cm) of phenyl-Sepharose equilibrated in 10 mM-sodium borate buffer, pH 7.00, containing 200 mM-NaCl, 3 mM- NaN_3 , 1 mM-dithiothreitol and 0.1 mM-EDTA at a rate of 50 ml/h. The column was washed with large volumes of equilibrating buffer and was then eluted with a linear gradient consisting of 150 ml of equilibrating buffer as starting buffer and 150 ml of 5 mM-sodium borate buffer, pH 8.0, containing 1.5 mM- NaN_3 , 1 mM-dithiothreitol and 0.1 mM-EDTA as final buffer. The flow rate during the gradient elution was approx. 20 ml/h. Fractions of volume size 10 ml were collected, and 200 μ l portions were assayed for proteolytic activity.

Proteinase assay

Casein. Casein was alkali-treated as described by Ashby & Walsh (1974) and stored at –20 °C in portions containing 5 mg of casein/ml in 50 mM-sodium borate buffer, pH 7.5, 3 mM- NaN_3 and 0.1 mM-EDTA. Immediately before an experiment the casein substrate was thawed and made to 1 mM in dithiothreitol and 4 mM in CaCl_2 (unless otherwise specified). A 500 μ l volume of substrate solution containing Ca^{2+} and dithiothreitol was added to 200 μ l of the enzyme solution in an open plastic tube. Incubations were for 1 h at 21 °C (unless otherwise stated), and were stopped by the addition of 200 μ l of 36% (w/v) trichloroacetic acid. The tubes were stored at 4 °C overnight. After centrifugation, 200 μ l of the trichloroacetic acid supernatant was mixed with 1.5 ml of

Fig. 1. Separation of calpain I and calpain II on a column of phenyl-Sepharose

For full experimental details see the text. —, A_{280} ; ·····, caseinolytic activity in the presence of 4 mM- Ca^{2+} [the incubation mixture contained 8 mM- Ca^{2+} when fractions (fractions 1–20) not bound to the phenyl-Sepharose were assayed]; ----, NaCl concentration. I and II denote proteolytic activity corresponding to the low- Ca^{2+} -requiring enzyme and the high- Ca^{2+} -requiring enzyme respectively.

0.5 M-sodium borate buffer, pH 8.5. Then 200 μl of fluorescamine solution (3.4 mg/10 ml of acetone) was added to each tube and the contents were mixed thoroughly. The amount of liberated amino groups was measured fluorimetrically (Nakai *et al.*, 1974) in a Farrand ratio fluorimeter and compared with standards of known amounts of glutamate in 10% trichloroacetic acid. One unit of proteinase activity is defined as the amount that causes the release of 1 μmol of amino groups/h during the described conditions. Zero-time values were always subtracted from values obtained after incubations. Results are expressed as units/ml of enzyme solution.

Endogenous labelled substrate. Rabbits were injected intraocularly with 50 μCi of L-[5- ^3H]proline (26 Ci/mmol). At different time intervals (20 h or 20 days) after this injection the animals were killed by an air embolus. The lateral geniculate bodies were dissected out and homogenized in 10 mM-Tris/HCl buffer, pH 7.5, containing 3 mM- NaN_3 and 1 mM-EDTA. The homogenate was centrifuged at 100 000 g for 60 min at 4 °C. The supernatant, containing soluble proteins, was diluted with homogenization buffer and stored in small portions at –80 °C. The pellet was resuspended in the same volume of buffer and stored in small portions at –80 °C. Portions of volume 200 μl were thawed and added to 800 μl of enzyme solution. CaCl_2 and MgCl_2 (5 mM and 2 mM respectively, final concentrations) were added to start degradation. The mixture was incubated in a shaking water bath at 39 °C for 1 h. Degradation was stopped by adding 100 μl of 100% trichloroacetic acid and 100 μl of bovine serum albumin (10 mg/ml). After centrifugation the radioactivity in the trichloroacetic acid supernatant was measured by liquid-scintillation counting. Proteolytic activity was defined as increase of trichloroacetic acid-soluble radioactivity during incubation (zero-time values subtracted). It should be noted that the substrate contained a certain amount of endogenous proteolytic activity that was detected in the absence of added enzyme (see the legend to Fig. 4).

Fig. 2. Proteolytic activity of calpain I and calpain II at 4 mM- Ca^{2+} on alkali-treated casein as a function of incubation time at 21 °C

For full experimental details see the text. ----, Calpain I; —, calpain II.

RESULTS

The amount of Ca^{2+} -dependent proteolytic activity in the high-speed supernatant from rabbit brain showed large variations between different experiments (results not shown). This may partly be due to the presence, in the extract, of endogenous inhibitor(s) of the enzymes.

When the supernatant was subjected to hydrophobic-interaction chromatography on a column of phenyl-Sepharose 4B (Fig. 1), the amount of Ca^{2+} -activated proteolytic activity in the eluate was very reproducible. With alkali-denatured casein as substrate, only two peaks (I and II) of proteolytic activity were detected in the

Fig. 3. Proteolytic activity of calpain I and calpain II at various concentrations of added Ca^{2+} to the assay mixture containing 0.1 mM-EDTA

Incubations were carried out for 30 min at 21 °C. For full experimental details see the text. ----, calpain I; —, calpain II. Proteolytic activity in the presence of 10 mM- Ca^{2+} is not significantly higher than the activity at 5 mM- Ca^{2+} .

presence of 4 mM- Ca^{2+} at neutral pH (Fig. 1). No proteolytic activity at all was detected in the absence of added Ca^{2+} .

Peak II was eluted at an approximate concentration of 100 mM-NaCl and at pH 7.4. Peak I was eluted at approx. 25 mM-NaCl and at pH 7.9. The relative amounts of proteolytic activity in the peaks compared with the total proteolytic activity eluted from the column were $31 \pm 3\%$ for peak I and $66 \pm 3\%$ for peak II (means $\pm$ S.D., $n = 5$).

Proteolytic degradation of casein was linear with time for up to 1 h at 21 °C (Fig. 2) at high Ca^{2+} concentrations. Proteolysis at 39 °C was linear with time for approx. 10 min. Proteolysis at 0 °C was negligible (results not shown).

Peaks I and II from the hydrophobic-interaction chromatography exhibited marked differences with respect to their Ca^{2+} requirements (Fig. 3). Peak I required 150 μM added Ca^{2+} for half-maximal activity, whereas peak II needed more than 1 mM- Ca^{2+} for half-maximal activity (Fig. 3). Peak I was assigned to calpain I and peak II to calpain II, in accordance with their Ca^{2+} requirements. The absolute activity of Ca^{2+} in the assay system is difficult to calculate at low concentrations of added Ca^{2+} . The assay system always contained 100 μM -EDTA. The substrate, casein, is a phosphoprotein with affinity for Ca^{2+} . Other proteins in the enzyme solution may also bind Ca^{2+} . It is probable that the free Ca^{2+} concentration in the assay system when 150 μM - Ca^{2+} is added is much below 50 μM . Addition of 200 μl portions of materials not bound to phenyl-Sepharose (fraction 8 in Fig. 1) to the assay system caused a complete inhibition of the proteolysis induced by calpain I. Calpain II was inhibited by 95% (results not shown).

Isolated calpain II was stable for about 1 week at 4 °C, whereas calpain I was stable for more than 1 month. Both enzymes showed a marked decrease in activity after freezing at -20 °C.

Calpains I and II were also capable of degrading slowly transported soluble proteins, as well as rapidly transported membrane-bound proteins, from axons and nerve terminals of retinal ganglion cells (Fig. 4). Besides peaks

Fig. 4. Separation of calpain I and calpain II on a column of phenyl-Sepharose, as shown by using activity against endogenously labelled substrates

Experimental conditions are similar to those indicated in Fig. 1 legend. —, Proteolytic activity against labelled rapidly transported membrane-bound proteins; the animal was injected with [^3H]proline 20 h before being killed; ·····, proteolytic activity against slowly transported soluble proteins; the animal was injected with [^3H]proline 20 days before being killed; ----, NaCl concentration. It should be noted that substrate containing labelled proteins showed an endogenous proteolytic activity (Ca^{2+} -stimulated), resulting in an elevated base-line of radioactivity. Material not bound to the phenyl-Sepharose is found in fractions 3–10. The major peaks of proteolytic activity (I and II) detected here corresponded exactly to peaks of caseinolytic activity. In this particular experiment the gradient consisted of 2×200 ml of buffer.

membrane-bound proteins, from axons and nerve terminals of retinal ganglion cells (Fig. 4). Besides peaks I and II, no other major proteolytic activity peak against the labelled substrates was detected in fractions from the phenyl-Sepharose column. It should be noted that the fractions that did not bind to the phenyl-Sepharose decreased the endogenous proteolytic degradation of the rapidly transported membrane proteins (Fig. 4). This effect was not apparent for the slowly transported soluble proteins.

DISCUSSION

The results presented here demonstrate that it is possible to separate calpain I and calpain II from each other, and from calpastatin, with a simple one-step procedure. The procedure is rapid, and the enzymes are obtained 1 day after the killing of the animal. The methodology is based on the difference in hydrophobicity between calpain I and calpain II (and the inhibitor). Hydrophobic-interaction chromatography has previously been used to separate calpain I from the endogenous inhibitor (Inomata *et al.*, 1983; Kawashima *et al.*, 1984) after prior separation on an ion-exchanger. Yoshimura *et al.* (1983) have suggested previously that rat kidney calpain I seems to be more hydrophobic, since it was eluted from a Blue Sepharose column more slowly than was calpain II. The present results with hydrophobic-interaction chromatography of the soluble fraction from rabbit brain clearly show that calpain I activity was more tightly bound to the hydrophobic matrix than was calpain II. In all nine experiments performed, calpain II was eluted from the phenyl-Sepharose in a larger volume, and occasionally with shoulders, compared with the relatively sharp peak of eluted calpain I activity. This may indicate that calpain II is more heterogeneous with respect to hydrophobicity compared with calpain I. Preliminary experiments with DEAE-cellulose chromatography of the separated enzymes do not indicate that calpain II is more heterogeneous than calpain I with respect to charge. Sodium dodecyl sulphate/polyacrylamide-gel electrophoresis of the fractions containing calpain I and calpain II revealed eight to 15 different polypeptide chains, including components of M_r 80000 and 20000.

The present study indicates that about one-third of the proteolytic activity in the soluble fraction from rabbit brain was due to calpain I, whereas the remaining proteolytic activity was due to calpain II. Similar results have been reported from rat brain supernatant (Zimmerman & Schlaepfer, 1984) and from rat brain synaptosomal membranes (Siman *et al.*, 1983). On the other hand, Malik *et al.* (1984) found more calpain I activity compared with calpain II in a calf brain synaptosomal fraction.

In agreement with many previous reports, alkali-denatured casein was a good substrate for the enzymes. It has been shown previously that calpains are able to degrade tubulin and neurofilament proteins (Malik *et al.*, 1983), as well as a number of different neuropeptides (Hirao & Takahashi, 1984). In addition, the present study demonstrates that both calpain I and calpain II have very broad specificities. Both enzymes were capable of

degrading slowly transported soluble proteins from axons of retinal ganglion cells. This fraction of neuronal proteins includes a large number of different polypeptides (for review see Karlsson, 1982). Rapidly transported membrane proteins in the neuron include hundreds of different polypeptide chains (Stone *et al.*, 1978; Wagner *et al.*, 1979; Karlsson, 1982). Both calpain I and calpain II were able to degrade a substantial part of these membrane-bound proteins.

It seems possible that the calpains are capable of degrading most of the cellular proteins as well as some exogenous proteins, such as casein, at least *in vitro*. We have previously shown that an endogenous Ca^{2+} -activated neutral proteinase is capable of degrading rapidly axonally transported neuronal membrane proteins in a homogenate (Gustavsson & Karlsson, 1985) and in intact tissue pieces (S. Gustavsson & J.-O. Karlsson, unpublished work).

This work was supported by the Swedish Medical Research Council (5932), Torsten and Ragnar Söderbergs Stiftelser and Magn. Bergvalls Stiftelse. We thank Mrs. Inger Jansson for excellent technical assistance, and Ms. Gull Grönstedt for careful secretarial work.

REFERENCES

- Ashby, C. D. & Walsh, D. A. (1974) *Methods Enzymol.* **38**, 350–358
- Croall, D. E. & De Martino, G. N. (1984) *Biochim. Biophys. Acta* **788**, 348–355
- Dayton, W. R., Schollmeyer, J. V., Lepley, R. A. & Cortés, L. R. (1981) *Biochim. Biophys. Acta* **659**, 48–61
- Guroff, G. (1964) *J. Biol. Chem.* **239**, 149–155
- Gustavsson, S. & Karlsson, J.-O. (1985) *Neurochem. Int.* **7**, 639–645
- Hirao, T. & Takahashi, K. (1984) *J. Biochem. (Tokyo)* **96**, 775–784
- Inomata, M., Hayashi, M., Nakamura, M., Imahori, K. & Kawashima, S. (1983) *J. Biochem. (Tokyo)* **93**, 291–294
- Karlsson, J.-O. (1982) in *Molecular Approaches to Neurobiology* (Brown, I. G., ed.), pp. 131–157, Academic Press, New York
- Kawashima, S., Nomoto, M., Hayashi, M., Inomata, M., Nakamura, M. & Imahori, K. (1984) *J. Biochem. (Tokyo)* **95**, 95–101
- Malik, M. N., Fenko, M. D., Iqbal, K. & Wisniewski, H. M. (1983) *J. Biol. Chem.* **258**, 8955–8962
- Malik, M. N., Fenko, M. D. & Wisniewski, H. M. (1984) *Neurochem. Res.* **9**, 233–240
- Mellgren, R. L. (1980) *FEBS Lett.* **109**, 129–133
- Murachi, T. (1983) *Trends Biochem. Sci.* **167**–169
- Murachi, T., Tanaka, K., Hatanaka, M. & Murakami, T. (1981) *Adv. Enzyme Regul.* **19**, 407–424
- Nakai, N., Lai, C. Y. & Horecker, B. L. (1974) *Anal. Biochem.* **58**, 563–570
- Siman, R., Baudry, M. & Lynch, G. (1983) *J. Neurochem.* **41**, 950–956
- Stone, G. C., Wilson, D. L. & Hall, M. E. (1978) *Brain Res.* **144**, 287–302
- Wagner, J. A., Kelly, A. S. & Kelly, R. B. (1979) *Brain Res.* **168**, 97–117
- Yoshimura, N., Kikuchi, T., Sasaki, T., Kitahara, A., Hatanaka, M. & Murachi, T. (1983) *J. Biol. Chem.* **258**, 8883–8889
- Zimmerman, U.-J. P. & Schlaepfer, W. W. (1984) *J. Biol. Chem.* **259**, 3210–3218