PRETEST AND OBJECTIVES

Articles are selected for CME credit designation on the basis of our assessment of the needs of readers of *The Primary Care Companion*, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. There are no prerequisites for participation in this CME activity.

To obtain credit, please study the designated article and complete the Posttest.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Designation

The CME Institute of Physicians Postgraduate Press, Inc. designates this educational activity for up to 1 Category 1 credit toward the American Medical Association Physician's Recognition Award. Each participant should claim only those credits that he/she actually spent in the educational activity.

Date of Original Release/Review

This educational activity is eligible for CME credit through June 30, 2007. The latest review of this material was May 2005.

Educational Objective

After studying the article by Kato et al., you will be able to:

- Select an appropriate body fat screening tool for predicting cardiovascular risk in patients with schizophrenia.

This pretest is designed to facilitate your study of the material.

1. **Waist circumference measurement is a subjective screening tool that can be used to identify risk for cardiovascular disease in the general population but not in patients with schizophrenia.**

- a. True
- b. False

Pretest answer and Posttest on page 119.

Disclosure of Off-Label Usage

The authors have determined that, to the best of their knowledge, no investigational information about pharmaceutical agents that is outside U.S. Food and Drug Administration–approved labeling has been presented in this article.

The Relation Between Body Fat Distribution and Cardiovascular Risk Factors in Patients With Schizophrenia: A Cross-Sectional Pilot Study

Martha M. Kato, M.D.; M. Beatriz Currier, M.D.;
Oscar Villaverde, M.D.; and Mercedes Gonzalez-Blanco, M.D.

Background: Obesity has recently become a concern for physicians treating schizophrenic patients. Obesity is associated with hypertension, dyslipidemia, and diabetes mellitus. In this pilot study, we investigate which anthropometric measurement, body mass index or waist circumference, is a better predictor of cardiovascular risk factors in patients with schizophrenia.

Method: This cross-sectional study, conducted from January 2001 to January 2002, examined body fat distribution and its relation to cardiovascular risk factors in 62 patients with schizophrenia (DSM-IV) recruited from an outpatient psychiatric clinic.

Results: Chi-square analysis revealed that an increased waist circumference was associated with dyslipidemia ($p < .01$), hypertension ($p < .05$), and abnormal serum glucose ($p < .05$), whereas an increased body mass index was only associated with dyslipidemia ($p < .05$). In logistic regression analysis, after controlling for age, gender, race, ethnicity, smoking, and body mass index, increased waist circumference remained significantly associated with dyslipidemia (odds ratio = 2.08, 95% CI = 1.01 to 1.15, $p < .05$) and hypertension (odds ratio = 2.05, 95% CI = 1.02 to 1.17, $p < .05$).

Conclusions: Waist circumference revealed a stronger correlation than body mass index to cardiovascular risk factors in patients with schizophrenia. We propose the measurement of waist circumference as a screening tool for cardiovascular risk factors in this population. Waist circumference measurement can provide an opportunity for primary prevention of coronary heart disease and diabetes mellitus in patients with schizophrenia.

(*Prim Care Companion J Clin Psychiatry* 2005;7:115-118)

Received Dec. 28, 2004; accepted April 7, 2005. From the Department of Psychiatry, University of Miami School of Medicine, Miami, Fla.

Presented in part at the 155th annual meeting of the American Psychiatric Association; May 18-23, 2002; Philadelphia, Pa.

In the spirit of full disclosure and in compliance with all ACCME Essential Areas and Policies, the faculty for this CME activity were asked to complete a full disclosure statement. The information received is as follows: Drs. Kato, Currier, Villaverde, and Gonzalez-Blanco have no significant commercial relationships to disclose relative to the presentation.

Corresponding author and reprints: Martha M. Kato, M.D., Department of Psychiatry, University of Miami School of Medicine, 1695 N.W. 9th Avenue, Room 2435, Miami, FL 33136 (e-mail: mkato@med.miami.edu).

Physicians need to be aware of the risks among schizophrenic patients of developing certain medical conditions in order to prevent and institute early and effective management of the comorbid medical illness. Weight gain has been associated with the use of both typical and atypical antipsychotic medications.¹⁻⁴ Many mechanisms have been postulated for the weight gain, but they remain speculative.

Hypertension, dyslipidemia, and diabetes mellitus are well-established cardiovascular risk factors. Obesity increases the risk for developing these medical conditions.⁵ Body mass index (BMI), which is defined as weight (kg)/height (m²), is the accepted measure to assess obesity.⁶ Obesity is defined by a BMI value ≥ 30 kg/m².⁶ BMI is an indicator of total body fat; however, it is not a good indicator of regional fat distribution. Studies suggest that fat distribution rather than total body fat is a better predictor for cardiovascular risk factors.^{7,8} Waist-hip ratio (WHR) is an index of body fat distribution. As the WHR is difficult to interpret biologically and less sensitive to changes in total body fat and visceral fat,⁹ waist circumference (WC) has been proposed to be a better predictor of cardiovascular risk factors.⁶ Recent guidelines recommend WC over the WHR as a predictor of obesity-related diseases, due to its simplicity and its correlation with abdominal fat as measured by computed tomography.⁶ Recent consensus guidelines for patients with schizophrenia recommend the measurement of both BMI and WC to monitor cardiovascular risk factors in this population.¹⁰

In this pilot cross-sectional study, we will investigate which anthropometric measurement, BMI or WC, is a

Table 1. Clinical Identification of Cardiovascular Risk Factors^a

Risk Factor	Defining Variable
Triglycerides	≥ 150 mg/dL
High-density lipoprotein cholesterol	
Men	< 35 mg/dL
Women	< 45 mg/dL
Low-density lipoprotein cholesterol	≥ 160 mg/dL
Blood pressure	
Systolic	≥ 140 mm Hg
Diastolic	≥ 90 mm Hg
Fasting serum glucose	≥ 126 mg/dL

^aBased on *The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults*¹¹ and the *Expert Panel on Detection and Treatment of High Blood Cholesterol in Adults*.¹²

better predictor of cardiovascular risk factors in patients with schizophrenia. The comparison of BMI and WC as predictor of cardiovascular risk factors has not been reported in patients with schizophrenia to our knowledge. The pattern of body fat distribution may be used by the treating physician to predict which patients are prone to develop cardiovascular risk factors, so that early preventive measurements can be taken.

METHOD

Patients with schizophrenia were recruited from an outpatient psychiatric clinic. All subjects, aged 20 through 73 years, met the DSM-IV criteria for schizophrenia. These patients had no history of chronic medical illnesses. Other exclusion criteria included patients who were pregnant, suicidal, active substance abusers, and those who in the investigators' opinion lacked capacity to give consent. This study was approved by the Subcommittee for the Protection of Human Subjects at the University of Miami. Informed consent was obtained from each subject. This study was conducted from January 2001 to January 2002.

Participants were seen after a 12-hour fast. All historical information was obtained by interview and from chart review. Sitting blood pressure was measured twice, and the mean reading was used. WC taken at the level of the umbilicus was measured twice, and the mean measurement was used. Fasting blood glucose and lipids were obtained. Height and weight were measured after the removal of shoes and with the patients wearing light clothing. BMI was calculated as weight (kg)/height (m²).

Cardiovascular risk factors shown in Table 1 are defined according to *The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults*¹¹: low-density lipoprotein (LDL)-cholesterol ≥ 160 mg/dL, high-density lipoprotein (HDL)-cholesterol < 35 mg/dL in men and < 45 mg/dL in women, systolic blood pressure (SBP) ≥ 140 mm Hg, diastolic blood pressure (DBP) ≥ 90 mm Hg, and fasting serum glucose ≥ 126

Table 2. Demographics and Characteristics of Schizophrenic Patients^a

Variable	Women (N = 30)	Men (N = 32)
Age, y	42.53 ± 12.76	41.53 ± 12.72
Weight, kg	86.35 ± 18.45	92.93 ± 20.45
Height, m	1.59 ± 0.11	1.72 ± 0.09
BMI, kg/m ²	34.34 ± 7.43	30.33 ± 4.55
WC, cm	102.79 ± 16.53	104.26 ± 13.34
Blood pressure, mm Hg		
Systolic	122.65 ± 17.40	126.14 ± 15.85
Diastolic	78.92 ± 10.11	82.13 ± 10.24
HDL-cholesterol, mg/dL	48.77 ± 12.84	41.60 ± 9.79
LDL-cholesterol, mg/dL	114.24 ± 35.83	116.79 ± 32.07
Triglycerides, mg/dL	194.10 ± 149.22	203.05 ± 105.31
Fasting glucose, mg/dL	113.63 ± 92.91	96.16 ± 23.01

^aData are mean ± SD.

Abbreviations: BMI = body mass index, HDL = high-density lipoprotein, LDL = low-density lipoprotein, WC = waist circumference.

mg/dL. Also, according to *The Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults*¹²: serum triglycerides level ≥ 150 mg/dL. Subjects with 1 or more of the above conditions were considered as having cardiovascular risk factors. Obesity was defined as a BMI value ≥ 30 kg/m² or WC > 102 cm in men and > 88 cm in women.⁶

All data were analyzed using the Statistical Program for Social Sciences 10.0 software program (SPSS Inc., Chicago, Ill.). Pearson's correlation coefficients were used to assess associations between adiposity measures and cardiovascular risk factors in women and men. The χ^2 test was used to compare the prevalence of the cardiovascular risk factors with WC and BMI as indices of obesity. The Fisher exact test was used if the expected cell size was < 5. Logistic regression analysis was utilized to allow for covariates. Triglyceride and glucose values were transformed to the natural logarithm to normalize skewed distribution for statistical testing; however, actual values are displayed. The p values were 2-sided, and the term *statistically significant* implies a p value < .05.

RESULTS

Sixty-two schizophrenic patients were enrolled in the study. The sample consisted of 30 women and 32 men, with a mean ± SD age of 41.98 ± 12.60 years. The ethnic background of the group included 40 Hispanics and 22 non-Hispanics. Seventy-nine percent (49/62) were white and 32% (20/62) were smokers. Sixty-nine percent (43/62) of the patients were taking atypical antipsychotics. The demographics and characteristics of the study patients are shown in Table 2.

Correlation coefficients for association among the adiposity measures revealed a strong and positive correlation between BMI and WC in both women (p < .01) and men

Table 3. Pearson's Correlation Coefficients for Association of Cardiovascular Risk Factors and Adiposity Measures

Variable	HDL	LDL	Log-Triglyceride	SBP	DBP	Log-Glucose
Women (N = 30)						
WC	-0.43*	0.27	0.16	0.37*	0.59**	0.25
BMI	-0.20	-0.21	0.28	0.35	0.31	0.32
Men (N = 32)						
WC	-0.12	0.35	0.44*	0.34	0.29	0.18
BMI	-0.06	0.39*	0.36*	0.12	0.16	0.20

*p < .05.

**p < .01.

Abbreviations: BMI = body mass index, DBP = diastolic blood pressure, HDL = high-density lipoprotein cholesterol, LDL = low-density lipoprotein cholesterol, SBP = systolic blood pressure, WC = waist circumference.

Table 4. Prevalence of Cardiovascular Risk Factors and Obesity in Patients With Schizophrenia (N = 62)

Variable	N	%
Dyslipidemia	36	58
Hypertension	28	45
Abnormal fasting glucose	9	15
Obesity		
WC	42	68
BMI	39	63

Abbreviations: BMI = body mass index, WC = waist circumference.

(p < .001). Table 3 displays Pearson's correlation coefficients for associations of cardiovascular risk factors and adipose measures in women and men. In women, there was an inverse association between HDL-cholesterol and WC, and a positive association was found between WC and both SBP and DBP. No association was found between BMI and cardiovascular risk factors in this group. In men, log-triglyceride positively correlated with both WC and BMI, and a significant association was also found between LDL-cholesterol and BMI.

The prevalence of cardiovascular risk factors and obesity defined by BMI and WC in patients with schizophrenia is shown in Table 4. Tables 5 and 6 show the prevalence of cardiovascular risk factors according to BMI and WC, respectively, in this population. In schizophrenic patients, an increased WC was significantly associated with all 3 risk factors: dyslipidemia, hypertension, and abnormal serum glucose, whereas an increased BMI was only significantly associated with dyslipidemia. In logistic regression analysis, after controlling for age, gender, race, ethnicity, smoking, and BMI, an increased WC remained significantly associated with dyslipidemia (odds ratio = 2.08, 95% CI = 1.01 to 1.15, p < .05) and hypertension (odds ratio = 2.05, 95% CI = 1.02 to 1.17, p < .05).

DISCUSSION

In this pilot study, data suggest that WC is a stronger predictor of cardiovascular risk factors than BMI in pa-

Table 5. Prevalence of Cardiovascular Risk Factors in Patients With Schizophrenia According to Body Mass Index (BMI)^a

Variable	Abnormal BMI (N = 39)	Normal BMI (N = 23)	p
Dyslipidemia	26 (67)	8 (35)	< .05
Normal lipids	13 (33)	15 (65)	
Abnormal glucose	8 (20)	1 (4)	.08
Normal glucose	31 (80)	22 (96)	
Hypertension	19 (49)	9 (39)	.46
Normal blood pressure	20 (51)	14 (61)	

^aData are N (%).

Table 6. Prevalence of Cardiovascular Risk Factors in Patients With Schizophrenia According to Waist Circumference (WC)^a

Variable	Abnormal WC (N = 42)	Normal WC (N = 20)	p
Dyslipidemia	28 (67)	6 (30)	< .01
Normal lipids	14 (33)	14 (70)	
Abnormal glucose	9 (21)	0 (0)	< .05
Normal glucose	33 (79)	20 (100)	
Hypertension	23 (55)	5 (25)	< .05
Normal blood pressure	19 (45)	15 (75)	

^aData are N (%).

tients with schizophrenia. Our findings in schizophrenic patients are comparable to those reported in the general population.^{13,14} Waist circumference, a surrogate measurement for intra-abdominal fat, has been associated with metabolic disturbances including insulin resistance.⁷ Insulin resistance is believed to be the underlying reason for the association between WC and cardiovascular risk factors.¹³

A strong correlation was found between BMI and WC, as has been shown in other studies. This points to WC as an indicator of both intra-abdominal fat and total body fat.¹⁵ The prevalence of obesity in patients with schizophrenia was high at 63% when defined by an abnormal BMI and 68% when defined by an abnormal WC. Patients with abnormal BMI or WC were found to have a higher prevalence of dyslipidemia, hypertension, and abnormal serum glucose than non-obese patients. A statistically significant association was found between increased WC and all 3 of the cardiovascular risk factors. In schizophrenic patients with increased BMI, a significant association was only found with dyslipidemia; but in logistic regression when age and WC were added as covariates, the association was no longer significant. The most common cardiovascular risk factor found in schizophrenic patients was dyslipidemia followed by hypertension. Notably, dyslipidemia and hypertension were the 2 measures that significantly correlated with an abnormal WC in logistic regressions. Elevated fasting glucose was the least frequent abnormality. Logistic regression revealed that the association between increased WC and both dyslipidemia and

hypertension was independent of BMI. Thus, WC, a measure of both general and central obesity, is a better anthropometric measure in patients with schizophrenia than BMI, which reflects only general obesity.

We propose the measurement of WC as a screening tool for cardiovascular risk factors. The standard weight and height measurements reflect body mass or obesity, whereas cardiovascular risk lies in the distribution of fat or namely central obesity. Waist circumference measurement provides an opportunity for primary prevention of coronary heart disease and diabetes mellitus in patients with schizophrenia. This screening tool is an objective measure that is easily obtained in any health care setting. Patients with increased WC (> 102 cm in men and > 88 cm in women) should be encouraged to lose weight and increase their physical activity.

Limitations in our study include lack of a control group and the small sample size. The power to identify further relationship between obesity measures and cardiovascular risk factors is reduced due to the small sample size. Another limitation is the cross-sectional relationship between predictors and cardiovascular risk factors used here. Further longitudinal studies are needed to corroborate our findings.

REFERENCES

1. Ganguli R. Weight gain associated with antipsychotic drugs. *J Clin Psychiatry* 1999;60(suppl 21):20–24
2. Allison DB, Mentore JL, Heo M, et al. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry* 1999;156:1686–1696
3. Wirshing DA, Wirshing WC, Kysar L, et al. Novel antipsychotics: comparison of weight gain liabilities. *J Clin Psychiatry* 1999;60:358–363
4. Wetterling T, Mubigbrodt HE. Weight gain: side effect of atypical neuroleptics? *J Clin Psychopharmacol* 1999;19:316–321
5. Kuczmarski RJ, Flegal KM, Troiano RP. Varying body mass index cutoff points to describe overweight prevalence among US adults: NHANES III (1988–1994). *Obes Res* 1997;5:542–548
6. Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Executive summary of the clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Arch Intern Med* 1998;158:1855–1867
7. Larsson B, Svarsdadd K, Welin L, et al. Abdominal adipose tissue distribution, obesity, and risk of cardiovascular disease and death: 13 year follow up of participants in the study of men born in 1913. *Br Med J (Clin Res Ed)* 1984;288:1401–1404
8. Lapidus L, Bengtsson C, Larsson B, et al. Distribution of adipose tissue and risk of cardiovascular disease and death: a 12 year follow up of participants in the population study of women in Gothenburg, Sweden. *Br Med J (Clin Res Ed)* 1984;289:1257–1261
9. Bouchard C, Bray GA, Hubbard VS. Basic and clinical aspects of regional fat distribution. *Am J Clin Nutr* 1990;52:946–950
10. American Diabetes Association, American Psychiatric Association, American Association of Clinical Endocrinologists, et al. Consensus development conference on antipsychotic drugs and obesity and diabetes. *J Clin Psychiatry* 2004;65:267–272
11. The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. US Dept Health Human Services publication 00-4084. Bethesda, Md: National Institutes of Health; 2000
12. Expert Panel on Detection and Treatment of High Blood Cholesterol in Adults. Executive summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486–2497
13. Schmidt MI, Watson RL, Duncan BB, et al. Clustering of dyslipidemia, hyperuricemia, diabetes, and hypertension and its association with fasting insulin and central and overall obesity in a general population. *Atherosclerosis Risk in Communities Study Investigators. Metabolism* 1996;45:699–706
14. Zhu SK, Wang Z, Heshka S, et al. Waist circumference and obesity-associated risk factors among whites in the Third National Health and Nutrition Examination Survey: clinical action thresholds. *Am J Clin Nutr* 2002;76:743–749
15. Han TS, Van Leer EM, Seidell JC, et al. Waist circumference action levels in the identification of cardiovascular risk factors: prevalence study in a random sample. *BMJ* 1995;31:1401–1405

For the CME Posttest for this article, see pages 119–120.

For more CME activities, visit

www.psychiatrist.com/pcc/cme

INSTRUCTIONS AND POSTTEST

Participants may receive up to 1 Category 1 credit toward the American Medical Association Physician's Recognition Award by reading the CME article and correctly answering at least 70% of the questions in the Posttest that follows:

Go to www.psychiatrist.com/cmehome to take this Posttest online and earn credit immediately.

Or

1. Read each question carefully and circle the answer on the Registration Form.
2. Type or print the registration information in the spaces provided and complete the evaluation.
3. Send the Registration Form to the address or fax number listed on the Registration Form.

All replies and results are confidential. Answer sheets, once graded, will not be returned. Unanswered questions will be considered incorrect and so scored. The CME Institute of Physicians Postgraduate Press, Inc. will keep only a record of participation, which indicates the completion of the activity and the designated number of Category 1 credits that have been awarded. Correct answers to the Posttest will be made available to the participants of this activity upon request after the submission deadline.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Answer to Pretest: 1. b

1. All of the following phrases describe body mass index (BMI) *except*:
 - a. Defined as weight (kg) divided by height (m²)
 - b. Is the accepted measure to assess obesity
 - c. Indicates total body fat
 - d. Indicates regional fat distribution
2. Recent consensus guidelines for patients with schizophrenia recommend that clinicians measure _____ to monitor cardiovascular risk factors.
 - a. Either BMI or waist-hip ratio (WHR)
 - b. Both BMI and WHR
 - c. Both BMI and waist circumference (WC)
 - d. Either WHR or WC
3. In this study, an increased WC was significantly associated with _____, after controlling for age, gender, race, ethnicity, smoking, and BMI.
 - a. Dyslipidemia and hypertension
 - b. Abnormal serum glucose and dyslipidemia
 - c. Hypertension and abnormal serum glucose
 - d. There was no significant association
4. In this study, an increased BMI was significantly associated with _____, until age and WC were added as covariates.
 - a. Dyslipidemia
 - b. Abnormal serum glucose and dyslipidemia
 - c. Hypertension
 - d. There was no significant association
5. WC measurement provides an opportunity for primary prevention of coronary heart disease and diabetes mellitus in patients with schizophrenia.
 - a. True
 - b. False

REGISTRATION FORM

Circle the one correct answer for each question.

1. a b c d 4. a b c d
2. a b c d 5. a b
3. a b c d

Print or type

Name _____

Last 4 digits of Social Security Number _____
(Required to issue CME credit)

Birth Date (mm, dd, yy) _____
(Required to issue CME credit)

Degree _____ Specialty _____

Affiliation _____

Address _____

City, State, Zip _____

Phone () _____

Fax () _____

E-mail _____

☐ Hospital ☐ Private Practice ☐ Resident ☐ Intern

Deadline for submission

For a credit certificate to be issued, please complete this Registration Form no later than June 30, 2007. Online submissions will receive credit certificates immediately. Faxed or mailed submissions will receive credit certificates within 6 to 8 weeks.

Keep a copy for your files

Retain a copy of your answers and compare them with the correct answers, which will be published after the submission deadline.

Payment

If you complete the test online, no payment is necessary. A \$10 payment must accompany this form. You may pay by check, money order, or credit card (Visa or MasterCard). Make check or money order payable to Physicians Postgraduate Press, Inc. If paying by credit card, please provide the information below.

Check one: ☐ Visa ☐ MasterCard

Card number _____

Expiration date _____

Your signature _____

Please evaluate the effectiveness of this CME activity by answering the following questions.

1. Was the educational content relevant to the stated educational objective? ☐ Yes ☐ No
2. Did this activity provide information that is useful in your clinical practice? ☐ Yes ☐ No
3. Was the format of this activity appropriate for the content being presented? ☐ Yes ☐ No
4. Did the method of presentation hold your interest and make the material easy to understand? ☐ Yes ☐ No
5. Achievement of educational objective:
 - A. Enabled me to select an appropriate body fat screening tool for predicting cardiovascular risk in patients with schizophrenia. ☐ Yes ☐ No
6. Did this CME activity provide a balanced, scientifically rigorous presentation of therapeutic options related to the topic, without commercial bias? ☐ Yes ☐ No
7. Does the information you received from this CME activity confirm the way you presently manage your patients? ☐ Yes ☐ No
8. Does the information you received from this CME activity change the way you will manage your patients in the future? ☐ Yes ☐ No
9. If you answered yes, what change(s) do you intend to make in your practice?

10. Please offer comments and/or suggested topics for future CME activities.

11. How much time did you spend completing this CME activity?

12. What is your preferred format for CME activities? Circle one.
 - a. Print media (eg, journals, supplements, and newsletters)
 - b. Internet text
 - c. Internet multimedia
 - d. Audio CD
 - e. Live meeting

TEAR OUT AND SEND THIS PAGE, ALONG WITH YOUR PAYMENT, TO:
CME INSTITUTE • PHYSICIANS POSTGRADUATE PRESS, INC. • P.O. BOX 752870 • MEMPHIS, TN 38175-2870

IF YOU ARE PAYING BY CREDIT CARD, YOU MAY FAX THIS PAGE TO: CME INSTITUTE AT 901-751-3444

QUESTIONS? CALL 1-800-489-1001 EXT. 8
WWW.PSYCHIATRIST.COM