Ku80 is required for immunoglobulin isotype switching

Rafael Casellas¹, Andre Nussenzweig², Robert Wuerffel³, Roberta Pelanda^{4,5}, Amy Reichlin¹, Heikyung Suh^{1,6}, Xiao-Feng Qin¹, Eva Besmer^{1,6}, Amy Kenter³, Klaus Rajewsky⁴ and Michel C.Nussenzweig^{1,6,7}

²Memorial Sloan-Kettering Institute, ³Department of Microbiology and Immunology, University of Illinois College of Medicine, IL 60612, ¹Laboratory of Molecular Immunology, The Rockefeller University, ⁶Howard Hughes Medical Institute, 1230 York Avenue, NY 10021, USA and ⁴Institute for Genetics, University of Cologne, D-50931 Weyertal, Germany

⁵Present address: Max Plank Institute for Immunobiology, D-79108 Freiburg, Germany

⁷Corresponding author
e-mail: nussen@rockvax.rockefeller.edu

Isotype switching is the DNA recombination mechanism by which antibody genes diversify immunoglobulin effector functions. In contrast to V(D)J recombination, which is mediated by RAG1, RAG2 and DNA double-stranded break (DSB) repair proteins, little is known about the mechanism of switching. We have investigated the role of DNA DSB repair in switch recombination in mice that are unable to repair DSBs due to a deficiency in Ku80 (Ku80^{-/-}). B-cell development is arrested at the pro-B cell stage in Ku80^{-/-} mice because of abnormalities in V(D)J recombination, and there are no mature B cells. To reconstitute the B-cell compartment in Ku80^{-/-} mice, pre-rearranged V_{B1-8}DJ_{H2} (μⁱ) and V₃₋₈₃J_{K2} (κⁱ) genes were introduced into the Ku80^{-/-} background (Ku80^{-/-}μ^{i/+}κ^{i/+}). Ku80^{-/-}μ^{i/+}κ^{i/+} mice develop mature mIgM⁺ B cells that respond normally to lipopolysaccharide (LPS) or LPS plus interleukin-4 (IL-4) by producing specific germline Ig constant region transcripts and by forming switch region-specific DSBs. However, Ku80^{-/-}μ^{i/+}κ^{i/+} B cells are unable to produce immunoglobulins of secondary isotypes, and fail to complete switch recombination. Thus, Ku80 is essential for overlap recombination *in vivo*, suggesting a significant overlap between the molecular machinery that mediates DNA DSB repair, V(D)J recombination and isotype switching.
Keywords: class switch recombination/DNA repair/immunoglobulin genes/Ku80

Introduction

During an immune response, B lymphocytes maintain their antigen-binding specificity but can change the antibody constant region subclass they produce by a DNA recombination process known as class switching (reviewed by Lorenz and Radbruch, 1996; Stavnezer, 1996). Switching occurs between highly repetitive DNA sequences, known

as switch regions, which are located 5' of the μ, γ, α and ε constant region (C_H) genes (Shimizu *et al.*, 1982). Switch recombination is preceded by DNA demethylation, increased deoxyribonuclease I hypersensitivity and germ-line transcription of the implicated C_H genes (Lorenz and Radbruch, 1996; Stavnezer, 1996).

Like V(D)J recombination (Tonegawa, 1983), switching involves DNA deletion (Honjo and Kataoka, 1978; Coleclough *et al.*, 1980; Cory *et al.*, 1980; Davis *et al.*, 1980a; Rabbitts, 1980; Sakano *et al.*, 1980; Yaoita and Honjo, 1980) by a mechanism whereby intervening sequences are excised in the form of circular DNA (Iwasato *et al.*, 1990; Matsuoka *et al.*, 1990; von Schwedler *et al.*, 1990). In addition, switching resembles V(D)J recombination in that a double-stranded break (DSB) intermediate appears to be part of the switch reaction (Wuerffel *et al.*, 1997). However, isotype switching is a unique process which differs from V(D)J recombination in several respects. V(D)J recombination is characterized by site-specific recombination between short well-defined recombination signal sequences (RSSs) (Tonegawa, 1983). In contrast, the precise signals that mediate switching are unknown, and switch regions range in size from 2.5 kb for S_ε to 10 kb for S_{γ1}. Unlike the RSSs which are shared by all V, D and J gene segments, the switch regions are composed of tandem nucleotide arrays that vary both in sequence and in length (Davis *et al.*, 1980b; Dunnick *et al.*, 1980; Sakano *et al.*, 1980; Kataoka *et al.*, 1981; Nikaido *et al.*, 1981, 1982; Obata *et al.*, 1981). Furthermore, RAG1 and RAG2, the enzymes that recognize the RSSs and activate V(D)J recombination, are not required for switching (Lansford *et al.*, 1998).

Mechanistically, V(D)J recombination involves a transesterification reaction that results in blunt-ended DNA DSBs and hairpin intermediates (van Gent *et al.*, 1996). Resolution of the broken DNA ends requires several non-lymphoid-specific proteins that were first defined by transfection and genetic studies in mutant cell lines and mice. These experiments showed that V(D)J recombination required proteins implicated in DNA DSB repair, including the catalytic subunit of the DNA-dependent protein kinase (DNA-PK_{cs}), Ku and XRCC4 (Bosma *et al.*, 1983; Schuler *et al.*, 1986; Hendrickson *et al.*, 1991; Pergola *et al.*, 1993; Taccioli *et al.*, 1993, 1994; Rathmell and Chu, 1994; Blunt *et al.*, 1995; Kirchgesner *et al.*, 1995; Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996).

Ku is a heterodimer composed of Ku70 and Ku80 subunits that binds to DNA DSBs, nicks, gaps and hairpins in a sequence-independent manner (Mimori *et al.*, 1986; Morozov *et al.*, 1994). DNA-bound Ku recruits and activates DNA-PK_{cs}, which is then thought to phosphorylate a series of factors which are themselves implicated in mediating DNA DSB repair (Gottlieb and Jackson, 1993; Jeggo *et al.*, 1995). The absence of either DNA-PK_{cs},

Ku80 or Ku70 results in a deficiency in both DNA DSB repair and V(D)J recombination (Bosma *et al.*, 1983; Schuler *et al.*, 1986; Hendrickson *et al.*, 1991; Pergola *et al.*, 1993; Taccioli *et al.*, 1993, 1994; Rathmell and Chu, 1994; Blunt *et al.*, 1995; Kirchgessner *et al.*, 1995; Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996; Ouyang *et al.*, 1997). However, the absence of Ku does not appear to affect DNA repair mediated by homologous recombination, which remains intact both in Ku-deficient yeast and mammalian cells (Boulton and Jackson, 1996a,b; Liang and Jasin, 1996; Liang *et al.*, 1996; Milne *et al.*, 1996; Siede *et al.*, 1996).

In contrast to V(D)J recombination, the role of DNA DSB repair in switch recombination in mature B cells has not been evaluated. Developing B cells in mice that carry mutations in DSB repair genes fail to complete V(D)J recombination, and lymphocyte development is arrested at early precursor stages (Bosma *et al.*, 1983; Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996; Ouyang *et al.*, 1997). Therefore, the cell type that normally undergoes class switching is absent, and this process cannot be assessed. Recombination between switch regions has been studied in DNA-PK_{cs} (SCID) mutant pro-B cells deprived of interleukin-7 (IL-7) (Rolink *et al.*, 1996). In the absence of IL-7, pro-B cells undergo programmed cell death and differentiate into pre-B cells (Grawunder *et al.*, 1993). When cell death is delayed in the IL-7-deprived pro-B cells by overexpression of Bcl-2, they undergo DNA recombination between Ig switch regions in a reaction that is independent of the assembly of a functional Ig transcription unit (Rolink *et al.*, 1996). Recombination between switch regions in pro-B cells *in vitro* requires DNA-PK_{cs}, but the requirements for this reaction may or may not be the same as authentic switching in mature B cells (Rolink *et al.*, 1996).

Here we report on switch recombination in mature Ku80^{-/-}μ^{i/+}κ^{i/+} B cells isolated from mice that carry pre-rearranged immunoglobulin heavy [V_{B1-8}DJ_H2 (Sonoda *et al.*, 1997)] and light [V₃₋₈₃J_K2 (Pelanda *et al.*, 1996)] chain genes. We find that B cells that are deficient in Ku80 are unable to complete switch recombination. These data implicate the DNA DSB repair machinery in class switch recombination.

Results

V_{B1-8}DJ_H2 and V₃₋₈₃J_K2 replacements reconstitute B-cell development in Ku80^{-/-} mice

Ku80^{-/-} mice are unable to repair DNA DSBs and show a phenotype that includes proportional dwarfism, and a profound disruption in both T- and B-lymphocyte development (Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996). In the absence of Ku80, developing lymphocytes cannot repair the DNA breaks produced by RAG1 and RAG2 during V(D)J recombination. The result of the DSB repair deficiency is that both T- and B-cell development are arrested at early precursor stages in Ku80^{-/-} mice (Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996).

To determine whether the absence of B cells in Ku80^{-/-} mice was due solely to impaired V(D)J recombination, we introduced pre-rearranged targeted Igμ (Sonoda *et al.*, 1997) and Igκ (Pelanda *et al.*, 1996) genes into the Ku80^{-/-} background (Figure 1). The mice resulting from these

Fig. 1. Map of the wild-type (+) Igκ and Igμ loci and their predicted structure after insertion (i) of the rearranged constructs. The 3' κ and H enhancers (E) are depicted as open circles, and the constant (C) κ region as open squares. Variable, diversity and joining regions are represented as black, open and gray bars respectively.

Fig. 2. Reconstitution of B-cell development in Ku80^{-/-} mice. Bone marrow and spleen samples from 6- to 10-week-old wild-type, Ku80^{+/-} and Ku80^{-/-}μ^{i/+}κ^{i/+} mice analyzed for B-cell maturation. Cell percentages were calculated from total gated populations. Bone marrow cells were stained with anti-B220 and anti-CD43, and splenocytes with anti-B220 and anti-IgM antibodies.

crosses are referred to as Ku80^{-/-}μ^{i/+}κ^{i/+} mice. Like Ku80^{-/-} controls, Ku80^{-/-}μ^{i/+}κ^{i/+} mice were proportional dwarfs and had no mature T cells (not shown). However, B-cell development in the bone marrow of Ku80^{-/-}μ^{i/+}κ^{i/+} mice differed from that of Ku80^{-/-} mice in that B cells progressed beyond the pro-B cell stage (Figure 2). FACS analysis of Ku80^{-/-}μ^{i/+}κ^{i/+} bone marrow showed that in the presence of pre-rearranged μ and κ Ig genes, Ku80-deficient B cells progressed to the B220⁺CD43⁻ pre-B cell stage and developed into immature and mature B220⁺IgM⁺ B cells (Figure 2). Mature B cells expressing surface IgM were also found in peripheral lymphoid organs such as spleen, but in 6- to 8-week-old mice the total number of B cells in Ku80^{-/-}μ^{i/+}κ^{i/+} was only 15–25% (*n* = 10) of that found in wild-type and Ku80^{+/-}μ^{i/+}κ^{i/+} littermate controls. This low number of mature B cells in the periphery of the reconstituted mice may be secondary to the absence of T cells and T Ku80^{-/-}μ^{i/+}κ^{i/+} B cell clonal expansion. Alternatively, the relative B

Fig. 3. IgG heavy chain (50 kDa) expression in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ mice and controls. Two ml of serum from wild-type (+/+) and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ (-/-) mice was analyzed by PAGE, and blotting with goat anti-mouse IgG visualized with alkaline phosphatase.

lymphopenia could be due to the documented inability of $Ku80^{-/-}$ cells to repair DNA DSBs incurred during normal proliferative responses (Nussenzweig *et al.*, 1996).

We conclude that pro-B cell arrest in $Ku80^{-/-}$ mice is a function of impaired resolution of DNA breaks resulting from V(D)J recombination and that Ku80 is not essential for other aspects of antigen-independent B cell development.

Ku80^{-/-} B cells are deficient in switch recombination

Consistent with the lower than normal number of peripheral B cells in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ mice, the level of circulating IgM was 62% of that found in wild-type mice as measured by an IgM-specific enzyme-linked immunosorbent assay (ELISA) (1.1 mg/ml in wild-type versus 0.68 in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ $n = 5$ mice of each type). In contrast to IgM, secondary Ig isotypes, which are normally found in the serum of un-immunized animals in the mg/ml range, were not detectable in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ mice. Western blotting with specific goat anti-mouse IgG showed no IgG heavy chains in the serum of the reconstituted mice (Figure 3).

The absence of secondary Ig isotypes in the serum of $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ mice could be due to a deficiency in switch recombination or might be a consequence of the absence of T cells in these mice. To determine whether the absence of secondary antibody isotypes in serum was due to a cell-autonomous defect, we isolated B cells and stimulated them with either lipopolysaccharide (LPS), or the combination of LPS plus IL-4 to activate switch recombination in cell culture.

LPS induces murine B cells to switch from μ to $\gamma 3$ and $\gamma 2b$, whereas LPS plus IL-4 activates switching to $\gamma 1$ and, to a lower extent, ϵ . In all cases, switch recombination is preceded by sterile transcription of the switch-targeted C_H genes (see Figure 4A). To determine whether $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells could respond to either LPS or LPS plus IL-4 by activating switch transcription, we measured germline sterile mRNAs by a semi-quantitative RT-PCR assay using $I\gamma 3$ -, $I\gamma 1$ - (Snapper *et al.*, 1996) and $I\gamma 2b$ -specific

Fig. 4. Ig germline and mature switch transcription. (A) Schematic representation of the $I\gamma 1$ locus before and after switching. The I exon (I), switch (S) and constant (C) regions for μ and $\gamma 1$ are shown. Their respective promoters are depicted as open circles. Switch recombination is preceded by the generation of sterile transcripts initiated at the I exons. Sterile transcripts were detected using I- and C-specific primers. (B) cDNA was prepared from wild-type (+/+), $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ (-/+ μ/κ) and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ (-/- μ/κ) unstimulated B cells (T_0) or B cells that were cultured with LPS or LPS plus IL-4 for 48 h (T_{48}). Germline ($I\gamma$) and mature (VDJ γ) transcripts for $\gamma 2b$, $\gamma 3$ and $\gamma 1$ were detected by RT-PCR using 30 cycles of amplification. Under the experimental conditions, the signal intensity was proportional to the amount of input cDNA. B cell-specific $Ig\beta$ mRNA was used to normalize each RT-PCR reaction.

primers. $Ig\beta$ mRNA was used as a B cell-specific mRNA loading control. $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ resembled control B cells in that $I\gamma 2b$, $I\gamma 3$ and $I\gamma 1$ germline transcripts were induced specifically by LPS and LPS plus IL-4 respectively, although the levels of sterile transcripts found in reconstituted mice were 10–30% lower than those found in wild-type mice, as assayed by phosphorimaging (Figure 4B, and data not shown). We conclude that $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells are competent to respond to signals that induce switch recombination *in vitro*.

To determine whether $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells can produce secondary antibodies in response to switch signals, we first measured cell surface expression of $\gamma 3$ and $\gamma 1$ after stimulation with LPS or LPS plus IL-4, respectively. B cells from $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ mice were compared with wild-type and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells after staining with isotype-specific antibodies (Figure 5). As expected, wild-type and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells expressed cell surface $\gamma 3$ and $\gamma 1$ following culture with LPS or LPS plus IL-4. In contrast, $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells showed no secondary isotype expression when cultured under the same conditions (Figure 5).

To enhance the sensitivity of the assay for secondary

Fig. 5. Cell surface expression of secondary Ig isotypes. Flow cytometry analysis of splenocytes from wild-type, $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$, and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ after LPS or LPS plus IL-4 stimulation. Cells were cultured for 3 days and stained with anti-B220 and anti-IgG3 (LPS) or anti-IgG1 (LPS-IL4). Percentages from total gated populations are shown.

isotypes and to determine whether $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells could produce mature secondary Ig mRNAs, we devised an isotype-specific PCR assay. $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ and $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ mice show a highly homogeneous B-cell compartment in which the great majority of the cells express the $V_{B1-8}DJ_{H2}$ replacement gene (Papavasiliou *et al.*, 1997). Primers specific for $V_{B1-8}DJ_{H2}$ combined with primers designed to hybridize $C\gamma3$, $C\gamma2b$ or $C\gamma1$ regions can therefore be used to measure mature $\gamma2b$, $\gamma3$ or $\gamma1$ mRNAs (see Figure 4A). As expected, mature $\gamma2b$, $\gamma3$ or $\gamma1$ mRNAs were abundant in the control $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ B cells stimulated with LPS or LPS and IL-4 (Figure 4B). In contrast, $V_{H}B1-8-C\gamma2b$ or $-C\gamma3$ mRNAs were detected at very low levels, and $V_{H}B1-8-C\gamma1$ was not detected in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells. Dilution analysis and quantitation by phosphorimaging showed that the levels of $V_{H}B1-8-C\gamma2b$ or $-C\gamma3$ were at least 10-fold lower than those found in $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ B cells.

When B cells are cultured with appropriate mitogens and cytokines, DNA DSBs are induced in the targeted switch regions (Wuerffel *et al.*, 1997). These breaks have been determined to be specific by several criteria including: (i) sequence; (ii) absence in T cells; (iii) absence in mitogen-stimulated T cells; and (iv) absence in resting B cells (Wuerffel *et al.*, 1997). To determine whether these DNA breaks were present in $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells, we made use of this $\gamma3$ switch region-specific DNA DSB assay (Wuerffel *et al.*, 1997) (Figure 6A). Figure 6B shows that DNA DSBs are generated in $\gamma3$ switch regions from $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ B cells and $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ controls upon activation with LPS for 4 h. These $\gamma3$ switch-specific breaks were not detected in unstimulated B cells or T cells stimulated with phytohemagglutinin (PHA). Further, B cells stimulated with anti- δ -dextran alone, which induces B-cell proliferation but little switch recombination, show only low levels of the $\gamma3$ switch-specific breaks (Figure 6B). In contrast, B cells treated with the combination of anti- δ -dextran plus IL-5, which induces high levels of $\gamma3$ switching, show high levels of $\gamma3$ switch-specific breaks. Thus, the $\gamma3$ switch breaks are seen in B cells stimulated

Fig. 6. $\gamma3$ switch breaks in stimulated B cells. (A) Schematic diagram of the DSB assay. A linker is ligated to 5'-phosphorylated blunt DNA ends in genomic DNA. DNA tagged with the linker is then PCR amplified for 15 cycles using the linker primer (L1) and a locus-specific primer (AP) (Wuerffel *et al.*, 1997). A second round of 15 cycle PCR amplification using AP and a nested linker primer (XL1) is then performed. The amplified products are then radioactively labeled and the precise positions of the DSBs are determined by denaturing PAGE. (B) $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ ($\mu/\kappa^{-/-}$) and control $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ ($\mu/\kappa^{+/-}$) splenocytes were stimulated with LPS for the indicated times and assayed for DNA breaks (Wuerffel *et al.*, 1997). Switch breaks are 104 nucleotides long and are found in live B cells undergoing switch recombination (Wuerffel *et al.*, 1997). As control, wild-type ($wt^{+/+}$) B cells were stimulated for 41 h with anti- δ -dextran (D) antibodies, and anti- δ -dextran plus IL-5 (D+IL5). L = DNA ladder.

with specific inducers of $\gamma3$ switching, and the breaks are seen in the presence and absence of Ku80 (Figure 6).

To determine whether the deficiency in switch transcription and protein production was due to a failure to complete switch recombination at the DNA level, we assayed for μ - $\gamma1$ switched DNA directly using a previously described $\gamma1$ digestion-circularization PCR (DC-PCR) assay. This assay, which detects recombination between μ and $\gamma1$ switch regions, was modified to increase sensitivity by including a second round of amplification with nested primers (Chu *et al.*, 1992) (see Figure 7A). We used the non-rearranging acetylcholine receptor (ACh) gene as a positive control for our digestion and ligation reactions (Chu *et al.*, 1992). All of the DNA samples were positive in the ACh DC-PCR reaction. In addition, μ - $\gamma1$ rearrangement was present in DNA from both wild-type and $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ control B cells stimulated with LPS and IL-4. In contrast, $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ showed a complete absence of μ - $\gamma1$ DNA recombination (Figure 7B). We conclude that there is no DNA recombination between μ and $\gamma1$ switch regions in $Ku80$ -deficient B cells stimulated with LPS and IL-4.

Discussion

Reconstitution of $Ku80$ -deficient mice

As previously reported, $Ku80^{-/-}$ mice exhibit impaired V(D)J recombination and B-cell maturation is arrested at the $B220^{+}CD43^{+}$ stage (Nussenzweig *et al.*, 1996; Zhu *et al.*, 1996). We bypassed this early block in development by introducing heavy and light chain variable region gene

Fig. 7. Digestion-circularization-mediated PCR assay for switch recombination. (A) Strategy for the DC-PCR adapted from Chu *et al.* (1992). Schematic map (not to scale) of the non-rearranged B1-8VDJ locus depicting the μ and $\gamma 1$ switching (S) and constant (C) regions. *Eco*RI digestion, ligation and PCR amplification results in a product only after $S\mu$ - $S\gamma 1$ recombination. A control PCR product from the nAChR is found regardless of chromosomal rearrangement. R represents *Eco*RI sites. (B) Semi-quantitative nested DC-PCR was performed on DNA extracted from wild-type, $Ku80^{+/-}\mu^{i/+}\kappa^{i/+}$ and $Ku80^{-/-}\mu^{i/+}\kappa^{i/+}$ splenocytes that had been stimulated with LPS and IL-4 for 3 days. The absence of switch DNA rearrangement in $Ku80$ -deficient B cells was confirmed by overexposure or by using higher concentrations of genomic DNA (up to 100 ng per reaction, data not shown). The size of the amplified products is indicated.

replacements that create fully functional Ig loci in the appropriate genomic context (Pelanda *et al.*, 1996; Sonoda *et al.*, 1997). These gene replacements were sufficient to reconstitute the peripheral B-cell compartment, indicating that, in early B-cell development, $Ku80$ is required solely for processing of V(D)J recombination intermediates. This result is consistent with previous reconstitution experiments using randomly integrated Ig transgenes in mice deficient for other components of the V(D)J recombinase (Spanopoulou *et al.*, 1994; Young *et al.*, 1994; Chang *et al.*, 1995). However, it was not possible to study switch recombination in mice with randomly integrated transgenes since switching normally occurs *in cis* on any given chromosome (Wabl *et al.*, 1985). In contrast to these studies, targeted V_H gene replacements provide physiologic substrates in mature B cells to study isotype switching (Lansford *et al.*, 1998).

Switching: a homology-based or an end-joining recombination mechanism?

At least two distinct biochemical pathways mediate DNA DSB repair in *Saccharomyces cerevisiae* and higher eukaryotes (Kramer *et al.*, 1994; Moore and Haber, 1996). The gene products of the RAD52 epistasis group function in DNA repair by homologous recombination, a process

in which damaged chromosomes restore genetic integrity by physically pairing with a sister chromatid or homolog. $Ku80$ and $Ku70$ are part of a separate group of proteins that are essential for DSB repair by a mechanism that does not require DNA homology (Boulton and Jackson, 1996a,b; Liang *et al.*, 1996; Milne *et al.*, 1996; Siede *et al.*, 1996). Consequently, Ku -deficient yeast strains are unable to repair DSBs properly by end-joining (Boulton and Jackson, 1996a,b; Milne *et al.*, 1996; Siede *et al.*, 1996). In mammalian cells, $Ku80$ is similarly dispensable for repair by homologous recombination, but required for rejoining of endonuclease-induced DNA breaks (Liang *et al.*, 1996) and for V(D)J recombination (Taccioli *et al.*, 1993, 1994; Rathmell and Chu, 1994; Smider *et al.*, 1994).

Although isotype switching is known to proceed through looping out and deletion of DNA (Iwasato *et al.*, 1990; Matsuoka *et al.*, 1990; von Schwedler *et al.*, 1990), the molecular details of the reaction remain to be determined. The similarity between different switch regions has led to the suggestion that short DNA stretches of identity could align and participate actively in the recombination process (reviewed in Stavnezer, 1996). However, if switch recombination were to proceed through a homology-based DNA repair mechanism, it would not be expected to be disrupted in Ku -deficient mice. Indeed, the finding that Ku is required for switching suggests that switch recombination involves a non-homologous DNA DSB mechanism. This idea is also supported by the observation that the catalytic subunit of DNA-PK is required for recombination between switch regions in pro-B cells *in vitro* (Rolink *et al.*, 1996).

The conclusion that DSB repair is required for switch recombination is consistent with the recent finding of DNA DSB intermediates in switch regions undergoing recombination (Wuerffel *et al.*, 1997). In these experiments, ligation-mediated PCR was used to document switch-specific, blunt and 5'-phosphorylated DNA ends in switch regions targeted for recombination (Wuerffel *et al.*, 1997). These switch ends are thus similar to the signal end intermediates found in V(D)J recombination (Roth *et al.*, 1993; Schlissel *et al.*, 1993). The requirement for Ku in switching and the detection of blunt 5'-phosphorylated switch DNA ends suggest a model for switching that resembles V(D)J recombination. In this model, switch donor and acceptor DNA tandem repeats are first cleaved by a switch-specific endonuclease to produce DNA DSB (Wuerffel *et al.*, 1997). Cleavage may be either imprecise or followed by limited processing leading to a cluster of breakpoints within a subregion of the tandem repeat (Wuerffel *et al.*, 1997). DNA ends produced by the switch endonuclease would then be held together by Ku and possibly other proteins in a synaptic complex analogous to the one described for signal ends during the V(D)J recombination reaction (Sheehan and Lieber, 1993; Eastman *et al.*, 1996; Agrawal, 1997). It has been suggested that the role of Ku in such a complex is to stimulate DNA end joining (Ramsden and Gellert, 1998).

An interesting prediction of this model is that switch region tandem repeats would not function in the DNA recombination process *per se*, but could act as recognition sites for a switch-specific endonuclease, much as RSSs are targeted by RAG1 and RAG2 in V(D)J recombination. Alternatively, transcription of these tandem repeats could

create DNA structures that would be recognized by a switch-specific endonuclease.

In conclusion, Ku80 is essential for switch recombination in mature B cells. Further, reconstitution of the mature B cell compartment using targeted Ig genes in mice deficient in DNA repair genes such as Ku70, DNA-PK α and XRCC4 provides a general method for evaluating the role of these and other genes in switch recombination *in vivo*.

Materials and methods

Mice

Mice were bred and maintained under specific pathogen-free conditions. Screening for targeted genes was as previously described (Nussenzweig *et al.*, 1996; Pelanda *et al.*, 1996; Sonoda *et al.*, 1997).

Cell culture and FACSscan analysis

Bone marrow and spleen cells from 6- to 8-week-old mice were cultured for 3–4 days in complete RPMI medium [RPMI 1640 (Gibco-BRL) with 10% fetal calf serum (Sigma), 1% antibiotic-antimycotic (Gibco-BRL), 1% L-glutamine (Gibco-BRL), 2% minimal essential medium amino acids solution, 1% sodium pyruvate solution (Cellgro), 10 mM HEPES buffer (Gibco-BRL) and 53 μ M β -mercaptoethanol (Fisher Scientific)] with the addition of 25 μ g/ml LPS (*Escherichia coli* 0111:B4; Sigma) or LPS and IL-4 (50 U/ml; Gibco-BRL). Bone marrow or spleen cell suspensions were stained with phycoerythrin-labeled anti-B220, and fluorescein isothiocyanate-labeled anti-CD43 or anti-IgM antibodies respectively (Pharmingen). LPS or LPS-IL-4-stimulated spleen cells were stained with phycoerythrin-labeled anti-B220 and biotin-conjugated anti-IgG3 and IgG1 antibodies and developed with streptavidin-Cy-Chrome (Pharmingen). Stained samples were gated according to standard forward- and side-scatter values and analyzed on a Becton-Dickinson FACSscan fluorescence-activated cell sorter with CELLQuest software.

DC-PCR

To increase the sensitivity of the DC-PCR assay, we added a nested amplification to the basic protocol (Chu *et al.*, 1992). Following ligation, 5 ng of DNA were denatured at 94°C for 5 min followed by 30 cycles of amplification with the first primer set at (94°C for 15 s, 66°C 1.5 min, 72°C 1 min) and a final amplification at 72°C for 10 min. From each first cycle reaction, 2 μ l was amplified further for 30 cycles with a second set of primers in the presence of [α - ^{32} P]dCTP (4 μ Ci). The cycle conditions were 94°C for 15 s, 68°C 1.5 min, 72°C 1 min. PCR products were analyzed by 8% PAGE and the products visualized and quantitated with a phosphorimager. PCR primers were S μ -S γ 1 first set: S1, 5' GAGCAGCTACCAAGGATCAGGGA 3' and S2, 5' CTTCACG-CCACTGACTGACTGAG 3'; S μ -S γ 1 second set: S3, 5' GGAGACCA-ATAATCAAGGGAAG 3' and S4, 5' GAGACAGGGTCTCC-TGGGTAGG 3'. For the nicotinic acetyl choline receptor (nAChR), primers used for the first set were: A1, 5' GCAAACAGGGCTGGATG-AGGCTG 3' and A2, 5' GTCCCATCTTAGAACCCAGCG 3'. For the second set, the primers were: A3, 5' GGACTGCTGTGGGTTT-CACCCAG 3' and A4, 5' GCCTTGCTTGCTTAAGACCTGG 3'.

RT-PCR

Total RNA was reverse transcribed with Superscript II (Life Technologies). Then 5–100 ng of cDNA was amplified in the presence of [^{32} P]dCTP (4 μ Ci; DuPont) in a 25 μ l PCR reaction with 10 pmol of each primer. To detect germline sterile transcripts, the following primers were paired: (i) I γ 3, γ 3-5' CAAGTGGATCTGAACACA and γ 3-3' GGCTCCATAGTTCCATT (expected product 350 bp); (ii) I γ 2b, γ 2b-5' CCTGACACCAAGGTCACG and γ 2b-3' CGACCAGGCAA-GTGAGACTG (expected product 345 bp); (iii) I γ 1, γ 1-5' CAGCCT-GGTGTCAACTAG and γ 1-3' GCAAGGGATCCAGAGTTCCAG (expected product 341 bp); (iv) Ig β , Ig β -5' GATGACGGCAA-GGCTGGGATGGAGGAA and Ig β -3' CTCATTCCTGGCCTGGATGC (expected product 142 bp). Mature γ 3 and γ 2b transcripts were amplified by combining a V μ B1-8-specific primer V μ B1-8 CAAGGGCAA-GGCCAACTG with either C γ 3 CCACTGCTGCCTGAGCCATCTC (expected product 313 bp) or C γ 2b CAGGTGACGGTCTGACTTGG (expected product 414 bp). Mature γ 1 transcripts were amplified by combining a second V μ B1-8-specific primer V μ B1-8' CCAGCT-ACTGGATGCACTG 3' with C γ 1 GGACAGCTGGGAAGGTGTG 3'

(expected product 440 bp). All reactions were performed for 30 cycles. Amplified samples were analyzed by 8% PAGE and visualized by phosphorimaging.

Western blotting

An aliquot of 2 μ l of mouse serum was diluted in 100 μ l of phosphate-buffered saline (PBS) and incubated with 20 μ l of protein A-Sepharose beads (Pierce) for 1 h. The beads were washed three times with PBS and the bound proteins separated by 8% reducing PAGE before blotting and visualization with alkaline phosphatase-conjugated goat anti-mouse IgG (Pierce).

DSB assay

Splenocytes were activated in culture as previously described (Wuerffel *et al.*, 1997). Genomic DNA was prepared using the Cell Culture DNA Kit (Qiagen) or the Puregene DNA Isolation Kit (Gentra). The DSB assay for S γ 3 DNA (Wuerffel *et al.*, 1997) was performed with modifications. The partially double-stranded linker was ligated directly to 1.5 μ g of unmodified genomic DNA for 18 h at 16°C. PCR amplification was done in two rounds of 15 cycles each (1 min at 95°C/2 min at 67°C/3 min at 76°C). Primers S γ 3 AP and L1 (Wuerffel *et al.*, 1997) were used in the first round. Ten μ l of product was taken to program the second round of amplification using primers S γ 3 AP and XL1, a nested primer (5'-GTGACCCGGGAGATCTGAATTCCTCC-3') specific for a subset of ligated broken fragments. Reaction products were radioactively labeled and analyzed by denaturing polyacrylamide gel electrophoresis. Broken fragments amplified by this method are three nucleotides shorter than the corresponding fragments of the ladder which are amplified using AP and L1 only.

Acknowledgements

The authors would like to thank members of the Nussenzweig laboratory for advice and suggestions. R.C. was supported by an NSF pre-doctoral fellowship. Supported by the Arthritis Foundation (A.N.), Council for Tobacco Research #3175 (A.K.), a Deutsche Forschungsgemeinschaft grant (K.R.), NIH grants (M.C.N.) and NIH Physician Scientist award (A.R.). M.C.N. is an Associate Investigator in the Howard Hughes Medical Institute.

References

- Agrawal, A. and Schatz, D. (1997) RAG1 and RAG2 form a stable poscleavage synaptic complex with DNA containing signal ends in V(D)J recombination. *Cell*, **89**, 43–53.
- Blunt, T. *et al.* (1995) Defective DNA-dependent protein kinase activity is linked to V(D)J recombination and DNA repair defects associated with the murine *scid* mutation. *Cell*, **80**, 813–823.
- Bosma, G.C., Custer, R.P. and Bosma, M. J. (1983) A severe combined immunodeficiency mutation in the mouse. *Nature*, **301**, 527–530.
- Boulton, S.J. and Jackson, S.P. (1996a) Identification of a *Saccharomyces cerevisiae* Ku80 homologue: roles in DNA double strand break rejoining and in telomeric maintenance. *Nucleic Acids Res.*, **24**, 4639–4648.
- Boulton, S.J. and Jackson, S.P. (1996b) *Saccharomyces cerevisiae* Ku70 potentiates illegitimate DNA double-strand break repair and serves as a barrier to error-prone DNA repair pathways. *EMBO J.*, **15**, 5093–5103.
- Chang, Y., Bosma, G.C. and Bosma, M.J. (1995) Development of B cells in *scid* mice with immunoglobulin transgenes: implications for the control of V(D)J recombination. *Immunity*, **2**, 607–616.
- Chu, C.C., Paul, W.E. and Max, E.E. (1992) Quantitation of immunoglobulin mu-gamma 1 heavy chain switch region recombination by a digestion-circularization polymerase chain reaction method. *Proc. Natl Acad. Sci. USA*, **89**, 6978–6982.
- Coleclough, C., Cooper, D. and Perry, R.P. (1980) Rearrangement of immunoglobulin heavy chain genes during B-lymphocyte development as revealed by studies of mouse plasmacytoma cells. *Proc. Natl Acad. Sci. USA*, **77**, 1422–1426.
- Cory, S., Jackson, J. and Adams, J.M. (1980) Deletions in the constant region locus can account for switches in immunoglobulin heavy chain expression. *Nature*, **285**, 450–456.
- Davis, M.M., Calame, K., Early, P.W., Livant, D.L., Joho, R., Weissman, I.L. and Hood, L. (1980a) An immunoglobulin heavy-chain gene is formed by at least two recombinational events. *Nature*, **283**, 733–739.

- Davis,M.M., Kim,S.K. and Hood,L.E. (1980b) DNA sequences mediating class switching in alpha-immunoglobulins. *Science*, **209**, 1360–1365.
- Dunnick,W., Rabbitts,T.H. and Milstein,C. (1980) An immunoglobulin deletion mutant with implications for the heavy-chain switch and RNA splicing. *Nature*, **286**, 669–675.
- Eastman,Q.M., Leu,T.M.L. and Schatz,D.G. (1996) Initiation of V(D)J recombination *in vitro* obeying 12/23 rule. *Nature*, **380**, 85–88.
- Gottlieb,T.M. and Jackson,S.P. (1993) The DNA-dependent protein kinase: requirement for DNA ends and association with Ku antigen. *Cell*, **72**, 131–142.
- Grawunder,U., Haasner,D., Melchers,F. and Rolink,A. (1993) Rearrangement and expression of kappa light chain genes can occur without mu heavy chain expression during differentiation of pre-B cells. *Int. Immunol.*, **5**, 1609–1618.
- Hendrickson,E.A., Qin,X.-Q., Bump,E.A., Schatz,D.G., Oettinger,M. and Weaver,D.T. (1991) A link between double-strand break-related repair and V(D)J recombination: the *scid* mutation. *Proc. Natl Acad. Sci. USA*, **88**, 4061–4065.
- Honjo,T. and Kataoka,T. (1978) Organization of immunoglobulin heavy chain genes and allelic deletion model. *Proc. Natl Acad. Sci. USA*, **75**, 2140–2144.
- Iwasato,T., Shimizu,A., Honjo,T. and Yamagishi,H. (1990) Circular DNA is excised by immunoglobulin class switch recombination. *Cell*, **62**, 143–149.
- Jeggo,P.A., Taccioli,G.E. and Jackson,S.P. (1995) Menage a trois: double strand break repair, V(D)J recombination and DNA-PK. *BioEssays*, **17**, 949–957.
- Kataoka,T., Miyata,T. and Honjo,T. (1981) Repetitive sequences in class-switch recombination regions of immunoglobulin heavy chain genes. *Cell*, **23**, 357–368.
- Kirchgesner,C.U., Patil,C.K., Evans,J.W., Cuomo,C.A., Fried,L.M., Carter,T., Oettinger,M.A. and Brown,J.M. (1995) DNA-dependent kinase (p350) as a candidate gene for the murine *scid* defect. *Science*, **267**, 1178–1183.
- Kramer,K.M., Brock,J.A., Bloom,K., Moore,J.K. and Haber,J.E. (1994) Two different types of double-strand breaks in *Saccharomyces cerevisiae* are repaired by similar RAD52-independent, nonhomologous recombination events. *Mol. Cell. Biol.*, **14**, 1293–1301.
- Lansford,R., Manis,J.P., Sonoda,E., Rajewsky,K. and Alt,F. (1998) Ig heavy chain class switching in rag-deficient mice. *Int. Immunol.*, **10**, 101–108.
- Liang,F. and Jasin,M. (1996) Ku-80 deficient cells exhibit excess degradation of extrachromosomal DNA. *J. Biol. Chem.*, **271**, 14405–14411.
- Liang,F., Romanienko,P.J., Weaver,D.T., Jeggo,P.A. and Jasin,M. (1996) Chromosomal double-strand break repair in Ku80-deficient cells. *Proc. Natl Acad. Sci. USA*, **93**, 8929–8933.
- Lorenz,M. and Radbruch,A. (1996) Developmental and molecular regulation of immunoglobulin class switch recombination. *Curr. Top. Microbiol. Immunol.*, **217**, 151–169.
- Matsuoka,M., Yoshida,K., Maeda,T., Usuda,S. and Sakano,H. (1990) Switch circular DNA formed in cytokine-treated mouse splenocytes: evidence for intramolecular DNA deletion in immunoglobulin class switching. *Cell*, **62**, 135–142.
- Milne,G.T., Jin,S., Shannon,K.B. and Weaver,D.T. (1996) Mutations in two Ku homologs define a DNA end-joining repair pathway in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.*, **16**, 4189–4198.
- Mimori,T., Hardin,J.A. and Seitz,J.A. (1986) Characterization of the DNA-binding protein antigen Ku recognized by autoantibodies from patients with rheumatic disorders. *J. Biol. Chem.*, **261**, 2274–2278.
- Moore,J.K. and Haber,J.E. (1996) Cell cycle and genetic requirements of two pathways of nonhomologous end-joining repair of double-strand breaks in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.*, **16**, 2164–2173.
- Morozov,V.E., Falzon,M., Anderson,W.C. and Kuff,E.L. (1994) DNA-dependent protein kinase is activated by nicks and larger single-stranded gaps. *J. Biol. Chem.*, **269**, 16684–16688.
- Nikaido,T., Nakai,S. and Honjo,T. (1981) Switch region of immunoglobulin Cmu gene is composed of simple tandem repetitive sequences. *Nature*, **292**, 845–848.
- Nikaido,T., Yamawaki-Kataoka,Y. and Honjo,T. (1982) Nucleotide sequences of switch regions of immunoglobulin C epsilon and C gamma genes and their comparison. *J. Biol. Chem.*, **257**, 7322–7329.
- Nussenzweig,A., Chen,C., da Costa Soares,V., Sanchez,M., Sokol,K., Nussenzweig,M.C. and Li,G.C. (1996) Requirement for Ku80 in growth and V(D)J recombination. *Nature*, **382**, 551–5.
- Obata,M., Kataoka,T., Nakai,S., Yamagishi,H., Takahashi,N., Yamawaki-Kataoka,Y., Nikaido,T., Shimizu,A. and Honjo,T. (1981) Structure of a rearranged gamma 1 chain gene and its implication to immunoglobulin class-switch mechanism. *Proc. Natl Acad. Sci. USA*, **78**, 2437–2441.
- Ouyang,H. *et al.* (1997) Ku70 is required for DNA repair but not for TCR gene recombination *in vivo*. *J. Exp. Med.*, **186**, 921–930.
- Papavasiliou,F., Casellas,R., Qin,X.F., Besmer,E., Suh,H., Pelanda,R., Nemazee,D., Rajewsky,K. and Nussenzweig,M.C. (1997) V(D)J recombination in mature B cells: a new mechanism for diversification of antibody responses. *Science*, **278**, 298–301.
- Pelanda,R., Schaal,S., Torres,R.M. and Rajewsky,K. (1996) A prematurely expressed Ig(kappa) transgene, but not V(kappa)J(kappa) gene segment targeted into the Ig(kappa) locus, can rescue B cell development in lambda5-deficient mice. *Immunity*, **5**, 229–239.
- Pergola,F., Zdzienicka,M.Z. and Lieber,M.R. (1993) V(D)J recombination in mammalian mutants defective in DNA double-strand break repair. *Mol. Cell. Biol.*, **13**, 3464–3471.
- Rabbitts,T., Forster,A., Dunnick,W. and Bentley,D.L. (1980) The role of gene deletion in the immunoglobulin heavy chain switch. *Nature*, **283**, 351–356.
- Ramsden,D.A. and Gellert,M. (1998) Ku protein stimulates DNA end joining by mammalian DNA ligases: a direct role for Ku in repair of DNA double-stranded breaks. *EMBO J.*, **17**, 609–614.
- Rathmell,W.K. and Chu,G. (1994) A DNA end-binding factor involved in double-strand break repair and V(D)J recombination. *Mol. Cell. Biol.*, **14**, 4741–4748.
- Rolink,A., Melchers,F. and Andersson,J. (1996) The SCID but not the RAG-2 gene product is required for S mu-S epsilon heavy chain class switching. *Immunity*, **5**, 319–330.
- Roth,D.B., Zhu,C. and Gellert,M. (1993) Characterization of broken DNA molecules associated with V(D)J recombination. *Proc. Natl Acad. Sci. USA*, **90**, 10788–10792.
- Sakano,H., Maki,R., Kurosawa,Y., Roeder,W. and Tonegawa,S. (1980) Two types of somatic recombination are necessary for the generation of complete immunoglobulin heavy-chain genes. *Nature*, **286**, 676–683.
- Schlissel,M., Constantinescu,U.A., Morrow,T., Baxter,M. and Peng,A. (1993) Double-strand signal sequence breaks in V(D)J recombination are blunt, 5'-phosphorylated, RAG-dependent, and cell cycle regulated. *Genes Dev.*, **7**, 2520–2532.
- Schuler,W., Weiler,I.J., Schuler,A., Phillips,R.A., Rosenberg,N., Mak,T.W., Kearney,J.F., Perry,R.P. and Bosma,M.J. (1986) Rearrangement of antigen receptor genes is defective in mice with severe combined immune deficiency. *Cell*, **46**, 963–972.
- Sheehan,K.M. and Lieber,M.R. (1993) V(D)J recombination: signal and coding joint resolution are uncoupled and depend on parallel synapsis of the sites. *Mol. Cell. Biol.*, **13**, 1363–1370.
- Shimizu,A., Takahashi,N., Yaoita,Y. and Honjo,T. (1982) Organization of the constant-region gene family of the mouse immunoglobulin heavy chain. *Cell*, **28**, 499–506.
- Siede,W., Friedl,A.A., Dianova,I., Eckardt-Schupp,F. and Friedberg,E.C. (1996) The *Saccharomyces cerevisiae* Ku autoantigen homologue affects radiosensitivity only in the absence of homologous recombination. *Genetics*, **142**, 91–102.
- Smider,V., Rathmell,W.K., Lieber,M.R. and Chu,G. (1994) Restoration of X-ray resistance and V(D)J recombination in mutant cells by Ku cDNA. *Science*, **266**, 288–91.
- Snapper,C.M., Zelazowski,P., Rosas,F.R., Kehry,M.R., Tian,M., Baltimore,D. and Sha,W.C. (1996) B cells from p50/NF- κ B knockout mice have selective defects in proliferation, differentiation, germ-line CH transcription, and Ig class switching. *J. Immunol.*, **156**, 183–191.
- Sonoda,E., Pewzner-Jung,Y., Schwes,S., Taki,S., Jung,S., Eilat,D. and Rajewsky,K. (1997) B cell development under the condition of allelic inclusion. *Immunity*, **6**, 225–233.
- Spanopoulou,E. *et al.* (1994) Functional immunoglobulin transgenes guide ordered B-cell differentiation in Rag-1-deficient mice. *Genes Dev.*, **8**, 1030–1042.
- Stavnezer,J. (1996) Immunoglobulin class switching. *Curr. Opin. Immunol.*, **8**, 199–205.
- Taccioli,G.E., Rathbun,G., Oltz,E., Stamato,T., Jeggo,P.A. and Alt,F.W. (1993) Impairment of V(D)J recombination in double-strand break repair mutants. *Science*, **260**, 207–210.
- Taccioli,G.E. *et al.* (1994) Ku80: product of the *XRCC5* gene and its role in DNA repair and V(D)J recombination. *Science*, **265**, 1442–1445.
- Tonegawa,S. (1983) Somatic generation of antibody diversity. *Nature*, **302**, 575–581.
- van Gent,D.C., Mizuuchi,K. and Gellert,M. (1996) Similarities between initiation of V(D)J recombination and retroviral integration. *Science*, **271**, 1592–1594.

- von Schwedler,U., Jack,H.M. and Wabl,M. (1990) Circular DNA is a product of the immunoglobulin class switch rearrangement. *Nature*, **345**, 452–456.
- Wabl,M., Meyer,J., Beck-Engeser,G., Tenkhoff,M. and Burrows,P.D. (1985) Critical test of a sister chromatid exchange model for the immunoglobulin heavy-chain class switch. *Nature*, **313**, 687–689.
- Wuerffel,R.A., Du,J., Thompson,R.J. and Kenter,A.L. (1997) Immunoglobulin S γ 3 DNA specific double stranded breaks are induced in mitogen activated B cells and are implicated in switch recombination. *J. Immunol.*, **159**, 4139–4144.
- Yaoita,Y. and Honjo,T. (1980) Deletion of immunoglobulin heavy chain genes from expressed allelic chromosome. *Nature*, **286**, 850–853.
- Young,F., Ardman,B., Shinkai,Y., Lansford,R., Blackwell,T.K., Mendelsohn,M., Rolink,A., Melchers,F. and Alt,F.W. (1994) Influence of immunoglobulin heavy- and light-chain expression on B-cell differentiation. *Genes Dev.*, **8**, 1043–1057.
- Zhu,C., Bogue,M.A., Lim,D.-S., Hasty,P. and Roth,D.B. (1996) Ku86-deficient mice exhibit severe combined immunodeficiency and defective processing of V(D)J recombination intermediates. *Cell*, **86**, 379–389.

*Received November 18, 1997; revised February 16, 1998;
accepted February 24, 1998*