The Biosynthesis of Prostaglandins in Thrombocytes

By J. CLAUSEN and K. C. SRIVASTAVA (*Department of Nutrition, Institute of Hygiene, University of Copenhagen and the Neurochemical Institute, 58 Radmandsgade-2200, Copenhagen N, Denmark*)

Prostaglandins are metabolic products of dihomog- γ -linoleate and arachidonate. Studies by Clausen *et al.* (1969) have shown that prostaglandins are able to inhibit the ADP-induced thrombocyte agglutination. Since the half-life of prostaglandins in the blood stream is only 2 min owing to enzymic inactivation in tissues such as lungs, the pathological significance of this inhibition in relation to the formation of thrombi in cardiovascular disorders can only be accepted if the prostaglandins are formed by enzymic systems of the thrombocytes. The present studies were designed to prove this theory.

Thrombocytes were isolated by selective spontaneous sedimentation from erythrocytes and leucocytes. Incubation of the platelets with [14 C]acetate in Krebs-Ringer bicarbonate solution in the presence of O_2 gave rise to significant incorporation of the label into the endogenous linoleate. The chain-elongating and desaturating enzymic systems gave rise to incorporation of the label into dihomog- γ -linoleate and arachidonate. T.l.c. on gel impregnated with $AgNO_3$ and t.l.c. of mercury adducts followed by assay of radioactivity revealed a significant desaturation activity of platelets.

Preparative separation and identification of prostaglandins from the platelets incubated as described above were investigated. Thus by means of silicic acid column chromatography and t.l.c. (J. Clausen & K. C. Srivastava, unpublished work) a significant amount of label was shown to be present in the prostaglandin E fraction when prostaglandin E was added to the thrombocyte reaction mixture before incubation. These data thus indicate the ability of the platelets to synthesize prostaglandin E and argue in favour of the theory that one of the factors of coronary thrombosis may be a relative linoleate deficiency leading to a lowering of prostaglandin E formation in thrombocytes.

These studies were supported by a grant from the Danish Heart Association.

Clausen, J., Kloeze, J., Houtmuller, U. M. T. & Vles, R. O. (1969) *Atheroscler. Res.* 9, 319

The Synthesis and Biological Role of Prostaglandins

By BENGT SAMUELSSON (*Department of Medical Chemistry, Royal Veterinary College, S-104 05 Stockholm, Sweden*)

Studies on the biosynthesis and the biological transformation of prostaglandins will be described. The latter studies have afforded the chemical background for quantitative work on the synthesis of prostaglandins. Recent experiments dealing with the endogenous synthesis of prostaglandins in relation to the physiological state of experimental animals have provided new information on the biological role of prostaglandins.

Fatty Acid Metabolism in Multiple Sclerosis

By R. H. S. THOMPSON (*Courtauld Institute of Biochemistry, The Middlesex Hospital Medical School, London W1P 5PR, U.K.*)

Studies of the changes in the fatty acid composition of the choline- and ethanolamine-containing phospholipids of the brain white matter taken from patients dying with multiple sclerosis will be reviewed. Changes in the linoleate content of the plasma lipids in patients suffering from this disease will also be described, together with work on the linoleate concentrations in erythrocyte and platelet phospholipids, indicating that there is growing evidence of a widespread abnormality in the handling of certain types of fatty acids in patients with multiple sclerosis. The effect of linoleate administered orally over a 5-day period on the plasma fatty acid concentrations, both in healthy controls and in patients with this disease, has been investigated. The results of this work, and their significance in terms of the aetiology and pathogenesis of multiple sclerosis, will be discussed.

Disorders of Lipid Metabolism

By ROSCOE O. BRADY (*Laboratory of Neurochemistry, National Institute of Neurological Diseases and Stroke, National Institutes of Health, Bethesda, Md. 20014, U.S.A.*)

The nature of the metabolic abnormalities in all of the known sphingolipid-storage diseases has been conclusively established within the past 6 years. This remarkable achievement is due largely to two developments. The first is the demonstration that the synthesis of the sphingolipids which accumulate in patients

Table 1. Constellation of metabolic diseases characterized by inability to degrade sphingolipids

Abbreviations: Cer, *N*-acylsphingosine (ceramide); Glc, glucose; PChol, phosphorylcholine; Gal, galactose; NAcNeu, *N*-acetylneuraminic acid; NAcGal, *N*-acetylgalactosamine.

Disease	Major sphingolipid accumulated	Enzyme defect
Gaucher's disease	<p>Ceramide glucoside (glucocerebroside)</p>	β -Glucosidase
Niemann-Pick disease	<p>Sphingomyelin</p>	Sphingomyelinase
Krabbe's disease	<p>Ceramide galactoside (galactocerebroside)</p>	β -Galactosidase
Metachromatic leucodystrophy	<p>Ceramide galactose 3-sulphate (sulphatide)</p>	Sulphatidase
Ceramide lactoside lipidosis	<p>Ceramide lactoside</p>	β -Galactosidase
Fabry's disease	<p>Ceramide trihexoside</p>	α -Galactosidase
Tay-Sachs disease	<p>Ganglioside GM₂</p>	Hexosaminidase A
Tay-Sachs disease variant	<p>Globoside (plus ganglioside (GM₂))</p>	Total hexosaminidase
Generalized gangliosidosis	<p>Ganglioside GM₁</p>	β -Galactosidase

with such storage diseases occurs at a normal rate. The second is the development of techniques for introducing radiotracer elements into the various lipid molecules. Investigations with labelled materials provided for the identification of the nature of the enzymic defects in Gaucher's disease, metachromatic leucodystrophy, Niemann-Pick disease and Fabry's disease in the short period of time from 1965 to 1967. These findings established the still unbroken precedent that all of the lipidoses are due to attenuation of specific catabolic enzymes normally required for the metabolic disposal of the accumulating lipid. There are now nine known diseases of this nature, and the respective enzymic abnormalities are indicated in Table 1. This hard-won knowledge has recently been put to practical clinical use. We can now accurately diagnose homozygous (afflicted) individuals, we can identify heterozygous carriers of these recessive diseases and, most importantly, procedures have now been developed for the accurate prenatal monitoring of pregnancies at risk for all of the lipid-storage diseases. The second portion of the presentation will deal with the profound alterations of ganglioside pattern and synthesis in DNA- and RNA-tumorigenic-virus-transformed cell lines.

Two Cerebral Sphingolipidoses (Metachromatic Leucodystrophy and Infantile Amaurotic Idiocy) as Human Inborn Errors of Metabolism

By H. JATZKEWITZ (*Max-Planck-Institut für Psychiatrie, Neurochemische Abteilung, München, West Germany*)

Both metachromatic leucodystrophy and infantile amaurotic idiocy arise from an accumulation of sphingoglycolipids. In cases of metachromatic leucodystrophy cerebroside sulphates (i.e. sulphatides or ceramide-galactose sulphate) accumulate. In patients suffering from infantile amaurotic idiocy gangliosides (i.e. ceramide-oligosaccharides + *N*-acetylneuraminic acid residues) and to a smaller extent their neuraminic acid-free residues are stored. This takes place especially in the brain, but also in other tissues. The major monosialoganglioside in the brain contains a single *N*-acetylneuraminic acid (sialic acid) residue, derived from a tetrahexosylceramide, and has the following structure:

where NeuNAc represents *N*-acetylneuraminic acid.

Both diseases, with a recessive mode of inheritance, are caused by a disturbance in the turnover of the accumulated sphingolipids due to a genetically

induced deficiency in the catabolizing enzyme system, which resides in the lysosomes.

Cerebroside sulphates are catabolized by arylsulphatase A (i.e. cerebroside sulphatase) to give cerebroside and sulphate. The major monosialoganglioside is degraded by different glycosidases, which hydrolyse the different hexoses of the oligosaccharide chain one after the other.

Sulphatides are constituents of brain white matter including myelin. A deficient cerebroside sulphatase gives rise to the demyelinating disease metachromatic leukodystrophy. Gangliosides are constituents of the grey matter. The inability to catabolize the major monosialoganglioside or its lower homologues leads to swollen and disintegrated neurons, giving rise to infantile amaurotic idiocy. Three types of glycosidases (β -galactosidase, EC 3.2.1.23; *N*-acetyl- β -D-hexosaminidase, EC 3.2.1.30; β -glucosidase, EC 3.2.1.21) are involved in the catabolism of the major monosialoganglioside. Since each of these glycosidases can be deficient, different variants of the disease have been observed. They are characterized by more or less visceral involvement. Even the classical type of the infantile amaurotic idiocy (Tay-Sachs disease), with the accumulation of the trihexosyl ganglioside (Tay-Sachs ganglioside), caused by a deficient *N*-acetyl- β -D-hexosaminidase, shows three variations. They depend on the different patterns of two multiple forms of this enzyme that occur in humans.

Diagnostic and therapeutic possibilities of these inborn errors of metabolism will be discussed.

The Metabolism of Phosphatidylcholine in Brain Tissue

By G. B. ANSELL [*Department of Pharmacology (Preclinical), University of Birmingham Medical School, Birmingham B15 2TJ, U.K.*]

Phosphatidylcholine is found in all the subcellular components of the nervous system; small amounts of choline plasmalogen are also present. Variations in its fatty acid composition in different subcellular components, deriving from a mixture of neuronal and glial cells, are known in some detail (Kishimoto *et al.*, 1969; Sun & Horrocks, 1970). Differences in the fatty acid composition of neuronal and glial cells have also been studied (Fewster & Mead, 1968; Hamberger & Svennerholm, 1971).

Both the concentration of phosphatidylcholine and its fatty acid composition change during the development of the brain. Thus in the rat a maximum rate of deposition occurs during a short period in early life, such that the concentration in the whole brain of a 6-day-old rat is only 60% of that of a