The Chemistry and Biochemistry of Polyprenyl Alcohols such as Solanesol and Dolichol

By R. A. MORTON (*Department of Zoology, University of Liverpool, Liverpool L69 3BX, U.K.*)

Rowland's (1956) discovery of solanesol ($C_{45}H_{93}OH$) in tobacco extended the geraniol-farnesol-geranylgeraniol series of unsaturated all-*trans*-polyprenols and gained additional significance by the discovery of quinones such as plastoquinones and ubiquinones with their all-*trans*-polyprenyl side chains.

Work carried out at Liverpool since then has shown that numerous other polyprenols occur naturally. The dolichols, obtained mainly from animal tissues, range from about C_{70} to C_{110} . The α -isoprene unit containing the hydroxyl groups is always saturated and *cis*, whereas the ω -isoprene unit and two internal isoprene units are *trans* with $n-4$ internal *cis*-isoprene units. A typical dolichol is thus $C_{100}H_{163}$ ($n=20$). Pig liver contains about 100mg of mixed dolichols ($n=17-22$)/kg.

Many plants contain significant quantities of deca-, undeca- and dodeca-prenols in which the ω -isoprene residue and three internal isoprene residues are *trans*, whereas the remainder, including the α -isoprene residue, are *cis*. Prenols from silver-birch wood include betulaprenol-7 ($n=7$), in which the ω -isoprene residue and the two other isoprene residues are known to be *trans*. An undecaprenol from *Lactobacillus plantarum* also has three *trans*-isoprene residues including the ω -isoprene residue, and eight *cis*-isoprene residues including the α -isoprene residue. *Aspergillus fumigatus* Fresenius polyprenols ($n=18-24$) have three isoprene residues (α , ψ and ω) saturated, and of the internal isoprene residues two are *trans* and the rest ($n=5$) are *cis*. More polyprenols are likely to be discovered.

Biosynthetic studies with stereospecifically double-labelled radioactive mevalonates are consistent with *cis* addition to all-*trans*-farnesyl pyrophosphate or all-*trans*-geranylgeraniol pyrophosphate, followed in some cases by reduction at one or both ends of the molecule.

In bacteria, the work of groups led by Robbins, Strominger, Lennarz, Baddiley and others indicates that undecaprenyl pyrophosphates act as carriers in the synthesis of complex polysaccharides, particularly in cell walls. This is an important advance, and dolichol metaphosphate glucose has been shown by Behrens & Leloir (1971) to act as an intermediate in glucose transfer in liver.

The study of polyprenols affords a good example of advances made possible by newer techniques of separation, and characterization by means of spectroscopic methods. The biosynthetic work owes much to stereospecific labelling, and the carrier role illustrates the confluence of different types of research.

The Liverpool contributions in this field owe much to the leadership shown by Dr. F. W. Hemming and Dr. J. F. Pennock and to the careful work of research students.

Behrens, N. H. & Leloir, L. F. (1971) *Arch. Biochem. Biophys.* **143**, 375

Rowland, R. L., Latimer, P. H. & Giles, J. A. (1956) *J. Amer. Chem. Soc.* **80**, 613

Metabolism of β -Apocarotenals

By HARMEET SINGH, A. KRISHNA MALLIA and H. R. CAMA (*Department of Biochemistry, Indian Institute of Science, Bangalore-12, India*)

β -Apocarotenals are formed during the chemical oxidation of β -carotene. Although apocarotenals were suggested to be the intermediates during the cleavage of β -carotene in animals, so far there has been no proof to support this hypothesis. In the present study metabolism of apocarotenals and their epoxides was undertaken in order to investigate the mechanism of cleavage of the β -ionone-ring-substituted carotenoids.

Various β -apocarotenals were prepared by oxidation of β -carotene with H_2O_2 in the presence of $KMnO_4$. They were separated and purified by chromatography on deactivated alumina columns. The apocarotenals separated were β -apo-6'-carotenal, β -apo-8'-carotenal, β -apo-10'-carotenal and β -apo-12'-carotenal, with absorption maxima in light petroleum at 414, 437, 457 and 470nm respectively. When tested by the rat-growth assay method, they were found to have biopotencies of 38.5, 31.3, 75.9 and 44.2% respectively of that of β -carotene. They also gave rise to vitamin A in rats, and different amounts of vitamin A were found in the liver.

5,6- and 5,8-Monoepoxides of various β -apocarotenals were also prepared and characterized, but these epoxides did not support the growth of earlier claim (Krishna Mallia *et al.*, 1970) that the double bond in the 5,6-position of the β -ionone ring is essential for exerting vitamin A activity.

Krishna Mallia, A., Jacob, J., John, K. V., Lakshmanan, M. R., Jungalwala, F. B. & Cama, H. R. (1970) *Indian J. Biochem.* **7**, 102

Recent Developments in the Study of the Biosynthesis of Carotenoids

By T. W. GOODWIN (*Department of Biochemistry, University of Liverpool, Liverpool L69 3BX, U.K.*)

Many of the recent developments in the study of carotenoid biosynthesis in our laboratory were achieved by using various species of stereospecifically labelled mevalonates, and have been reviewed in