Co-ordinated Synthesis of Membrane Phospholipids with the Formation of Ribosomes and Accelerated Protein Synthesis during Growth and Development

By J. R. TATA (*National Institute for Medical Research, Mill Hill, London NW7 1AA, U.K.*)

In several hormone-dependent growth and developmental systems studied, the rate of labelling of membrane phospholipids is enhanced in all major subcellular particulate fractions (nuclear, mitochondrial and microsomal) after hormone administration. The net accumulation of phospholipids, as well as protein and RNA, is most marked in the rough endoplasmic reticulum. Growth of the liver induced by growth hormone, by thyroid hormone, by testosterone or during regeneration is characterized by a preferential acceleration of the labelling with ^{32}P of sphingomyelin relative to that of phosphatidylcholine or phosphatidylethanolamine. Time-course analysis of six growth and developmental systems studied in our laboratory has shown that the enhancement of the rate of membrane phospholipid synthesis coincides with the rather abrupt increase in ribosomes, especially in the rough endoplasmic reticulum. There is some mechanism in the cell that tightly co-ordinates the formation of membranes when an increased demand is made for protein synthesis, as during growth and development.

Some recent histochemical, electron-microscopic and subcellular fractionation studies in induced amphibian metamorphosis will also be presented, which suggest that newly formed protein may be initially restricted to small areas of the rough endoplasmic reticulum in the perinuclear region. The significance of these findings is that, besides secretion of proteins, attachment of ribosomes to membranes of the endoplasmic reticulum during growth and development may serve to topographically segregate different populations of ribosomes synthesizing different groups of proteins.

Studies on the Cell Membrane of Bacilli

By WILLIAM J. LENNARZ (*Department of Physiological Chemistry, The Johns Hopkins University School of Medicine, Baltimore, Md. 21205, U.S.A.*)

Because phospholipids comprise 25–50% of the dry weight of bacterial membranes, the enzymes responsible for their synthesis must play an important role in the biogenesis of the cytoplasmic membrane. Studies with 'ghosts' of *Bacillus megaterium* have revealed that all the enzymes responsible for synthesis of phosphatidylethanolamine and phosphatidylglycerol from phosphatidic acid are localized

in the cell membrane. Current studies on the disaggregation, fractionation and reconstitution of bacterial membranes are based on the premise that formation of all, or at least part, of the basis membrane structure is a self-assembly process. One criterion of reconstitution of membranes, namely regain of enzymic activity, is being studied with the enzymes of phospholipid synthesis. In addition, techniques are being developed for the selection of membrane mutants of *bacilli* that are deficient in lipids or proteins essential for the structural integrity of the membrane.

Turnover and Exchange of Phospholipids in the Membranes of the Nervous System

By R. M. C. DAWSON, ELAINE MILLER and F. B. JUNGALWALA (*Department of Biochemistry, Agricultural Research Council Institute of Animal Physiology, Babraham, Cambridge CB2 4AT, U.K.*)

The brain is an organ that, unlike the liver, does not secrete phospholipids. Consequently phospholipid synthesis at the termination of growth is required only for turnover and replacement of the cerebral membranes. Synthesis of the membrane phospholipids *de novo* largely occurs in the endoplasmic reticulum, although formation of some phosphatidic acid and diphosphatidylglycerol takes place in mitochondria. The synaptic vesicles in the nerve terminal (synaptosome) can synthesize phosphatidylcholine from CDP-choline. The endoplasmic reticulum can carry out base-exchange reactions of phospholipids, including that of choline by a non-energy-requiring Ca^{2+} -dependent enzymic exchange process. On intraventricular injection of phospholipid precursors into the brain of adult animals a substantial turnover of myelin phospholipids can be demonstrated, but a small more slowly exchanging pool is also apparent. ^{32}P -labelled brain microsomal fraction can exchange phospholipids on incubation with unlabelled mitochondria, and vice versa. The exchange is dependent on a heat-labile macromolecular factor in the supernatant, and the concentration of this is rate-limiting. ^{32}P -labelled microsomal fraction exchange phospholipid more slowly with intact synaptosomes, and during the incubation the intraneural mitochondria in the synaptosomes acquire phospholipid most actively. However, on subfractionation of synaptosomes all the isolated membranes can participate in phospholipid exchange processes. ^{32}P -labelled microsomal fraction does not exchange phospholipids on incubation with myelin particles, and the latter particles inhibit exchange processes between other membranes. The reasons for and mechanism of phospholipid exchange in the intact brain will be discussed.