The membrane is isolated from the purified virus by treatment with detergents and subsequent sucrose-density-gradient centrifugation. It contains only one glycoprotein species (mol.wt. 50000), which is synthesized during the virus infection. One virus particle contains about 500 molecules of this glycoprotein. The host (baby-hamster kidney cell line BHK21) plasma membrane, on the other hand, is very complex and contains more than 15 different proteins.

The lipids of the virus membrane consist mainly of free cholesterol, phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine, sphingomyelin and sialolactosylceramide. The cholesterol/phospholipid molar ratio is 1:1. One virus particle contains about 10000 cholesterol-phospholipid pairs. Each class of the polar lipids is characterized by a specific fatty acid chain composition that is nearly identical in Semiliki Forest virus and host plasma membrane. Virus lipids are synthesized in the cell both before and during the infection. The virus membrane resembles in its lipid composition the host plasma membrane but is different from the endoplasmic reticulum.

- Acheson, N. H. & Tamm, I. (1967) *Virology* **32**, 128
 Compans, R. W. (1971) *Nature New Biol. (London)* **229**, 114
 Friedman, R. M. & Pastan, I. (1969) *J. Mol. Biol.* **40**, 107
 Harrison, S. C., Caspar, D. L. D., Camerini-Otero, R. D. & Franklin, R. M. (1971) *Nature New Biol. (London)* **229**, 197
 Klenk, H.-D. & Choppin, P. W. (1970) *Virology* **40**, 939
 Pfefferkorn, E. R. & Hunter, H. S. (1963) *Virology* **20**, 433
 Simons, N. & Kääriäinen, J. (1970) *Biochem. Biophys. Res. Commun.* **38**, 981
 Strauss, J. H., Burge, B. W., Pfefferkorn, E. R. & Darnell, J. E. (1968) *Proc. Nat. Acad. Sci. U.S.* **59**, 533
 Tiffany, J. M. & Blough, H. A. (1970) *Proc. Nat. Acad. Sci. U.S.* **65**, 1105

The Role of Lipids in Membrane Structure

By FREDERICK L. CRANE (*Division of Biological Sciences, Purdue University, Lafayette, Ind. 47907, U.S.A.*)

Cytochrome oxidase from mitochondria contains two molecules of haem, a protein (mol.wt. 26000) and one core protein IV (mol.wt. 56000). When free of lipid it forms pentameric clusters of rods 9.0nm (90Å) in diameter made up of 5.0–6.0nm (50–60Å) subunits and has no enzyme activity. On addition of phospholipids it disperses into a membrane sheet with 5.0–6.0nm (50–60Å) subunits separated by dark regions (negative stain). Sections show unit membrane structure 5.0nm (50Å) thick. Freeze-etch

surfaces have a rough appearance, and globular units appear in the sheet structure in cross-section. Membrane formation and activity restoration are saturated at 65 g-atom of phosphorus/mol of haem *a*. At high concentrations excess of lipid can be removed by centrifugation. As lipid content is increased the subunits show increased separation until saturation is reached, indicating that the white globular subunits (negative stain) are the oxidase. Association of lipid and protein in membrane is accompanied by a loss of the i.r. rocking band of lipid at 720cm⁻¹, increase of the thermal stability of enzyme activity, and increase of negative circular dichroism at 225 nm and increase of positive circular dichroism at 200 nm as when a protein is transferred from water to 2-chloroethanol. 1-Anilino-naphthalene-8-sulphonate-fluorescence induced by Triton-dispersed oxidase is 30% decreased by the addition of phospholipid. The results are consistent with a mosaic membrane formed by insertion of protein into the hydrocarbon phase of a phospholipid bilayer. A stabilizing influence of protein on the bilayer is shown, since oxidase plus diolefin form membrane, whereas diolefin alone does not form a bilayer. Formation of a diolefin-oxidase membrane does not restore oxidase activity. Specificity of protein in lipid association to form membrane is shown, since isolated haem *a* protein will form membrane, which shows smaller 3.0–4.0nm (30–40Å) subunits, but core protein IV will not form membrane. Addition of core protein IV to haem *a*-lipid membrane does, however, increase oxidase activity. Thus cytochrome oxidase provides a simple model mosaic membrane in which hydrophobic association of lipid with a functional protein can be studied.

A Soluble Mitochondrial Adenosine Triphosphate Synthetase Complex and its Stimulation by the Membrane Fraction

By D. RAO SANADI, R. J. FISHER, S. S. KAPLAY, J. C. CHEN and B. P. SANI (*Department of Cell Physiology, Boston Biomedical Research Institute, 20 Stamford Street, Boston, Mass. 02114, U.S.A.*)

This communication reports that a highly purified, soluble, energy-transfer factor preparation carries out an oligomycin- and uncoupler-sensitive ATP-P_i exchange reaction. It has been named the ATP synthetase complex. The exchange activity (up to 35 nmol/min per mg of protein) is dependent on Mg²⁺ and is stimulated threefold by dithiothreitol. The reaction is linear for at least 2 h. The activity is inhibited by low concentrations of oligomycin or carbonyl cyanide trifluoromethoxyphenylhydrazone.