deteriorating as a result of the unplanned location of industries and urban development.

Experience in other countries has indicated certain basic requirements for tackling the problems of pollution. These are legislation, technical expertise, funds, the co-operation of industries and the need for a master plan for the location of industries. It is time that the industrialists realize their responsibilities and take effective action in the prevention and abatement of environmental pollution arising out of industrial wastes.

Influence of D-Alanine, Glycine and Egg Albumin on Biochemical Oxygen Demand

By B. L. C. RAJAN (*Visvesvaraya College of Engineering, University of Bangalore, Bangalore-1, India*)

Experiments were carried out by adding D-alanine, glycine and egg albumin in different combinations to domestic-sewage and industrial-waste waters and following the 5-day biological oxygen demand of these mixtures. The results indicated the following. (1) There was variation in the increase in biological oxygen demand in the presence of equal amounts of the amino acids and egg albumin; the increase was greatest with D-alanine and least with glycine. Of these chemicals D-alanine apparently stimulated bacterial growth to the greatest extent and glycine to the least extent. (2) The increase in biological oxygen demand in the presence of egg albumin was not appreciable during the first 36h, presumably owing to a lag phase in the growth of bacteria. (3) About 90% of the biological oxygen demand was satisfied in 5 days in the presence of D-alanine, whereas 80 and 70% of the biological demand were satisfied during that time in the presence of egg albumin and glycine respectively. (4) With increased concentrations of amino acids and egg albumin in the waste water the increases in biological oxygen demand were not proportional. (5) The 'K' values also closely followed the values for the biological oxygen demand.

These observations suggest that addition of such organic chemicals as those used in this work is useful in the treatment of organic industrial-waste waters deficient in nitrogen.

Effect of Ultraviolet Irradiation on Chlorine Residuals in Polluted Waters

By T. M. AHMEER AHMED (*Civil Engineering Department, Visvesvaraya Engineering College, Bangalore-1, India*)

In this study over 50 samples of river water polluted with sewage and industrial wastes were used. They contained high concentrations of suspended solids and dissolved impurities. All the samples were ex-

posed to ultraviolet irradiation for 6h. There was a decrease of about 60% in the chlorine residuals in the first 2h. The decrease was less at greater depths when the samples were kept in experimental tanks. The decrease was increased with a greater intensity of illumination. Along with the decrease in the chlorine residuals there was a fall in pH value.

A simplified water system obtained by passing distilled water through a column of Amberlite MB-1 resin at pH7 was compared with the polluted waters for the decrease in chlorine residuals. There was about an 80% decrease in 6h. In the same period the polluted waters showed a decrease of about 70%. This was due to the presence of suspended and dissolved impurities.

These results show that the depth of the tank, intensity of illumination, time, pH and the dissolved and suspended impurities of the sample influence the decrease in chlorine residuals effected by ultraviolet irradiation. These observations have a bearing on chlorination practice in the treatment of waste waters.

Biochemical Evaluation of Pesticides

By LEELA IYENGAR, M. P. PANDEY, C. VENKOBACHAR and A. V. S. PRABHAKARA RAO (*Department of Civil Engineering, Indian Institute of Technology, Kanpur, Kanpur-16, India*)

In view of the increasing use of pesticides and insecticides in agriculture and of the problem of development of resistance and effects of insecticide combinations, a biochemical evaluation of pesticides was undertaken. The effects of chemicals such as DDT [1,1,1-trichloro-2,2-di-(*p*-chlorophenyl)ethane], BHC (lindane, 1,2,3,4,5,6-hexachlorocyclohexane), aldrin (1,2,3,4,10,10-hexachloro-1,4,4a,5,8,8a-hexahydro-1,4,5,8-dimethanonaphthalene), endrin (1,2,3,4,10,10-hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-1,4,5,8-endo-endo-dimethanonaphthalene), malathion (*S*-1,2-dicarbethoxyethyl *OO*-dimethyl phosphorothiolothionate), Rogor [*OO*-dimethyl *S*-(*N* - methylcarbamoylmethyl)phosphorodithioate], parathion (*OO*-diethyl *O*-*p*-nitrophenyl phosphorothionate), Diazinon (*OO*-diethyl *O*-2-isopropyl-6-methylpyrimidin-4-yl phosphorothionate), chlordane (1,2,4,5,6,7,8,8-octachloro-2,3,3a,4,7,7a-hexahydro-4,7-methanoindene) and Thiometon *OO*-dimethyl *S*-2-(ethylthioethyl)phosphorodithioate on amylase, catalase, transaminases, proteases, lipases and dehydrogenases from such different sources as seeds and mosquito larvae were studied. The results indicated that there was a delay or inhibition of germination of some seeds. The inhibitory patterns

of some of the enzymes were indicative of the concentrations of the insecticides. Synergistic or antagonistic effects of combinations of pesticides were obtained by using simple techniques.

Studies on Microbial Degradation of Herbicides in Soil

By K. SRIRAMA RAJU and G. RANGASWAMI (*Department of Microbiology, University of Agricultural Sciences, Bangalore-24, India*)

The hazards due to cumulative effects of agricultural chemicals, particularly pesticides, added to cultivated soil have been a subject of investigation in recent years. In order to understand their persistence in the soil under tropical conditions, as in India, a series of studies were undertaken in the laboratory on the microbial degradation of herbicides in soil.

When four common herbicides, atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine), simazin [2-chloro-4,6-bis(ethylamino)-sym-triazine], diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] and Tok-E25 (2,4-dichlorophenyl 4-nitrophenyl ether), were applied to the soil, there was a change in the populations of bacteria, actinomycetes and fungi in the soil. Atrazine caused the maximum change. When the soil was enriched with 1000 p.p.m. of the herbicides, several bacteria and fungi resistant to the herbicides were isolated. Some of the fungi and bacteria were able to resist the herbicides at concentrations as high as 7500 p.p.m., but the actinomycetes were not resistant. Some of the bacteria, fungi and actinomycetes were able to utilize the herbicides as a source of carbon or nitrogen, and a few were able to grow in artificial media containing the herbicides as the sole carbon and nitrogen source. The growth of the latter organisms was, however, slow in the laboratory media, indicating their low efficiency in utilizing these compounds. Of the four herbicides tested, diuron was relatively the most resistant to microbial degradation.

Seasonal Biochemical Acclimatization in *Channa punctatus*

By ASIT BARAN DAS [*Department of Biological Sciences (University of Sambalpur), V.S.S. Medical College, Burla (Sambalpur), Orissa, India*]

The seasonal biochemical changes in different tissues of the *Channa punctatus*, such as gill, brain, liver and muscle, started with a gradual rise in the total RNA/DNA ratio from a low summer value (1.2–2.0:1.0) to a peak winter value (2.1–3.2:1.0).

This was closely followed by an augmentation of the total free amino acids/DNA ratio with a summer value of 0.35–0.44:1.0 and a winter value of 0.45–0.75:1.0. The winter peaks in both RNA and amino acid concentrations in the tissues were maintained from December through to March. The peaks in total protein accumulation (protein/DNA ratio) in different tissues were found to be 13.7:1.0, 17.3:1.0, 34.7:1.0 and 158.6:1.0 in gill, brain, liver and muscle respectively towards the end of winter (February–March), and the lowest values were found to be 9.4:1.0, 10.6:1.0, 16.3:1.0 and 87.9:1.0 in gill, brain, liver and muscle respectively during the summer and the rainy season (April–July). The hepatic glycogen content was maintained relatively constant at 0.31–0.55 g/100 g of liver throughout the year, except for an abrupt increase to 2.1 g/100 g of liver during the winter. The possible significance of the concurrent cyclic variations in the different biochemical parameters and their relationships to the rhythmic fluctuations in metabolic rate and in cytological parameters of liver will be discussed.

Breeding of an Indian Carp (*Cirrhana mrigala*) under Different Environmental Temperatures

By A. K. DAS (*Kandaswami Lane, Sultan Bazaar, Hyderabad 1, India*)

The breeding season for the major Indian carps is the monsoon from the end of June to August, when the maturation of gonads usually occurs. In a representative of this group of fish, *Cirrhana mrigala*, adaptation and spawning during the hot summer was successfully carried out by supplementing the diet used by the breeders with protein-rich pulses and lowering the environmental temperature from 42°C to 27°C by cooling with artificial 'rain-drops'. There was a carry-over effect of the above treatment in the progeny, which showed a faster growth rate.

Studies on the Development of Catalysts for the Removal of Carbon Monoxide from Motor-Car Exhausts

By C. R. RAMANATHAN and M. RAVINDRAM (*Department of Chemical Engineering, Indian Institute of Science, Bangalore-12, India*)

Experimental investigations were undertaken with a view to developing stable catalysts capable of converting CO contained in motor-car exhausts into CO₂. Initially CO (in low concentrations in air) oxidation was studied in a semi-micro flow catalytic reactor in the temperature range 200–400°C over pure CuO