where E-OH represents the enzyme with its non-thiol binding site and ACP-SH represents acyl-carrier protein.

- Joshi, V. C., Plate, C. A. & Wakil, G. J. (1970) *J. Biol. Chem.* **245**, 2857
 Lynen, F. (1967) in *Organizational Biosynthesis* (Vogel, H. J., Lampen, J. O. & Bryson, V.), p. 243, Academic Press, New York
 Phillips, G. T., Nixon, J. E., Abramovitz, A. S. & Porter, J. W. (1970) *Arch. Biochem. Biophys.* **138**, 357.
 Plate, C. A., Joshi, V. C. & Wakil, S. J. (1970) *J. Biol. Chem.* **245**, 2868

Incorporation of Fatty Acids into Monogalactosyl Diglycerides by Spinach Leaf Cell-Free Preparations

By S. S. BAIWA and P. S. SASTRY (*Department of Biochemistry, Indian Institute of Science, Bangalore-12, India*)

Mono- and di-galactosyl diglycerides are the major lipid components of chloroplasts and, in contrast with phospholipids, these plant galactolipids contain unsaturated fatty acids, usually α -linolenate, in both the 1- and the 2-position of the diglyceride moiety (Sastry & Kates, 1964). Although it has been shown in spinach that monogalactosyl diglyceride is synthesized from diglyceride and UDP-galactose (Neufeld & Hall, 1964; Ongun & Mudd, 1968), the enzymic mechanisms for the biosynthesis of the dilinolenin moiety in monogalactosyl diglyceride are not clearly understood.

Dialysed spinach (*Tetragonia expansa*) leaf homogenate and chloroplast preparations were found to incorporate 1-¹⁴C-labelled palmitate, stearate, oleate and linolenate into monogalactosyl diglyceride. This incorporation was much higher in the chloroplast fraction. It was enhanced by CoA and ATP. The activity increased with time, showing that it was enzymic, and was optimum at 30°C. A higher rate of incorporation was obtained with unsaturated fatty acids than with saturated fatty acids. Addition of galactosyl glycerol significantly stimulated the incorporation of oleate and linolenate only. These studies indicate the possible existence of deacylation and reacylation with specific fatty acids as a mechanism for attaining the peculiar fatty acid distribution in monogalactosyl diglyceride of higher plants.

- Neufeld, E. F. & Hall, C. W. (1964) *Biochem. Biophys. Res. Commun.* **14**, 503.
 Ongun, A. & Mudd, J. B. (1968) *J. Biol. Chem.* **243**, 1558
 Sastry, P. S. & Kates, M. (1964) *Biochemistry*, **3**, 1271

The Fatty Acid Composition of Different Classes of Lipids in Developing Human Brain

By P. SRINIVASA RAO and KALLURI SUBBA RAO (*National Institute of Nutrition, Hyderabad-7, A.P., India*)

Information on the lipid composition of developing human brain, particularly during its intrauterine growth, is scanty. Whatever data available so far are concerned mainly with the quantitative changes in various lipids. However, it is equally important to consider the qualitative changes in these lipids that would be reflected in the fatty acid constituents of given class of lipids. An attempt has therefore been made to determine the fatty acid composition of various classes of lipids in different regions (namely cerebrum, cerebellum and medulla oblongata) of brains of foetuses aged 13, 22, 26, 30, 34 and 40 weeks. The extracted lipids were fractionated into neutral lipids (cholesteryl esters, glycerides etc.), glycolipids and phospholipids on a silicic acid column. The fatty acid pattern, after methanolysis of these fractions, was determined by g.l.c. in a column packed with diethylene glycol succinate in Chromosorb W. The fatty acid profile of neutral lipids indicated palmitate, stearate and oleate as major components in all the regions of brain studied. This trend remained almost constant with the progress of gestation. Analysis of the glycolipid fraction showed that lignocerate (C_{24:0}) and nervonate (C_{24:1}) were absent at earlier periods of intrauterine life, but made their appearance towards term, though at a much lower concentration than expected. Studies with the phospholipid fraction also pointed out palmitate, stearate and oleate as predominant components and this pattern did not alter with the gestational age. Analysis of fatty acids in the glycolipid fraction of the brain of a child (4 years old) who had died of kwashiorkor revealed a considerably low content of lignocerate and nervonate, indicating the possibility of defective myelination.

The Gross Fatty Acid Composition of the Body Fat from the Termite (*Odontotermes assmuthi*) Queen

By S. BASALINGAPPA,* R. C. BADAMI† and S. M. KUDARI† (**Department of Zoology and †Department of Chemistry, Karnatak University, Dharwar, Mysore, India*)

An analysis of the total body fat of the termite (*Odontotermes assmuthi*) queen by reversed-phase partition column chromatography (Badami, 1964, 1965) reveals a total of 13 fatty acids as against nine