primarily involved in this inhibition; other metal ions such as Fe^{2+} and Zn^{2+} are not effective. At optimum pH (5.5) the enzyme is not inhibited by *p*-hydroxymercuribenzoate; however, at pH 5.0 the enzyme is markedly inhibited by *p*-hydroxymercuribenzoate, though this inhibition can be reversed by cysteine. The possible role of chicken brain arylsulphatase A in the metabolism of sulphated compounds will be discussed.

Austin, J. H., Balasubramanian, A. S., Pattabiraman, T. N., Saraswathi, S., Basu, D. K. & Bachhawat, B. K. (1963) *J. Neurochem.* **10**, 805

Farooqui, A. A. & Bachhawat, B. K. (1971) *J. Neurochem.* **18**, 635

Mehl, E. & Jatzkewitz, H. (1968) *Biochim. Biophys. Acta* **151**, 619

Roy, A. B. (1970) *Biochim. Biophys. Acta* **198**, 76

Ribonuclease in Yoshida Ascites Sarcoma

By ARUNA BHANDARKAR, G. RAMANANDA RAO and M. SIRSI (*Microbiology and Pharmacology Laboratory, Indian Institute of Science, Bangalore-12, India*)

In animal cells the ribonuclease-ribonuclease inhibitor complex provides possible control at the level of RNA degradation through the regulation of ribonuclease activity (Kraft & Shortman, 1970). Anti-cancer drugs such as L-asparaginase and 9 α -fluoroprednisolone, which cause regression of lymphoid tumours, induce a significant increase in acid ribonuclease activity. It has been postulated that increased acid ribonuclease activity is causally linked with the involution of lymphoid tissue (Ambellan & Hollander, 1966). The ribonucleases in Yoshida ascites-sarcoma cells and the inhibitors acting on them were studied to elucidate the mechanism of this control in cancer cells.

Two types of ribonucleases, distinguished on the basis of their pH optima, were found in cell-free extracts of 4-day-old Yoshida ascites sarcoma. The specific activity of free acid ribonuclease (pH 5.8) was 1.58 units/mg of protein, whereas that of free alkaline ribonuclease (pH 7.8) was 0.32 unit/mg of protein (1 unit is the amount of enzyme that causes an increase of 1.0 E_{260} unit in 30 min under the assay conditions used). Increased alkaline ribonuclease activity was observed on the addition of thiol-group reagents such as *p*-hydroxymercuribenzoate to the assay system; this increase was proportional to the concentration of *p*-hydroxymercuribenzoate from 0.1 to 1 mM. Addition of *N*-ethylmaleimide and iodoacetate also resulted in increased activity, indicating the presence of an inhibitor protein with reactive thiol groups. No increased acid ribonuclease activity was found on adding *p*-hydroxymercuribenzoate, indicating thereby the absence of an inhibitor of acid ribonuclease.

The specific activities of alkaline ribonuclease and acid ribonuclease during tumour progression were studied.

A. B. is grateful to the Lady Tata Memorial Trust, Bombay, for financial support.

Ambellan, E. & Hollander, V. P. (1966) *Cancer Res.* **26**, 903-909

Kraft, N. & Shortman, K. (1970) *Biochim. Biophys. Acta* **217**, 164-175

Bull Semen Phosphodiesterase

By ASHOK KHER and S. R. ANAND (*National Dairy Research Institute, Karnal, India*)

Semen is a rich source of various nucleotide-splitting enzymes (Mann, 1964). Adenylate cyclase, which catalyses the conversion of ATP into cyclic AMP (adenosine 3':5'-cyclic monophosphate), has been recognized as a second messenger in the action of various hormones. Robinson *et al.* (1968) have proposed that the intracellular concentration of cyclic AMP is regulated by two enzymes: (1) adenylate cyclase; (2) phosphodiesterase, which hydrolyses cyclic nucleotide to AMP. The presence of adenylate cyclase has been reported in monkey spermatozoa (Hoskin & Stephens, 1969) and in sea-urchin spermatozoa (Sutherland, 1970). There is no report of the presence of phosphodiesterase activity in bull semen. We therefore tested this and detected phosphodiesterase activity in bull spermatozoa and seminal plasma with bis-(*p*-nitrophenyl) phosphate as substrate. Seminal plasma phosphodiesterase was purified five- to ten-fold by precipitation with ammonium sulphate followed by DEAE-cellulose chromatography. The enzyme exhibits a broad pH optimum at 7-8. Its ability to attack DNA, RNA, cyclic AMP and other nucleotides, as well as other properties, will be reported.

Hoskin, D. D. & Stephens, D. T. (1969) *Biochim. Biophys. Acta* **191**, 292

Mann, T. (1964) *The Biochemistry of Semen and of the Male Reproductive Tract*, John Wiley and Sons, New York

Robinson, G. A., Butcher, R. W. & Sutherland, E. W. (1968) *Annu. Rev. Biochem.* **37**, 149

Sutherland, E. W. (1970) *Fed. Proc. Fed. Amer. Soc. Exp. Biol.* **29**, 608