last 2 weeks of a 4-week incubation period. Simazin depressed the actinomycete population in the beginning and stimulated it during the last 2 weeks. The others did not have any effect on actinomycete population. Fungal population was stimulated by Tok-E-25, whereas Benlate depressed it initially and stimulated it during the last 2 weeks of incubation.

Simazin drastically decreased the soil respiration initially and thereafter it had no effect. The others did not affect the respiration. The effect of Benlate at two concentrations on soil ammonification and nitrification rates was studied, with media containing peptone or ammonium sulphate or nitrite. In peptone medium, ammonification and nitrite formation were unaffected but nitrate formation was drastically depressed, resulting in accumulation of nitrite. In ammonium sulphate medium and nitrite medium Benlate at both concentrations did not have any marked effect initially, but during the latter half of incubation period nitrification was gradually depressed or it remained steady at a low rate. The implications of these effects will be discussed.

Influence of Growth Hormone on Thymus Metabolism *in vivo* and *in vitro*

By M. R. PANDIAN and G. P. TALWAR (*Department of Biochemistry, All India Institute of Medical Sciences, New Delhi-16, India*)

Administration of bovine growth hormone leads to an enhancement of DNA synthesis in lymphoid organs and recovery of the immune response in hypophysectomized rats. Similar effects of the hormone are observed in 'plateaued' rats (i.e. rats which have attained maximum weight). Injection of rabbit anti-(bovine growth hormone) globulins, in contrast with normal rabbit globulins, over 5 days causes a decrease in the weight of the thymus and in the rate of DNA synthesis in this organ. Thymus is also the organ in which stimulation of DNA synthesis is observed at a time earlier than that in spleen and lymph nodes after a single injection of bovine growth hormone. The hormone stimulates not only the incorporation of [³H]thymidine into DNA in the cortical cells, but also the incorporation of [³⁵S]sulphate into trichloroacetic acid-insoluble biopolymers reported to be elaborated in the medullary area of the thymus.

A system for study of the action of bovine growth hormone on thymus *in vitro* has been devised. There is an obligatory requirement for Ca²⁺, but not for foetal calf serum, in the medium for the hormone effect. An early action of the hormone is the enhanced incorporation of [³H]uridine into RNA in thymocytes, this being followed by a stimulation of the synthesis of proteins and DNA. The stimulatory action of the hormone on RNA synthesis is not due to

a facilitated uptake of the radioactive uridine by the cells under hormonal influence.

Mechanism of Regulation of the Immune Response

By R. KRISHNARAJ and G. P. TALWAR (*Department of Biochemistry, All India Institute of Medical Sciences, New Delhi-16, India*)

It is currently believed that in the development of the immune response the interaction with specific 'receptors' on the surface of lymphocytes is the first event leading to the proliferation and differentiation of these cells. The proliferation of lymphocytes can also be triggered by general mitogens such as phytohaemagglutinin.

On the premise that the initial interaction takes place at the membrane, we have studied the adenylate cyclase activity in the primary T&B lymphoid cells, as well as in lymphocytes present in secondary lymphoid organs. Incubation of cells with phytohaemagglutinin stimulates the adenylate cyclase activity of thymocytes and thymus-derived lymphocytes. The obvious implication has been tested by the ability of cyclic AMP (adenosine 3':5'-cyclic monophosphate) to cause blast transformation. Some information is also available on the probable sites affected by cyclic AMP that ultimately lead to the replication of these cells.

Studies on some Aspects of Ascorbate Metabolism in Adrenalectomized Rats

By MAHESH G. NATHANI, IDREES BHAI, N. NATH, H. F. DAGINAWALA and M. C. NATH (*Department of Biochemistry, University of Nagpur, Nagpur, India*)

Male albino rats were adrenalectomized and the ascorbate, dehydroascorbate and dioxogulonate contents in liver and urine, but only the total ascorbate in blood, were determined at the intervals of 4, 8 and 12 days after adrenalectomy. The activities of ascorbate-synthesizing enzymes (D-glucurono- δ -lactone hydrolase, L-gulonon- γ -lactone hydrolase and L-gulonon- γ -lactone oxidase) in liver were measured, as were the activities of ascorbate-degrading enzymes (dehydroascorbate and 2,3-dioxoaldonate decarboxylase) in liver and kidney.

In adrenalectomized rats there was a slight decrease in the ascorbate content and an appreciable decrease in the dehydroascorbate content of the liver. On the other hand there was about a twofold increase in the dioxogulonate content of liver. The ascorbate and dehydroascorbate contents in the