streptozotocin into the intact animal before excision of the heart, glycogen synthesis from $[U-^{14}C]$ glucose was decreased by over a half whereas glucose oxidation to $^{14}CO_2$ was not affected. The rate of glycogen synthesis could be restored to normality by the addition of insulin (2munits/ml) to the perfusion medium. Insulin has been shown to activate *in vitro* adipose-tissue cyclic AMP (adenosine 3':5'-cyclic monophosphate) phosphodiesterase (EC 3.1.4.-) (Loten & Sneyd, 1970) and purified ox heart cyclic AMP phosphodiesterase (Senft *et al.*, 1968). A direct effect of insulin on cyclic AMP phosphodiesterase or cyclic GMP (guanosine 3':5'-cyclic monophosphate) phosphodiesterase has been postulated but is still controversial (Sutherland & Robinson, 1969; Thompson & Appleman, 1971).

The heart-perfusion method, apparatus and preparation of the streptozotocin-diabetic rats were essentially as described by Chain *et al.* (1969). Cyclic AMP phosphodiesterase and cyclic GMP phosphodiesterase activities were assayed in 1000g sediment and supernatant by the method described by Butcher & Sutherland (1962), with some modifications; cyclic GMP phosphodiesterase activity was assayed by using cyclic GMP as substrate in place of cyclic AMP. There were consistent and significant decreases (25–35%) in both activities in the streptozotocin-diabetic perfused rat heart. The decreases in these activities were more evident in nuclear and membrane fractions. Insulin (10munits/ml) in the perfusate significantly increased both cyclic AMP phosphodiesterase and cyclic GMP phosphodiesterase activities in normal diabetic perfused hearts.

The patterns of decrease or increase in phosphodiesterase activity are similar whether we use cyclic AMP or cyclic GMP as substrate. The increase in the phosphodiesterase activities with insulin is more marked in the 1000g-sediment fraction. This is the first report of the stimulation of cyclic AMP phosphodiesterase and cyclic GMP phosphodiesterase activities in an isolated perfused organ by insulin. No stimulation of the phosphodiesterase activities by insulin was observed when the homogenate was treated with insulin (3units/ml) *in vitro*. Senft *et al.* (1968) described an activation of purified ox heart cyclic AMP phosphodiesterase *in vitro* by insulin. The regulation of cyclic AMP phosphodiesterase and cyclic GMP phosphodiesterase activities may control the tissue concentrations of cyclic nucleotides, the latter in turn regulating the rate of glycogen synthesis and degradation as suggested by Senft *et al.* (1968). The increase in cyclic nucleotide phosphodiesterase activities here demonstrated may therefore be partially responsible for the increased glycogen synthesis found in the perfused insulin-treated rat heart (Chain *et al.*, 1969). The magnitude of the increase in glycogen synthesis in normal and diabetic perfused hearts (Chain *et al.*, 1969) suggests, how-

ever, that there might be some other additional factor(s) also contributing to the stimulatory action of insulin on glycogen synthesis in the perfused rat heart.

The skilled technical assistance of Miss Helga Wille is gratefully acknowledged. We thank Dr. Anne Beloff-Chain for useful discussions.

- Butcher, R. W. & Sutherland, E. W. (1962) *J. Biol. Chem.* **237**, 1244
 Chain, E. B., Mansford, K. R. L. & Opie, L. H. (1969) *Biochem. J.* **115**, 537
 Loten, E. G. & Sneyd, J. G. T. (1970) *Biochem. J.* **120**, 187
 Senft, G., Schultz, G., Munske, K. & Hoffmann, M. (1968) *Diabetologia* **4**, 322
 Sutherland, E. W. & Robinson, A. G. (1969) *Diabetes* **18**, 797
 Thompson, W. J. & Appleman, M. M. (1971) *J. Biol. Chem.* **246**, 3145

Glucose-Mediated Insulin Release from Mouse Islets of Langerhans: Adenylate Cyclase

By B. DAVIS and NORMAN R. LAZARUS (*Diabetes Research Unit, The Wellcome Foundation, Dartford, Kent, U.K.*)

Cyclic AMP (adenosine 3':5'-cyclic monophosphate) has been implicated in mediating insulin release from the β -cells of the pancreas. There is virtually no delay between the arrival of glucose at the β -cell and insulin secretion. It is reasonable to assume therefore that the glucose effect may be mediated by direct interaction with the plasma membrane. Insulin release takes place in two distinct phases (Grodsky *et al.*, 1967; Blackard & Nelson, 1971). The first phase is over within 10min, but the second phase is more prolonged. It is the events underlying this first phase that are the subject of the present investigation. The effects of glucose on mouse islet adenylate cyclase were investigated since it may be expected that this enzyme could be one of the primary points of interaction.

Large numbers of islets (5000–18000) were isolated from mouse pancreas by the collagenase technique. After homogenization, the islets were centrifuged and the 1000g pellet was assayed for adenylate cyclase activity. The pH optimum of the enzyme was found to be 7.6. The apparent K_m was approx. $2 \times 10^{-4}M$ for ATP. The effects of various substances and hormones on the initial velocity of the reaction under the same conditions of enzyme and substrate concentration (0.4mM-ATP) were tested. The cyclase was stimulated by the following substances: NaF (2.4 and 8mM), glucagon (0.3, 0.6 and $1.6\mu M$), pancreozymin (0.15, 0.3 and $0.6\mu M$), secretin (0.3 and $0.6\mu M$). The enzyme was inhibited by Ca^{2+}

(50% inhibition obtained at 0.3 mM). Glucose (2, 8 and 17 mM) had no effect on the activity of the enzyme, nor did galactose at the same concentrations. Groups of islets were incubated separately in two concentrations of glucose (2 and 17 mM) for periods of 5 or 15 min. After incubation islets were harvested and homogenized, and the 1000g pellet was assayed as before for adenylate cyclase activity. No differences between any groups of islets were observed. The following conclusions emerge: the adenylate cyclase of islets is similar in behaviour to other tissue adenylate cyclases. The preparation obtained retained its hormone-sensitive properties. Glucose had no effect on adenylate cyclase activity either *in vivo* or *in vitro*. Likewise galactose was without effect *in vitro*. It is concluded that glucose does not mediate insulin secretion via an effect on adenylate cyclase.

Blackard, W. G. & Nelson, N. C. (1971) *Diabetes* **19**, 302–306

Grodsky, G. M., Bennett, L. L., Smith, D. & Nemecek, K. (1967) *Proc. Brook Lodge Symp.* 11–21

Glucose-Mediated Insulin Release from Mouse Islets of Langerhans: Cyclic Nucleotide Phosphodiesterase

By V. BOWEN and NORMAN R. LAZARUS (*Diabetes Research Unit, The Wellcome Foundation, Dartford, Kent, U.K.*)

Islets (4000–18000) were isolated from mouse pancreas by the collagenase technique in batches. After homogenization the 30000g supernatant was dialysed overnight against 10 mM-NaCl at 4°C and then concentrated by freeze-drying. Cyclic nucleotide phosphodiesterase activity was assayed radiochemically by using cyclic [³H]AMP ([8-³H]adenosine 3':5'-cyclic monophosphate) as substrate, with the precautions recommended by Gulyassy & Oken (1971). The enzyme had a pH optimum of 8.2 and an apparent K_m 0.9×10^{-5} M for cyclic AMP. The addition of Mg^{2+} (10 mM) caused a 5-fold increase in V_{max} . The enzyme is 50% inhibited by 10 mM-theophylline. Enzyme activity is abolished by the addition of 0.5 mM-*p*-chloromercuribenzoate. The activity of the enzyme was unaffected by the addition of 17 mM-glucose or 1 mM-glucose 6-phosphate. Groups of islets were incubated separately with two concentrations of glucose (2 and 17 mM) for periods of 5 or 15 min. After incubation, islets were harvested and homogenized and the enzyme was isolated as before. No decrease in activity was found in samples exposed to 17 mM-glucose. Further, when non-dialysed heated (60°C) supernatant from incubated islets was added to a non-incubated enzyme preparation no difference in enzyme activity was detected. It is concluded

that glucose has no direct inhibitory effect on the phosphodiesterase enzyme. Further, incubation of islets in the presence of glucose likewise does not alter enzyme activity. This further substantiates the view suggested by Montague & Cook (1971) and Atkins & Matty (1971) that glucose-mediated insulin release is independent of the cyclic nucleotide phosphodiesterase system.

Atkins, T. & Matty, A. J. (1971) *J. Endocrinol.* **51**, 67–78
Gulyassy, P. F. & Oken, R. L. (1971) *Proc. Soc. Exp. Med. Biol.* **137**, 361–365

Montague, W. & Cook, J. R. (1971) *Biochem. J.* **122**, 115–120

Oestrogen-Induced Secretion of Egg-Yolk Proteins by Male *Xenopus* Liver in Organ Culture

By M. J. CLEMENS, R. LOFTHOUSE and J. R. TATA (*National Institute for Medical Research, Mill Hill, London N.W.7, U.K.*)

In egg-laying animals synthesis and secretion of yolk proteins by the liver are controlled by oestrogens. There are many reports of the induction of egg-yolk protein synthesis *in vivo* by administration of oestradiol-17 β to the amphibian *Xenopus laevis* (Wallace & Dumont, 1968; Follett *et al.*, 1968; Dolphin *et al.*, 1971), and such a response in male animals must necessarily represent synthesis of new proteins *de novo*. It has been demonstrated that incubated liver slices or cultured liver explants can secrete yolk proteins into the medium (Dolphin *et al.*, 1971; Wallace & Jared, 1969). We have used the technique of organ culture to study the effects of oestradiol on the synthesis of secreted and non-secreted protein by male *Xenopus* liver.

Small pieces of tissue were prepared from livers of adult males that had been injected 3 days previously with either 1 mg of oestradiol-17 β in propylene glycol or the solvent alone. Tissue was cultured in amphibian medium (Wolf & Quimby, 1964) in which $NaHCO_3$ was replaced by 15 mM-HEPES (*N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulphonic acid) buffer, pH 7.5, with added antibiotics. Duplicate cultures were incubated at 18°C in air with [¹⁴C]-leucine or [¹⁴C]-serine, and incorporation of radioactivity into tissue protein and proteins secreted into the medium was assayed.

When [¹⁴C]-serine was the labelled precursor at least 3–4 times as much acid-insoluble radioactivity was secreted into the medium by explants from oestrogen-treated animals, compared with control tissue, for at least 4 days. In contrast very little stimulation of tissue protein labelling was observed. A less dramatic effect on the labelling of secreted protein with [¹⁴C]-leucine was evident. It is probable