(50% inhibition obtained at 0.3 mM). Glucose (2, 8 and 17 mM) had no effect on the activity of the enzyme, nor did galactose at the same concentrations. Groups of islets were incubated separately in two concentrations of glucose (2 and 17 mM) for periods of 5 or 15 min. After incubation islets were harvested and homogenized, and the 1000g pellet was assayed as before for adenylate cyclase activity. No differences between any groups of islets were observed. The following conclusions emerge: the adenylate cyclase of islets is similar in behaviour to other tissue adenylate cyclases. The preparation obtained retained its hormone-sensitive properties. Glucose had no effect on adenylate cyclase activity either *in vivo* or *in vitro*. Likewise galactose was without effect *in vitro*. It is concluded that glucose does not mediate insulin secretion via an effect on adenylate cyclase.

Blackard, W. G. & Nelson, N. C. (1971) *Diabetes* **19**, 302–306

Grodsky, G. M., Bennett, L. L., Smith, D. & Nemecek, K. (1967) *Proc. Brook Lodge Symp.* 11–21

Glucose-Mediated Insulin Release from Mouse Islets of Langerhans: Cyclic Nucleotide Phosphodiesterase

By V. BOWEN and NORMAN R. LAZARUS (*Diabetes Research Unit, The Wellcome Foundation, Dartford, Kent, U.K.*)

Islets (4000–18000) were isolated from mouse pancreas by the collagenase technique in batches. After homogenization the 30000g supernatant was dialysed overnight against 10 mM-NaCl at 4°C and then concentrated by freeze-drying. Cyclic nucleotide phosphodiesterase activity was assayed radiochemically by using cyclic [³H]AMP ([8-³H]adenosine 3':5'-cyclic monophosphate) as substrate, with the precautions recommended by Gulyassy & Oken (1971). The enzyme had a pH optimum of 8.2 and an apparent K_m 0.9×10^{-5} M for cyclic AMP. The addition of Mg^{2+} (10 mM) caused a 5-fold increase in V_{max} . The enzyme is 50% inhibited by 10 mM-theophylline. Enzyme activity is abolished by the addition of 0.5 mM-*p*-chloromercuribenzoate. The activity of the enzyme was unaffected by the addition of 17 mM-glucose or 1 mM-glucose 6-phosphate. Groups of islets were incubated separately with two concentrations of glucose (2 and 17 mM) for periods of 5 or 15 min. After incubation, islets were harvested and homogenized and the enzyme was isolated as before. No decrease in activity was found in samples exposed to 17 mM-glucose. Further, when non-dialysed heated (60°C) supernatant from incubated islets was added to a non-incubated enzyme preparation no difference in enzyme activity was detected. It is concluded

that glucose has no direct inhibitory effect on the phosphodiesterase enzyme. Further, incubation of islets in the presence of glucose likewise does not alter enzyme activity. This further substantiates the view suggested by Montague & Cook (1971) and Atkins & Matty (1971) that glucose-mediated insulin release is independent of the cyclic nucleotide phosphodiesterase system.

Atkins, T. & Matty, A. J. (1971) *J. Endocrinol.* **51**, 67–78
Gulyassy, P. F. & Oken, R. L. (1971) *Proc. Soc. Exp. Med. Biol.* **137**, 361–365

Montague, W. & Cook, J. R. (1971) *Biochem. J.* **122**, 115–120

Oestrogen-Induced Secretion of Egg-Yolk Proteins by Male *Xenopus* Liver in Organ Culture

By M. J. CLEMENS, R. LOFTHOUSE and J. R. TATA (*National Institute for Medical Research, Mill Hill, London N.W.7, U.K.*)

In egg-laying animals synthesis and secretion of yolk proteins by the liver are controlled by oestrogens. There are many reports of the induction of egg-yolk protein synthesis *in vivo* by administration of oestradiol-17 β to the amphibian *Xenopus laevis* (Wallace & Dumont, 1968; Follett *et al.*, 1968; Dolphin *et al.*, 1971), and such a response in male animals must necessarily represent synthesis of new proteins *de novo*. It has been demonstrated that incubated liver slices or cultured liver explants can secrete yolk proteins into the medium (Dolphin *et al.*, 1971; Wallace & Jared, 1969). We have used the technique of organ culture to study the effects of oestradiol on the synthesis of secreted and non-secreted protein by male *Xenopus* liver.

Small pieces of tissue were prepared from livers of adult males that had been injected 3 days previously with either 1 mg of oestradiol-17 β in propylene glycol or the solvent alone. Tissue was cultured in amphibian medium (Wolf & Quimby, 1964) in which $NaHCO_3$ was replaced by 15 mM-HEPES (*N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulphonic acid) buffer, pH 7.5, with added antibiotics. Duplicate cultures were incubated at 18°C in air with [¹⁴C]-leucine or [¹⁴C]-serine, and incorporation of radioactivity into tissue protein and proteins secreted into the medium was assayed.

When [¹⁴C]-serine was the labelled precursor at least 3–4 times as much acid-insoluble radioactivity was secreted into the medium by explants from oestrogen-treated animals, compared with control tissue, for at least 4 days. In contrast very little stimulation of tissue protein labelling was observed. A less dramatic effect on the labelling of secreted protein with [¹⁴C]-leucine was evident. It is probable