Ca¹⁴CO₃ (Swick, 1958), single injection of a large dose of [¹⁴C]lysine (Henshaw *et al.*, 1971), repeated injections of [¹⁴C]glucose (Austin *et al.*, 1972) and constant infusion of tracer doses of labelled amino acids. In this laboratory we favour the last-mentioned method. Results from the infusion method can be used to determine protein synthesis rates in both individual tissues and in the whole animal, and short infusions of a few hours give the same synthesis rates with different amino acids (Waterlow & Stephen, 1968; Garlick, 1969, 1972).

In both uptake and decay experiments mixed samples of protein cannot be treated as a single protein with a turnover rate equal to the mean of the mixture (Koch, 1962). The apparent rate decreases as time progresses. The problem is more serious with the decay rate, since the faster-turning-over components receive more of the pulse dose of label and the early part of the decay curve is biased by these components, so that the slope is too steep. Hence decay experiments cannot be used to obtain a unique value for the mean turnover rate of a mixture. With uptake experiments the rate at zero time is equal to the mean synthesis rate, and therefore short experiments lasting a few hours can give results that are not seriously in error.

Each experimental situation requires careful consideration of the method used to measure turnover rates. The usefulness and validity of the various methods in different situations will be discussed.

- Austin, L., Lowry, O. H., Brown, J. G. & Carter, J. G. (1972) *Biochem. J.* **126**, 351
 Awwad, H. K., Sheraky, A. S., Helmi, S. A., Shetaiwy, S. K. & Potchen, E. J. (1970) *J. Biol. Chem.* **245**, 469
 Berlin, C. M. & Schimke, R. T. (1965) *Mol. Pharmacol.* **1**, 149
 Garlick, P. J. (1969) *Nature (London)* **223**, 61
 Garlick, P. J. (1972) *Biochem. J.* **126**, 23 P
 Henshaw, E. C., Hirsch, C. A., Morton, B. E. & Hiatt, H. H. (1971) *J. Biol. Chem.* **246**, 436
 Koch, A. L. (1962) *J. Theor. Biol.* **3**, 283
 Levin, W. & Kuntzman, R. (1969) *J. Biol. Chem.* **244**, 3671
 McFarlane, A. S. (1963) *Biochem. J.* **89**, 277
 Millward, D. J. (1970) *Clin. Sci.* **39**, 577
 Schimke, R. T. (1964) *J. Biol. Chem.* **239**, 3808
 Schimke, R. T. (1970) in *Mammalian Protein Metabolism* (Munro, H. N., ed.), vol. 4, pp. 178–223, Academic Press, London and New York
 Schoenheimer, R., Ratner, S. & Rittenberg, D. (1939) *J. Biol. Chem.* **127**, 333
 Swick, R. W. (1958) *J. Biol. Chem.* **231**, 751
 Waterlow, J. C. & Stephen, J. M. L. (1968) *Clin. Sci.* **35**, 287

The Pattern of Protein Turnover in the Whole Animal and the Effect of Dietary Variation

By D. J. MILLWARD and P. J. GARLICK (*Clinical Nutrition and Metabolism Unit, Hospital for Tropical Diseases, St. Pancras Way, London NW1 2PE, U.K.*)

In animal cells it is now clear that most if not all proteins are continuously degraded and replaced, i.e. they turn over, and the rate is distinct for each protein. The heterogeneity in turnover within individual tissues is very marked. For example, in the liver δ -aminolaevulinase synthetase and α -glycerophosphate dehydrogenase, both mitochondrial enzymes, are said to have half-lives of 1 h and 4 days respectively (Schimke, 1970). Heterogeneity is also a feature of turnover among different fractions of brain (Austin *et al.*, 1972) and skeletal muscle (Millward, 1970a).

The concentration of a protein in a tissue depends on the balance between its synthesis and degradation, and there is evidence that the control of protein concentrations in the tissues is indeed exerted through both processes and often independently (Schimke & Doyle, 1970). This is especially true for secreted proteins such as albumin, where the site of breakdown is remote from the site of synthesis.

Measurements of whole-body protein turnover in the rat (Waterlow & Stephen, 1967) and in the infant (Picou & Taylor-Roberts, 1969) have shown that under a wide range of nutritional states and dietary intakes the rate does not markedly change, especially in the latter case. We have shown, however, that the pattern of protein turnover in different tissues is susceptible to dietary variation (Millward, 1970b). The rates of synthesis and breakdown of liver and muscle protein were measured in starved rats or rats fed on a protein-low diet. When the total synthesis and catabolism in the two tissues were calculated, it was shown that the low-protein diet caused a shift in synthesis towards liver and an opposite shift in catabolism, whereas starvation altered synthesis and catabolism proportionately in each tissue.

The pattern of total protein synthesis between the tissues varies with age. In muscle tissue the replacement rate is low and therefore, in the young animal during growth, half of the total synthesis represents net growth (the rest being turnover or replacement) (Millward, 1970b). In the adult, growth is much diminished and so total muscle protein synthesis may be halved (Waterlow & Stephen, 1968). By contrast, in rapidly-turning-over tissues such as liver, the growth component or rate of net synthesis is relatively small compared with the rate of turnover or replacement, so that when growth ceases there is little change in the total synthesis rate. The result is that the muscles' share of total body protein synthesis is lower in the adult than in the young animal.

The intensity of protein synthesis in different tissues corresponds to the concentrations of RNA in the tissues. Thus in cardiac muscle the rate of protein synthesis and the concentration of RNA are in each case higher than in skeletal muscle. Changes in RNA concentration can occur very rapidly in response to food intake. We have shown, for example, that, in liver, heart and skeletal muscle of rats on cyclic feeding schedules, periodic changes in RNA concentration occur in all three tissues. That such changes can effect a change in protein synthesis is shown by parallel alterations in the rate of muscle protein synthesis and RNA concentration. Also, the loss of RNA that occurs when food is withdrawn is equally rapid from liver, heart and skeletal muscle.

The control of protein breakdown in different tissues is in no way understood. Changes in the amount of dietary protein can cause immediate changes in either direction in the breakdown rate of arginase, whereas the breakdown rate of albumin responds only slowly to changes in dietary protein. A decrease in the rate of breakdown represents an energetically economical way of increasing the mass of protein. Evidence that such a mechanism occurs has been reported by Young *et al.* (1971) during the rapid catch-up growth of skeletal muscle that occurs on re-feeding after protein depletion. We have reported decreases in the specific activities of two proteolytic enzymes in rat skeletal muscle during similar periods of catch-up growth (Millward, 1972).

It is therefore obvious that protein breakdown and synthesis are equally important in controlling the concentration of tissue protein. These results and current work on the problem will be discussed.

- Austin, L., Lowry, O. H., Brown, J. G. & Carter, J. G. (1972) *Biochem. J.* **126**, 351
 Millward, D. J. (1970a) *Clin. Sci.* **39**, 577
 Millward, D. J. (1970b) *Clin. Sci.* **39**, 591
 Millward, D. J. (1972) *Proc. Nutr. Soc.* **31**, 3A
 Picou, D. & Taylor-Roberts, T. (1969) *Clin. Sci.* **36**, 283
 Schimke, R. T. (1970) in *Mammalian Protein Metabolism* (Munro, H. N., ed.), vol. 4, pp. 178–223, Academic Press, London and New York
 Schimke, R. T. & Doyle, D. (1970) *Annu. Rev. Biochem.* **39**, 929
 Waterlow, J. C. & Stephen, J. M. L. (1967) *Clin. Sci.* **33**, 489
 Waterlow, J. C. & Stephen, J. M. L. (1968) *Clin. Sci.* **35**, 287
 Young, V. R., Stothers, S. C. & Vilaire, G. (1971) *J. Nutr.* **101**, 1379

Measurement of the Synthesis of Liver-Produced Plasma Proteins with Particular Reference to Dietary Protein and Amino Acid Supply

By R. HOFFENBERG (*Division of Metabolism, Clinical Research Centre, Watford Road, Harrow, Middx. HA1 3UJ, U.K.*)

Production of plasma proteins by the liver can be measured *in vivo* (a) indirectly, by following the metabolism of injected radioactively labelled protein molecules, or (b) directly, by the [^{14}C]carbonate method. *In vitro*, use may be made of the isolated perfused liver, liver slices or isolated liver cells, and a direct assay of protein production by immunological means can be applied. The advantages and disadvantages of the various techniques will be discussed, with particular reference to the adaptation of albumin production to protein or amino acid deprivation. *In vivo*, it can be shown that albumin synthesis falls rapidly in the face of such deprivation, and a similar response can be demonstrated with the isolated perfused liver or isolated liver cells removed from rats subjected to a low-protein diet. These findings will be discussed in relation to changes in liver mass, cell size and polyribosome structure.

Influence of Amino Acid Supply on Protein Synthesis by Rat Liver *in vitro*

By MICHAEL J. CLEMENS (*National Institute for Medical Research, Mill Hill, London NW7 1AA, U.K.*)

The rate of protein synthesis in rat liver is sensitive to changes in the availability of amino acids. This has been shown *in vivo* (Wunner *et al.*, 1966), in a perfusion system (Jefferson & Korner, 1969) and with a cell-free preparation (Baliga *et al.*, 1968). I have studied the effects of changes in amino acid supply in a liver-slice incubation system and have shown that the protein-synthetic activity of cell-free systems subsequently prepared from the incubated tissue is elevated after exposure to high concentrations of amino acids (Clemens & Korner, 1971). Further, this effect is localized in the cell-sap fraction of the liver and may represent an effect of amino acids distinct from the stimulation of polyribosome formation reported by others.

Liver slices were prepared from male Sprague-Dawley rats (100–120g) that were either fed *ad lib.* until death or starved overnight (16h). The tissue was incubated for 1h in a medium containing either normal plasma concentrations of 20 amino acids or six times these concentrations (Clemens & Korner, 1970). Post-mitochondrial supernatants, ribosomes