Nucleic Acid Precursor Uptake into Normal and Regenerating Rat Liver

By MARGERY G. ORD and LLOYD A. STOCKEN
(Department of Biochemistry, University of Oxford,
Oxford OX1 3QU, U.K.)

Growth-promoting hormones may alter precursor uptake into their target tissues (Tata, 1970), and in cell cultures uptake may be promoted by factors from serum (Cunningham & Pardee, 1969). In regenerating rat liver UTP concentrations are increased by 50% within 3 h after partial hepatectomy (Bucher & Swaffield, 1969). Experiments with [^3H]-orotate, -thymidine and -uridine, with [$\text{carboxy-}^{14}\text{C}$]inulin as marker for extracellular space, indicate twofold increases in intracellular acid-soluble ^3H radioactivity in the residual livers 1 h after partial hepatectomy. This response was not found in pancreas, thymus, salivary gland or muscle, which also differ from liver by their failure, with [^3H]orotate, to concentrate ^3H radioactivity intracellularly with respect to plasma. Actinomycin D (0.55 mg/kg body wt.) decreased total nuclear RNA synthesis in the livers of partially hepatectomized rats by 50%, but did not prevent the increased precursor uptake at 1 h. [^3H]Adenine distribution between liver and plasma did not alter after partial hepatectomy. Liver is sensitive to surgical stress (Moolten *et al.*, 1970), but neither unilateral nephrectomies nor sham operations affected [^3H]orotate uptake.

The peak of DNA synthesis after partial hepatectomy in our rats occurs at 21–22 h. [^3H]Thymidine uptake into the intracellular acid-soluble compartment in liver at this time is still doubled, although that for [^3H]orotate is now only 30% above values for sham-operated animals.

Work with cell cultures indicates that uridine transport shows saturation kinetics and that permeation is the rate-limiting step for the incorporation of uridine into the intracellular pool (Plagemann & Roth, 1969). The uptake of [^3H]orotate into intact livers is linearly related to the administered dose up to about 2 mg/kg body wt., and the ratio of intracellular acid-soluble ^3H radioactivity to that of plasma did not exceed 6.8:1, in contrast with the ratio 10–12:1 for the partially hepatectomized animals. These findings are consistent with a carrier-mediated mechanism for the transport of orotate. The possibilities that this mechanism may be promoted by a humoral factor, which is increased by partial hepatectomy, and that the increased penetration of precursors plays a role in the initial stages of liver regeneration will be discussed.

Bucher, N. L. R. & Swaffield, M. N. (1969) *Biochim. Biophys. Acta* **174**, 491–502

Cunningham, D. S. & Pardee, A. B. (1969) *Proc. Nat. Acad. Sci. U.S.A.* **64**, 1049–1056
Moolten, F. L., Oakman, N. J. & Bucher, N. L. R. (1970) *Cancer Res.* **30**, 2353–2357
Plagemann, P. G. W. & Roth, M. F. (1969) *Biochemistry* **8**, 4782–4789
Tata, J. R. (1970) in *Biochemical Actions of Hormones* (Litwak, G., ed.), vol. 1, pp. 89–133, Academic Press New York and London

α -Adrenergic Blockers and Early Stages in Liver Regeneration

By SEAN THROWER and MARGERY G. ORD (Department of Biochemistry, University of Oxford, Oxford OX1 3QU, U.K.)

Adrenalectomy does not prevent liver regeneration in partially hepatectomized rats (Ferrari & Harkness, 1954), but both catecholamines and glucocorticoids are known to facilitate the induction of some liver enzymes, including ornithine decarboxylase (Panko & Kenney, 1971), which is markedly increased early in liver regeneration (Jänne & Raina, 1968; Russell & Snyder, 1968).

Preliminary experiments in this laboratory established that propranolol, a β -adrenergic blocker, did not affect regeneration, but that phenoxybenzamine, an α -adrenergic blocker, did. A dose of 5 mg of phenoxybenzamine/kg body wt. given immediately before operation prevented the normal induction of ornithine decarboxylase; the effect, which was transient, delayed the peak of induction by 2 h. Under these conditions no effects were detected on DNA synthesis. Adrenalectomized rats behaved similarly to phenoxybenzamine-treated animals, showing only 26% of the usual decarboxylase activity at 4 h but no change in thymidine incorporation into DNA at 21–24 h.

If phenoxybenzamine was given 5–10 h after operation, after decarboxylase induction had taken place, the peak of DNA synthesis was affected. A dose of 5 mg of phenoxybenzamine/kg body wt. given 9 h after operation shifted the peak from 21.5 to 24.5 h. Administration of the drug 1 h before the peak had no effect, indicating that phenoxybenzamine does not act directly on the enzymes involved in DNA synthesis. Nuclear histone phosphokinase (Siebert *et al.*, 1971) is typical of the enzymes induced 12–15 h after partial hepatectomy; this induction was delayed by a dose of 5 mg of phenoxybenzamine/kg body wt. given at 9 h, suggesting that catecholamines facilitate not only the early wave of enzyme induction in regenerating liver but also that of enzymes, particularly those concerned with replication, that are induced later in G_1 phase.

Phenoxybenzamine did not alter the increased pyrimidine uptake detected 1 h after partial hepatectomy

(Ord & Stocken, 1972), confirming that this membrane effect is not dependent on catecholamines.

- Ferrari, V. & Harkness, R. D. (1954) *J. Physiol. (London)* **124**, 443–463
 Jänne, J. & Raina, A. (1968) *Acta Chem. Scand.* **22**, 1349–1351
 Panko, W. B. & Kenney, F. T. (1971) *Biochem. Biophys. Res. Commun.* **43**, 346–350
 Russell, D. H. & Snyder, S. H. (1968) *Proc. Nat. Acad. Sci. U.S.A.* **60**, 1420–1427
 Siebert, G., Ord, M. G. & Stocken, L. A. (1971) *Biochem. J.* **122**, 721–725
 Thrower, S., Ord, M. G. & Stocken, L. A. (1972) *Biochem. Pharmacol.* in the press

Prolactin-Stimulated Deoxyribonucleic Acid Synthesis in Rabbit Mammary Tissue

By R. A. BOURNE, J. A. BRYANT, D. GRIERSON and I. R. FALCONER* (*University of Nottingham School of Agriculture, Sutton Bonington, Loughborough, Leics. LE12 5RD, U.K.*)

In an earlier communication Fiddler & Falconer (1969) reported that prolactin stimulated the incorporation of thymidine into DNA in rabbit mammary gland. This work has been extended with the aim of investigating the role of DNA synthesis in cyto-differentiation in the mammary gland.

Pseudo-pregnancy was induced as described previously (Fiddler & Falconer, 1969). At 12 days after the induction of pseudo-pregnancy an individual mammary gland was injected with prolactin (125 µg). Further glands were injected at 6h intervals. At 24h after the first prolactin injection all injected glands and one un-injected gland were excised. Tissue slices from the glands were incubated for 2h in [Me^3H]thymidine (1 µCi/ml). DNA was extracted from the tissue by a method based on that of Munro & Fleck (1966). Measurement of the specific radioactivity of the DNA showed that prolactin markedly stimulated the incorporation of thymidine into DNA, with the greatest stimulation occurring 18–24h after the injection. Addition of chloramphenicol (25 µg/ml) and streptomycin (25 µg/ml) to the incubation medium had no effect on the incorporation of thymidine into DNA. The incorporation of thymidine in both prolactin-stimulated and unstimulated tissue was inhibited by hydroxyurea (50 µg/ml).

Radioautographic studies indicated that prolactin caused an increase in the number of cells that incorporated thymidine into their nuclei. Further, prolactin-stimulated tissue that had been treated with

colchicine contained more arrested metaphases than did unstimulated tissue. These results suggest that prolactin stimulates DNA synthesis and cell division in mammary tissue.

Partial characterization of the radioactive DNA from prolactin-stimulated and unstimulated tissue has been carried out. The DNA was extracted by a detergent–chloroform method and centrifuged in a CsCl density gradient. The gradients were fractionated and the specific radioactivity of each fraction was determined. In prolactin-treated tissue the DNA at the heavy side of the peak, corresponding in position to the genes coding for rRNA (Birnstiel *et al.*, 1972), was much more heavily labelled than was the bulk of the DNA. In unstimulated tissue the same region of the peak was slightly more heavily labelled than was the bulk of the DNA. This suggests that prolactin may also stimulate amplification of the cistrons coding for rRNA. This suggestion is currently being tested by using hybridization techniques.

We thank Dr. Margaret Birkinshaw for help with the radioautography.

- Birnstiel, M. L., Sells, B. H. & Purdom, I. F. (1972) *J. Mol. Biol.* **63**, 21–39
 Fiddler, T. J. & Falconer, I. R. (1969) *Biochem. J.* **115**, 58P–59P
 Munro, H. N. & Fleck, A. (1966) *Methods Biochem. Anal.* **14**, 113

Role of Proteins in the Condensation of Hen Erythrocyte Chromatin

By MICHAEL A. BILLETT and J. MICHAEL BARRY (*Department of Agricultural Science, University of Oxford, Oxford OX1 3PF, U.K.*)

Chromatin in hen erythrocytes is condensed and genetically inactive, whereas rat liver chromatin contains both inactive condensed and active uncondensed fractions. Chromatin isolated from erythrocytes is swollen in media of low cation concentration, but condenses as the unbound cation concentration is raised to the values that occur in the intact cell (Leake *et al.*, 1972). We report here some observations on the influence of chromatin proteins on this condensation.

Proteins were extracted from hen erythrocyte chromatin or nuclei with increasing concentrations of NaCl. They were separated from the residual chromatin either by centrifugation or by gel filtration on Sepharose 4B. Proteins were extracted from the residual chromatin with 0.25M-HCl. All protein fractions were characterized by polyacrylamide-gel electrophoresis (Johns, 1967). The proportion of condensed chromatin at different Mg^{2+} concentrations was determined by its sedimentation on low-speed centrifugation (Leake *et al.*, 1972).

* Present address: Department of Applied Biochemistry and Nutrition, University of New England, Armidale, N.S.W., Australia.