ether analogue of griseofulvin, which is not porphyrogenic, does not inhibit the activity of the chelatase.

The stimulation of 5-aminolaevulinatase synthetase activity by both 2-allyl-2-isopropylacetamide and 3,5-diethoxycarbonyl-1,4-dihydrocollidine can be prevented by prior administration to the rat of either compound SKF 525-A (2-diethylaminoethyl 3,3-diphenylpropylacetate) or cycloheximide. An important difference is that compound SKF 525-A also prevents the destruction of liver haem by 2-allyl-2-isopropylacetamide and the inhibition of the chelatase by 3,5-diethoxycarbonyl-1,4-dihydrocollidine, whereas cycloheximide does not (De Matteis, 1971; F. De Matteis, G. Abbritti & A. Gibbs, unpublished work). The significance of these findings and the possible mechanism of action of the two inhibitors will be discussed in the light of the hypothesis (Granick, 1966) that the stimulation of the synthetase caused by drugs involves synthesis *de novo* of the enzyme protein.

In clear contrast with the porphyrogenic compounds considered so far, a second group of drugs, including phenobarbitone and phenylbutazone, increases rather than decreases the concentration of cytochrome *P*-450 and haem in the microsomal fraction without causing accumulation of porphyrins in the liver. The small stimulation of 5-aminolaevulinatase synthetase activity caused by these drugs possibly reflects an increased haem utilization rather than an interference with the regulation of the pathway. However, when either phenobarbitone or phenylbutazone is given together with a small dose of 3,5-diethoxycarbonyl-1,4-dihydrocollidine, the stimulation of 5-aminolaevulinatase synthetase activity caused by this latter drug is greatly enhanced (De Matteis, 1972). The following interpretation is possible for this potentiation effect. 3,5-Diethoxycarbonyl-1,4-dihydrocollidine may, by inhibiting the synthesis of liver haem, decrease the concentration of haem available for control of the synthetase. Phenobarbitone and phenylbutazone, by promoting an accumulation of cytochrome *P*-450 apoprotein, may stimulate haem utilization and thereby decrease the concentration of this 'regulatory' haem still further.

- Abbritti, G. & De Matteis, F. (1971-72) *Chem.-Biol. Interact.* **4**, 281-286
 De Matteis, F. (1971) *Biochem. J.* **124**, 767-777
 De Matteis, F. (1972) *Biochem. J.* **127**, 21P-22P
 De Matteis, F. & Gibbs, A. (1972) *Biochem. J.* **126**, 1149-1160
 Gibson, K. D., Neuberger, A. & Scott, J. J. (1955) *Biochem. J.* **61**, 618-629
 Granick, S. (1966) *J. Biol. Chem.* **241**, 1359-1375
 Granick, S. & Sassa, S. (1971) *Metab. Pathways* **5**, 77
 Granick, S. & Urata, G. (1963) *J. Biol. Chem.* **238**, 821-827
 Hayashi, N., Yoda, B. & Kikuchi, G. (1969) *Arch. Biochem. Biophys.* **131**, 83-91
 Lockhead, A. C., Dagg, J. H. & Goldberg, A. (1967) *Brit. J. Dermatol.* **79**, 96-102

- Marver, H. S., Collins, A., Tschudy, D. P. & Rechcigl, M. (1966) *J. Biol. Chem.* **241**, 4323-4329
 Meyer, U. A. & Marver, H. S. (1971) *Science* **171**, 64-66
 Miyagi, K., Cardinal, R., Bossenmaier, I. & Watson, C. J. (1971) *J. Lab. Clin. Med.* **78**, 683-695
 Nakao, K., Wada, O., Takaku, F., Sassa, S., Yano, Y. & Urata, G. (1967) *J. Lab. Clin. Med.* **70**, 923-932
 Onisawa, J. & Labbe, R. F. (1962) *Biochim. Biophys. Acta* **56**, 618-620
 Strand, L. J., Felsher, B. F., Redeker, A. G. & Marver, H. S. (1970) *Proc. Nat. Acad. Sci. U.S.A.* **67**, 1315-1320
 Tschudy, D. P. & Bonkowski, H. L. (1972) *Fed. Proc. Fed. Amer. Soc. Exp. Biol.* **31**, 147-159

Mechanisms and Consequences of the Induction of Microsomal Enzymes of Mammalian Liver

By DENNIS V. PARKE (*Department of Biochemistry, University of Surrey, Guildford, Surrey, U.K.*)

The mixed-function oxidase system present in the endoplasmic reticulum of mammalian liver is responsible, through its terminal oxygenase cytochrome *P*-450, for the oxidative metabolism of cholesterol, steroids, fatty acids, drugs and other xenobiotics (Parke, 1968). The induction of this enzyme system has been widely studied and may be effected by a great diversity of drugs, steroids, carcinogens and other foreign compounds (Conney, 1967; Parke, 1971a). Several theoretical models for translational and transcriptional control of the relevant gene expression have been examined (Venkatesan *et al.*, 1971). An important criterion for induction, at least at the translational level, would appear to be that the inducing agent is a substrate of this microsomal enzyme system. One of the major physiological roles of this enzyme system is undoubtedly the oxidative metabolism of anutrient xenobiotics of the diet, thus facilitating their elimination from the body. It might therefore be expected that natural dietary anutrients would also be effective inducing agents of the microsomal mixed-function oxidase system, and a number of such compounds, such as β -ionone and safrole, have been shown to induce these enzymes (Parke & Rahman, 1969, 1970). If enzyme induction is primarily an adaptive process, it might be expected that this most versatile enzyme system with its wide diversity of substrates would be among the foremost of mammalian enzymes to manifest induction, and, with the present widespread use of synthetic chemicals such as drugs, food additives, pesticides etc., this induction would have high potential and might exhibit certain abnormalities.

One of the most widely studied inducing agents of this enzyme system is the barbiturate phenobarbital. Studies of the inducing potential of a series of barbiturates, including pentobarbital, secobarbital, allobarbitol, thiopental and phenobarbital, have shown

that the extent of induction of cytochrome *P*-450 and various drug-metabolizing enzyme activities is directly related to the plasma half-lives of these drugs (Parke, 1971*b*). This suggests that induction of this enzyme system is greater for substrates that are more slowly metabolized, and hence are present in the hepatic endoplasmic reticulum, and probably in contact with the enzyme, for the longest periods. The induction by phenobarbital involves the synthesis *de novo* of enzyme proteins, including cytochrome *P*-450 (Conney & Gilman, 1963; Kato *et al.*, 1965; Orrenius *et al.*, 1965), but the increase in the phospholipid component of the endoplasmic reticulum is attributed to a decrease in turnover (Holtzman & Gillette, 1966). Induction by phenobarbital may also affect the nature of cytochrome *P*-450 or its micro-environment, causing qualitative changes in the ligand interactions with substrates (von Bahr *et al.*, 1972) and manifesting latent enzymic activity (Cram *et al.*, 1965).

The induction of cytochrome *P*-450 and the hepatic microsomal drug-metabolizing enzymes by β -ionone also involves synthesis *de novo*, but the increases in enzymic activities that result from pretreatment with safrole and isosafrole are due only in part to increased enzyme synthesis. With the substrate biphenyl it has been shown that hydroxylation in the 4-position may be induced by both drugs and carcinogenic polycyclic hydrocarbons, whereas hydroxylation in the 2-position is increased primarily by the carcinogens (Creaven & Parke, 1966). Safrole, a component of oil of sassafras and other essential oils and flavours, is a weak hepatocarcinogen (Epstein *et al.*, 1970). Administration to rats results in a marked increase in cytochrome *P*-450 and in the 4-hydroxylation of biphenyl, both of which increases are inhibited by actinomycin D and thioacetamide, inhibitors of protein synthesis, and by 3-amino-1,2,4-triazole, an inhibitor of haem synthesis. An even more marked increase occurs in the 2-hydroxylation of biphenyl, but only part of this increased activity appears to be due to increased synthesis of enzyme protein. Like other carcinogens, safrole and isosafrole also induce the activity of benzopyrene hydroxylase in extra-hepatic tissues (Lake & Parke, 1972*b*).

In the induction of the hepatic microsomal mixed-function oxidase system by 3-methylcholanthrene and other carcinogens, certain fundamental differences from phenobarbital induction have been observed in the nature of the terminal cytochrome and the mechanism of induction (Alvares *et al.*, 1967, 1968; Bresnick & Mosse, 1969). In the induction by safrole and other methylenedioxyaryl compounds, such as the insecticide synergist piperonyl butoxide, evidence for a new or modified cytochrome has been obtained that may result from the binding of a metabolite to cytochrome *P*-450 (Parke & Rahman, 1971; Franklin, 1971). This binding to the cytochrome

modifies the enzymic characteristics (Lake & Parke, 1972*a*), so that despite large increases in enzyme protein the increases in enzymic activity are at first modest and later decline below normal (Gray *et al.*, 1972). A comparative study of the long-term administration to rats of safrole and butylated hydroxytoluene (BHT), a food antioxidant and another inducer of these enzymes, has shown that, whereas safrole results in the formation of substantial amounts of the modified cytochrome *P*-450 with little or no increase in activity of the various hydroxylating enzymic activities, butylated hydroxytoluene results in a continuous induction of normal cytochrome *P*-450 and the expected accompanying increase in enzymic activity (Gray *et al.*, 1972). This modification of the cytochrome *P*-450 by methylenedioxyaryl compounds may therefore be responsible for the inhibitory effects that lead to their use as pesticide synergists, and may also be the cause of the hepatotoxic effects of safrole and isosafrole. Similar binding to cytochrome *P*-450 has also been suggested for carcinogenic hydrocarbons (Shoeman *et al.*, 1969), and it is possible, with the diversity of reactive groups to be found in the many synthetic chemicals in current use, that other substances, possibly drugs, might similarly interact with the endoplasmic reticulum to modify the processes of detoxication and alter the normal processes of deactivation of drugs and toxic chemicals.

Enzyme induction is responsible for the tolerance developed with barbiturates and other drugs (Parke, 1971*b*), and may give rise to interactions between drugs that may result in increased metabolism and decreased pharmacological activity (Parke, 1972). Enzyme-inducing agents such as phenobarbital and *N*-phenylbarbital have also been used to enhance the glucuronyltransferase of the hepatic endoplasmic reticulum in the treatment of congenital non-haemolytic unconjugated hyperbilirubinaemia (Crigler & Gold, 1966). A better understanding of the mechanisms of induction of the drug-metabolizing enzymes would benefit our knowledge of pharmacokinetics and drug interactions, and might also lead to the design of more effective therapeutic inducing agents and pesticide synergists.

- Alvares, A. P., Schilling, G. R., Levin, W. & Kuntzman, R. (1967) *Biochem. Biophys. Res. Commun.* **29**, 521
 Alvares, A. P., Schilling, G. R. & Kuntzman, R. (1968) *Biochem. Biophys. Res. Commun.* **30**, 588
 Bresnick, E. & Mosse, H. (1969) *Mol. Pharmacol.* **5**, 219
 Conney, A. H. (1967) *Pharmacol. Rev.* **19**, 317
 Conney, A. H. & Gilman, A. G. (1963) *J. Biol. Chem.* **238**, 3682
 Cram, R. L., Juchau, M. R. & Fouts, J. R. (1965) *Proc. Soc. Exp. Biol. Med.* **118**, 872
 Creaven, P. J. & Parke, D. V. (1966) *Biochem. Pharmacol.* **15**, 7
 Crigler, J. F., Jr. & Gold, N. I. (1966) *J. Clin. Invest.* **48**, 42

- Epstein, S. S., Fujii, K., Andrea, J. & Mantel, N. (1970) *Toxicol. Appl. Pharmacol.* **16**, 321
- Franklin, M. R. (1971) *Xenobiotica* **1**, 581
- Gray, T., Parke, D. V., Grasso, P. & Crampton, R. F. (1972) *Biochem. J.* **130**, 91P
- Holtzman, J. L. & Gillette, J. R. (1966) *Biochem. Biophys. Res. Commun.* **24**, 639
- Kato, R., Loeb, L. & Gelboin, H. V. (1965) *Biochem. Pharmacol.* **14**, 1164
- Lake, B. G. & Parke, D. V. (1972a) *Biochem. J.* **127**, 23P
- Lake, B. G. & Parke, D. V. (1972b) *Biochem. J.* **130**, 86P
- Orrenius, S., Ericson, J. E. & Ernster, L. (1965) *J. Cell Biol.* **25**, 627
- Parke, D. V. (1968) *Int. Ser. Monogr. Pure Appl. Biol. (The Biochemistry of Foreign Compounds)*, Pergamon Press, Oxford
- Parke, D. V. (1971a) in *Effects of Drugs in Cellular Control Mechanisms* (Rabin, B. R. & Freedman, R. B. eds.), p. 69, Macmillan, London
- Parke, D. V. (1971b) in *Acute Barbiturate Poisoning* (Matthew, H. ed.), p. 7, Excerpta Medica Foundation, Amsterdam
- Parke, D. V. (1972) *Roy. Soc. Health J.* **92**, 65
- Parke, D. V. & Rahman, H. (1969) *Biochem. J.* **113**, 12P
- Parke, D. V. & Rahman, H. (1970) *Biochem. J.* **119**, 53
- Parke, D. V. & Rahman, H. (1971) *Biochem. J.* **123**, 9P
- Shoeman, D. W., Chaplin, M. D. & Mannering, G. J. (1969) *Mol. Pharmacol.* **5**, 412
- Venkatesan, N., Arcos, J. C. & Argus, M. F. (1971) *J. Theor. Biol.* **33**, 517
- von Bahr, C., Schenkman, J. B. & Orrenius, S. (1972) *Xenobiotica* **2**, 89

COMPUTER AND INSTRUMENTATION GROUP COLLOQUIUM ON 'CHROMATOGRAPHIC EQUIPMENT'

Trends in Chromatographic Equipment

By E. REID (*Wolfson Bioanalytical Centre, University of Surrey, Guildford, Surrey, U.K.*)

An outlay of £1000 or more on g.l.c. (vapour-phase chromatography) equipment is a good investment for any laboratory that has to analyse vaporizable substances, the technique being notable for resolving power if not speed. The following contribution, by G. Guiochon, outlines present-day trends in g.l.c. equipment. Although the usefulness of g.l.c. apparatus is enormously enhanced by coupling it to a mass spectrometer, this entails expense that, even if it becomes less formidable during the next few years, will remain prohibitive to most laboratories. Another approach, still with few practitioners, is to interface g.l.c. with t.l.c.; this may not facilitate actual measurement, but it does help with identification or with resolving a mixed peak into its components (Kaiser, 1969).

Already g.l.c. has proved useful for preparative as well as for analytical work. This may also be the case in future for high-pressure chromatography of liquids. For the latter up-and-coming technique a satisfactory name is badly needed in place of the unpardonable term 'liquid chromatography' that has found its way into sales literature. To the uninitiated this term might signify conventional column chromatography such as has been practised for decades. Even where literature does emphasize the high-pressure aspect, and the superior resolution and speed achievable, seldom is it pointed out that the key feature of the technique is the use of micro-granular column materials, through which the flow of liquid would be hopelessly slow if merely under gravity. For high performance a daunting outlay of more than

£5000 has to be made. Fair performance may, however, be achievable with medium-pressure apparatus such as is now appearing on the market, sometimes glibly termed 'low-cost' (actual cost £1000-2000); components for do-it-yourself assemblies are also becoming available.

Gravity-flow column chromatography will continue to be the method of choice for many workers, on the grounds of cheapness and/or convenience; indeed, high pressure must be avoided if present-day molecular-sieve materials are to be used. The fraction collector represents the only significant outlay, and need cost no more than £100 if of turntable type rather than modular-rack type; manufacturers are still introducing new models into what seems to be a saturated market. With overnight running, and possibly use of an automatic solvent-changing device, the slowness of gravity flow is often no handicap.

Speed of operation may likewise, in some contexts, be a minor factor when it comes to choosing between paper and thin-layer materials. One advantage of paper is that it stays intact when elution of separated spots is performed. Yet t.l.c. is nowadays the method of choice except where paper chromatography has an over-riding advantage for a particular reason. Centrifugal or other methods to speed up t.l.c. seem to be 'gilding the lily'. The tedious stages in layer chromatography are commonly the processes of applying the material (particularly where multiple spotting has to be done) and of measuring the amounts of the separated components quantitatively. For the latter purpose scanning devices that obviate elution are deservedly becoming popular, although users should beware of unsuspected errors in the readings obtained.

'On-line' quantitative measurement may come to occupy a limited but useful place in the t.l.c. analysis