Rat Liver Mitochondrial Monoamine Oxidase, a Flavin-Containing Enzyme

By M. B. H. YODIM. (*Bernhard Baron Memorial Research Laboratories and Institute of Obstetrics and Gynaecology, Queen Charlotte's Maternity Hospital, London W.6, U.K.*) and T. L. SOURKES. (*Laboratory of Chemical Neurobiology, Department of Psychiatry, McGill University, Montreal, Que., Canada*)

Mitochondrial monoamine oxidase [monoamine-oxygen, oxidoreductase (deaminating), EC 1.4.3.4] of rat liver was purified (Youdim & Sourkes, 1966). The enzyme was pale yellow in colour. When the difference absorption spectrum (oxidized minus reduced) was determined, peaks at 405 and 460 nm appeared. The absorption at 460 nm suggests a flavin moiety. Erwin & Hellerman (1967) reported a similar finding with bovine kidney enzyme. The fluorescent emission and activation spectra of the purified enzyme resembles that of authentic FAD. However, some preparations of the enzyme did not fluoresce, probably owing to the impurity of the enzyme. Treatment of the enzyme with 10% (w/v) trichloroacetic acid resulted in the release of a small amount of flavin-like material. If the purified preparation (1.23 mg/ml) is treated with pronase (1.5 mg/ml) in 0.05 M-phosphate buffer, pH 7.4, at 37°C, then a material fluorescing at 520 nm (activation at 460 nm) is found. The release of this fluorescent material is progressive in the course of digestion with pronase over 4 h. There are losses of fluorescence after 8 h (H. Birnbaum, M. B. H. Youdim & T. L. Sourkes, unpublished work; cf. Sourkes, 1968). This fluorescent material is identical with FAD as regards fluorescent emission and activation. One mol of flavin is associated with 150 000 g of the enzyme. Similar results, starting with enzyme purified from organs of other species, have been reported by Erwin & Hellerman (1967), Tipton (1968) and Gomes, Igaue, Kloeppfer & Yasunobu (1969). The flavin obtained from pronase-treated enzyme and standard solutions of authentic FAD and FMN were subjected to silica-gel t.l.c. in a dark room. After development in butanol-acetic acid-water (3:2:1, by vol.) the chromatogram was dried and examined under u.v. light. The flavin extracted from enzyme showed a major fluorescent spot, with some trailing, behind FAD. No fluorescent material running with FMN could be detected. When the plate was treated with ninhydrin, the enzyme-extracted flavin spot gave a positive reaction, whereas FAD and FMN did not.

The treatment of rats with galactoflavin (2 mg/kg of diet), an anti-metabolite of riboflavin, up to 42 days resulted in the loss of monoamine oxidase

activity in the liver and brain of about 81 and 33% respectively.

From these observations it is concluded that rat liver monoamine oxidase is associated with FAD.

We thank the Medical Research Council of Canada and the Wellcome Trust, U.K., for their support in this project.

Erwin, V. G. & Hellerman, L. (1967). *J. biol. Chem.* **242**, 4230.

Gomes, B., Igaue, I., Kloeppfer, H. G. & Yasunobu, K. T. (1969). *Archs Biochem. Biophys.* **132**, 16.

Sourkes, T. L. (1968). *Adv. Pharmac.* **6A**, 61.

Tipton, K. F. (1968). *Biochim. biophys. Acta*, **159**, 451.

Youdim, M. B. H. & Sourkes, T. L. (1966). *Can. J. Biochem.* **44**, 1397.

The Kinetic Properties of Human Placental Choline Acetyltransferase

By D. MORRIS* and C. O. HEBB. (*Agricultural Research Council Institute of Animal Physiology, Babraham, Cambridge, U.K.*)

Previous studies on the mechanism of choline acetyltransferase (EC 2.3.1.6) have yielded conflicting results. Schuberth (1966) obtained kinetics with human placental enzyme characteristic of a 'Ping Pong' mechanism in which a central acetyl-enzyme complex is formed from acetyl-CoA (in the forward direction) or from acetylcholine (in the reverse direction) independently of the presence of the second substrate (choline or CoA). Potter, Glover & Saelens (1968), on the other hand, obtained evidence of a sequential mechanism for the rat brain enzyme in which both substrates must interact with the enzyme before any products are released.

A study of isotopic exchange between acetylcholine and [*Me*-¹⁴C]choline catalysed by the placental enzyme (Morris & Grewaal, 1969) showed that the 'Ping Pong' mechanism could not be correct since the exchange occurred only in the presence of CoA, a result consistent with a 'sequential' mechanism. The present work is a reinvestigation of the kinetic behaviour of the placental enzyme, which has been extended to include product-inhibition studies.

Choline acetyltransferase activity was assayed by using [*acetyl*-¹⁴C]CoA, by a procedure (D. Morris, A. Maneckjee & C. O. Hebb, unpublished work) modified from that described by McCaman & Hunt (1965) and Schrier & Shuster (1967). Double-reciprocal plots of the initial velocity of acetylcholine synthesis versus the concentration of one

* Present address: Department of Physiology and Biochemistry, University of Southampton, Southampton SO9 5NH, U.K.

substrate at different fixed concentrations of the second substrate gave convergent lines intersecting to the left of the ordinate and below the abscissa. These results are characteristic of a sequential mechanism in which the Michaelis constants for both the substrates are greater than the respective dissociation constants. However, this procedure does not distinguish between the various possible sequential mechanisms, which include (Cleland, 1963) Ordered Bi Bi, Theorell-Chance, Random Bi Bi and Rapid-Equilibrium Random Bi Bi. Product-inhibition studies have now served to distinguish between these possibilities.

Inhibition by CoA was competitive with respect to acetyl-CoA and non-competitive with respect to choline; these results eliminate both the random mechanisms and suggest that an ordered addition of substrates is required. Inhibition by acetylcholine was competitive with respect to choline at saturating acetyl-CoA concentrations. In the presence of a saturating concentration of choline, acetylcholine produced no inhibition with respect to acetyl-CoA. These findings are consistent with the Theorell-Chance mechanism and inconsistent with the Ordered Bi Bi mechanism, which would require acetylcholine to produce non-competitive and uncompetitive inhibition respectively under the above conditions. Considerable problems were encountered in the experiments with acetylcholine owing to the small amount of choline that it contained; this difficulty was overcome by assaying the choline content and making the necessary allowance in the kinetic experiments. A further problem was that eserine, a competitive inhibitor of cholinesterase present in the enzyme preparation, was not totally effective in the presence of the high concentrations of acetylcholine used in the inhibition experiments, resulting in a considerable release of choline. This problem was overcome by pre-incubating the enzyme with 0.1 mM-di-isopropyl phosphorofluoridate, an irreversible inhibitor of the esterase.

The Theorell-Chance mechanism involves an ordered addition of substrates and the results described above are consistent with acetyl-CoA being the leading substrate. Enzyme binding studies with radioactive acetyl-CoA and choline are now being undertaken.

- Cleland, W. W. (1963). *Biochim. biophys. Acta*, **67**, 104.
McCaman, R. E. & Hunt, J. M. (1965). *J. Neurochem.* **12**, 253.
Morris, D. & Grewaal, D. S. (1969). *Biochem. J.* **114**, 85P.
Potter, L. T., Glover, V. A. S. & Saelens, J. K. (1968). *J. biol. Chem.* **243**, 3864.
Schrier, B. K. & Shuster, L. (1967). *J. Neurochem.* **14**, 977.
Schubert, J. (1966). *Biochim. biophys. Acta*, **122**, 470.

A Quantitative Investigation of the Phospholipid-Dependence of Rat Liver Microsomal Glucose 6-Phosphohydrolase

By JANE POULTER, B. R. CATER and T. HALLINAN.
(Department of Biochemistry, Royal Free Hospital School of Medicine, Hunter Street, London WC1N 1BP, U.K.)

Since the classical studies of Beaufay & de Duve (1954), much work has been done on the phospholipid dependence of glucose 6-phosphohydrolase (glucose 6-phosphatase). However, the specificity of this dependence is still unknown. We have initiated a quantitative investigation of glucose 6-phosphatase dependence on phosphatidylcholine and phosphatidylethanolamine.

Rat liver microsomal phospholipids were hydrolysed with phospholipase C (*Clostridium welchii*; Sigma) at 5 and 37°C. Glucose 6-phosphatase was assayed after hydrolysis had been stopped with EGTA,* which removes calcium essential for this phospholipase. Hydrolysis of phospholipids was measured quantitatively by using a sensitive radiochemical method involving direct counting of radioactive phospholipids in acid-washed microsomal precipitates (Hallinan, Nagley, Murty, Bennett & Grant, 1969). This simple radioassay was possible because conditions had been devised giving virtually specific labelling of phosphatidylcholine by radioactive choline *in vivo*, over 97% of the incorporated radioactivity entering phosphatidylcholine (Nagley & Hallinan, 1968). Similar specific labelling of phosphatidylethanolamine by radioactive ethanolamine has now been developed (B. R. Cater, unpublished work).

With EGTA present, phospholipase C caused no phospholipid hydrolysis and no inhibition of glucose 6-phosphatase. With calcium present, phospholipase C hydrolysed 90–100% of the microsomal phosphatidylcholine and 50–70% of the phosphatidylethanolamine at 37°C. Under these conditions glucose 6-phosphatase activity decreased approximately in parallel to phosphatidylcholine, approaching zero at 90–100% hydrolysis of the latter. However, at 5°C the kinetics were completely different. Though the same degree of phospholipid hydrolysis occurred, the enzyme activity no longer decreased in parallel with the phospholipid. Rather, enzyme activity decreased only 40–60% and then reached a plateau, following somewhat phosphatidylethanolamine, which reached a plateau at 50–70% hydrolysis.

Removal of exactly the same proportion of phospholipids has different effects at 5°C compared with 20 or 37°C, because when some phospholipid is hydrolysed glucose 6-phosphatase is

* Abbreviation: EGTA, ethanedioxybis(ethylamine)-tetra-acetic acid.