relatively small length of the PL1 molecule eliminate the possibility of cross-linkage of small free DNA zones in dNP by preparations PL1 and PL2. Thus 38% of the 'free' DNA zones are sufficiently long to accommodate either preparation PL1 or PL2. With preparation PL3, cross-linkage probably does occur and may account for its slightly higher titration value. The even higher values for preparation PL1 and for Toluidine Blue are presumably related to their smaller size.

Previous enzyme studies (R. F. Itzhaki, unpublished work) had shown that there are few, if any, extensive truly free DNA zones in dNP. This and the present result can be explained on the basis that dNP contains zones of phosphate groups unbound but hidden beneath protein. The distribution of basic residues in histones is very irregular (DeLange, Farnbrough, Smith & Bonner, 1969), and model-building studies (Richards & Pardon, 1970) with the F2a1 histone suggest that less than 60% of the DNA phosphate groups in dNP can be neutralized by the basic residues. The remaining phosphate groups would not be estimated as free by the enzyme method because of their associated protein, but would be free to bind dyes and polylysine. The relevance of the results to the role of histones in control of gene transcription will be discussed.

DeLange, R. J., Farnbrough, D. M., Smith, E. L. & Bonner, J. (1969). *J. biol. Chem.* **244**, 319.
Richards, B. M. & Pardon, J. F. (1970). *Expl Cell Res.* **62**, 184.

Investigations into the Nucleotide Sequence of Cysteine Transfer Ribonucleic Acid from Baker's Yeast

By N. J. HOLNESS and G. ATFIELD. (*Department of Experimental Biochemistry, The London Hospital Medical College, London E.1, U.K.*)

Cysteine tRNA has been isolated from baker's yeast RNA by a modification of the reversed-phase partition-chromatographic system described by Kelmers, Novelli & Stulberg (1965).

Cysteine tRNA represents about 1% of the RNA applied to the column and after chromatography is enriched at least 20-fold in its ability to accept [¹⁴C]cysteine in the standard assay system with a partially purified aminoacyl-tRNA synthetase prepared from rat liver. In our hands the maximum incorporation of [¹⁴C]cysteine corresponded to about 40% of that calculated, assuming mol.wt. 25000 for cysteine tRNA.

Our preparation does not accept significant quantities of serine, phenylalanine, tyrosine, iso-

leucine, proline and alanine. Further, there is no incorporation of radioactivity from a mixture of all ¹⁴C-labelled amino acids lacking cysteine. It is not further resolved by chromatography in other systems nor by countercurrent distribution for 500 transfers in a phosphate-formamide-propan-2-ol system (Apgar, Holley & Merrill, 1962). The elution patterns resulting from enzymic degradation described below are consistent with the view that this RNA preparation is substantially pure.

The nucleotide composition was established by DEAE-cellulose chromatography of an alkaline or ribonuclease T2 digest, with 0.07M-sodium acetate buffer, pH 4.10 or 4.15, as eluent. Oligonucleotides resulting from complete pancreatic and T1 ribonuclease hydrolyses have been isolated and partially characterized. Minor nucleotides identified as constituents of these oligonucleotides include ψ p, Tp, 5-Me-Cp, diMe-Gp and a nucleotide resembling isopentenyl-Ap.

Treatment with ribonuclease T1 at 2°C for 45 min affords two fragments that can be separated by DEAE-cellulose chromatography in 7M-urea at pH 7.9. With a higher enzyme:RNA ratio a more complex series of products is obtained. The sequences of these larger oligonucleotides are still to be determined.

Apgar, J., Holley, R. W. & Merrill, S. H. (1962). *J. biol. Chem.* **237**, 796.
Kelmers, A. D., Novelli, G. D. & Stulberg, M. P. (1965). *J. biol. Chem.* **240**, 3979.

Chemotaxonomy and Biosynthetic Relationships of Boletineae Pigments

By R. J. MEAD and W. SEGAL (introduced by G. M. GRAY). (*Department of Biochemistry, University of Western Australia, Nedlands, W.A., Australia*)

Morphological taxonomy of the Boletineae is attended with the disadvantage of subjective assessment of essentially qualitative features. The nature of the pigments and the presence or absence of a polyphenol oxidase in eight species of three Boletineae genera have therefore been examined as a possible aid in taxonomy.

T.l.c. of fungal extracts separates polyhydroxy-diphenylbenzoquinones and pulvinic acid derivatives, which have been identified by spectroscopic methods and colour reactions. Some *Boletus* species turn blue on bruising and this has been attributed to enzymic oxidation of variegatic and xerocommic acids, which are pulvinic acid derivatives (Steglich, Furtner & Prox. 1968; Beaumont, Edwards & Elsworth, 1968). In a study of eight Boletineae species four pulvinic acids have been isolated.