- Buffoni, F. (1966). *Pharmac. Rev.* **18**, 1163.
 Buffoni, F. & Blaschko, H. (1964). *Proc. R. Soc. B*, **161**, 153.
 Cleland, W. W. (1967). *A. Rev. Biochem.* **36**, 77.
 Estabrook, R. W. (1967). In *Methods in Enzymology*, vol. 10, p. 41. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Oi, S., Inamasu, M. & Yasunobu, K. T. (1970). *Biochemistry, Easton*, **9**, 3378.
 Reed, D. J. & Swindell, R. (1969). *Fedn Proc. Fedn Am. Socs exp. Biol.* **28**, 891.

Kinetic Studies of the N-Demethylation of Ethylmorphine by a Cytochrome P-450-Dependent Enzyme System in Human Liver Microsomes

By S. S. THORGEIRSSON and DONALD S. DAVIES.
 (Department of Clinical Pharmacology, Royal Postgraduate Medical School, London W.12, U.K.)

Oxidation is an important pathway for the inactivation of many lipid-soluble drugs and other foreign chemicals. Studies with liver microsomes from different species of laboratory animals have established a good inverse correlation between the duration of action of many lipid-soluble drugs *in vivo* and the activity of the drug-oxidizing enzymes of the different species (Quinn, Axelrod & Brodie, 1958). However, there have only been a few studies of human liver microsomal enzymes and in many cases the liver samples were obtained either at post mortem or were kept for periods of 1–3 h before use (Ackerman, 1970; Alvares *et al.* 1970). In view of the important role these enzymes may play in determining individual differences in response to various drugs in man (Davies, 1969) it was important to make kinetic studies of the oxidation of drugs with fresh samples of human liver.

Samples of liver were taken during abdominal surgery with the informed consent of the patient. Isolation of the microsomal fraction was commenced within 5–10 min of the samples being taken.

Concentrations of microsomal protein per g of liver ranged from 9.5 to 23 mg in the ten samples analysed. The concentration of cytochrome P-450 ranged from 0.67 to 1.02 nmol/mg of microsomal protein. These values are within the range of concentrations of cytochrome P-450 previously found in different species of laboratory animals (Davies, Gigon & Gillette, 1969). The maximal rates ($V_{\max}$) of N-demethylation of ethylmorphine showed a greater than fourfold range (2.0–8.8 nmol/min per mg), the highest value being similar to that found in the adult male rat. The apparent Michaelis constant (K_m) for the N-demethylation of ethylmorphine ranged from 0.8 to 4.0 mM, indicating

that the substrate has a lower affinity for human liver enzymes than for rat enzymes (K_m 0.32 mM). In addition the K_m values increased with $V_{\max}$, as would be the case in a diffusion-limited system (Davies, Gigon & Gillette, 1968).

The rate of reduction of cytochrome c varied from 176 to 282 nmol/min per mg, the equivalent value for the rat being 158 ± 24.3 . The rate of reduction of cytochrome P-450 ranged from 2.76 to 4.29 nmol/min per mg, which is less than in the rat. Reduction of cytochrome P-450, which is thought to be rate-limiting in the oxidation of drugs (Davies, 1969), correlated with $V_{\max}$ values and was increased after the addition of 2 mM-ethylmorphine. However, as has been previously reported (Gigon, Gram & Gillette, 1969), 0.5 mM-aniline decreased the initial rate of reduction of the cytochrome.

These observations demonstrate that there are individual differences in the rates of oxidation of ethylmorphine by cytochrome P-450-dependent enzymes in human liver microsomes. These differences are apparently not due to variations in the concentration of cytochrome P-450, but rather to differing rates of reduction of the cytochrome.

- Ackerman, E. (1970). *Biochem. Pharmac.* **19**, 327.
 Alvares, A. P., Schilling, G., Levin, W., Kuntzman, R., Brand, L. & Mark, L. C. (1970). *Clin. Pharmac. Ther.* **10**, 655.
 Davies, D. S. (1969). *Biochem. J.* **115**, 23–24P.
 Davies, D. S., Gigon, P. L. & Gillette, J. R. (1968). *Biochem. Pharmac.* **17**, 1865.
 Davies, D. S., Gigon, P. L. & Gillette, J. R. (1969). *Life Sci.* **8**, 85.
 Gigon, P. L., Gram, T. E. & Gillette, J. R. (1969). *Molec. Pharmac.* **5**, 109.
 Quinn, G., Axelrod, J. & Brodie, B. B. (1958). *Biochem. Pharmac.* **1**, 512.

The Electrophoretic Mobilities of Native and Chemically Modified Human Placental Alkaline Phosphatase Isoenzymes

By DILYS M. THOMAS and D. W. MOSS. (Department of Chemical Pathology, Royal Postgraduate Medical School, London W.12, U.K.)

At least 15 phenotypes of human placental alkaline phosphatase (EC 3.1.3.1) resulting from the action of nine structural genes have been recognized on the basis of the different electrophoretic mobilities of the enzymes at several pH values (Robson & Harris, 1967). However, the enzyme variants are closely similar in other respects (Thomas, 1970). To attempt to infer some of the structural differences that presumably underlie the differences in electrophoretic behaviour, we have compared the electrophoretic mobilities over a range of pH values of three common placental phosphatase

variants ('fast', 'slow' and 'intermediate') in their native forms and after acetylation, which increases the anodal migration of alkaline phosphatase without altering the characteristics of the residual enzyme activity (Moss, 1970).

All three forms are accelerated by acetylation: on starch-gel electrophoresis at pH 8.6 (tris-citrate-borate buffer) the slow isoenzyme is more accelerated than the fast variant, the modified forms having closely similar though not identical mobilities. The intermediate form behaves similarly to the fast isoenzyme at this pH value. Succinylation confirms the results of acetylation. The ratios of the mobilities of the acetylated and native forms are dependent on pH over the range 6.0–7.5 (phosphate buffers) and 8.0–10.4 (glycine buffers). The shape of these curves suggests the presence of modifiable charged groups with a pK_a in the region of pH 6.8, and that these groups make a greater contribution to the net charge of the intermediate isoenzyme in the pH range 6–8 than to that of the other two variants. Further groups with pK_a values about 9.0, which can be acetylated, are indicated, and the slow variant appears to possess a greater proportion of these groups than the fast and intermediate isoenzymes, accounting for its lower mobility and the more pronounced effect of acetylation.

Although these observations account for some of the electrophoretic differences between the three placental phosphatase isoenzymes, small residual differences in the mobilities of the acetylated forms suggest that further variations in structure also exist involving groups that are not acetylated. The groups in the protein or carbohydrate portions of the enzyme molecule that are modified by acetylation cannot be identified from these data alone, but extension of the method to include reagents of greater specificity may permit this, provided that such reagents do not also abolish enzymic activity.

Moss, D. W. (1970). *Enzymologia* (in the Press).

Robson, E. B. & Harris, H. (1967). *Ann. hum. Genet.* **30**, 219.

Thomas, D. M. (1970). Ph.D. Thesis: University of London.

Characterization of an Alkaline Phosphatase from Rat Heart

By EDITH MÜLLER, A. G. E. PEARSE and D. W. MOSS. (*Departments of Histochemistry and Chemical Pathology, Royal Postgraduate Medical School, London, W.12, U.K.*)

An alkaline phosphatase has been detected histochemically in sections of rat heart, activity being particularly marked in the tissues of the right atrium. The amount of this enzyme is increased after injections of catecholamines or isoprenaline

(Müller & Pearse, 1969). The enzyme activity has now been extracted from the atria of isoprenaline-treated and control male rats and certain of its properties have been investigated.

Homogenization of atria in sucrose solution or extraction of the tissue with butan-1-ol or autolysis give essentially equal recoveries of phosphatase activity. The activity in the left atrium of unstressed control animals averages about $0.003 \mu\text{mol}$ of *p*-nitrophenyl phosphate hydrolysed/min per mg of tissue by the method of Bowers & McComb (1966), compared with $0.007 \mu\text{mol}/\text{min}$ per mg in the right atrium. After subcutaneous injection of isoprenaline (40 mg/kg body wt.), right-atrial activity rises from three- to ten-fold, depending on the age of the animals, to values averaging $0.025 \mu\text{mol}/\text{min}$ per mg, but that of the left atrium reaches only $0.006 \mu\text{mol}/\text{min}$ per mg.

The atrial phosphatase has a pH optimum in the region of 9.7 at 5 mM substrate concentration and is activated by Mg^{2+} ions. It is not inhibited by L-phenylalanine (Fishman, Green & Inglis, 1962) and is thus of non-intestinal type by this criterion. Starch-gel electrophoresis shows only one major zone of alkaline phosphatase activity in each of the right and left atria. The zones in extracts of the two tissues have similar electrophoretic mobilities. The increase in activity after injection of isoprenaline does not appear to be accompanied by the appearance of additional enzyme zones.

Bowers, G. N. & McComb, R. B. (1966). *Clin. Chem.* **12**, 70.
Fishman, W. H., Green, S. & Inglis, N. R. (1962). *Biochim. biophys. Acta*, **62**, 363.

Müller, E. & Pearse, A. G. E. (1969). *Cardiovasc. Res.* **3**, 391.

Protease and Ribonuclease Activities in Bovine Pituitary Lobes

By J. C. PICKUP and D. B. HOPE. (*Department of Pharmacology, University of Oxford, Oxford OX1 3QT, U.K.*)

The acid and alkaline proteinase activity first described by Adams & Smith (1951) in bovine anterior and posterior pituitary lobes has been reinvestigated by an assay in which the trichloroacetic acid-soluble digestion products of denatured haemoglobin are estimated by measuring both the extinction at 280 nm and the colour produced by Folin-Ciocalteu reagent. The pH-dependence curve for fresh and acetone-dried posterior, and fresh anterior, bovine pituitary lobes confirms the presence of a cathepsin with a pH optimum at 3.8, but an optimum between pH 8.3 and 9.7 is given by u.v.-absorbing material that is Folin-negative and therefore not the result of proteolytic activity.

Controls showed that the u.v.-absorbing products