variants ('fast', 'slow' and 'intermediate') in their native forms and after acetylation, which increases the anodal migration of alkaline phosphatase without altering the characteristics of the residual enzyme activity (Moss, 1970).

All three forms are accelerated by acetylation: on starch-gel electrophoresis at pH 8.6 (tris-citrate-borate buffer) the slow isoenzyme is more accelerated than the fast variant, the modified forms having closely similar though not identical mobilities. The intermediate form behaves similarly to the fast isoenzyme at this pH value. Succinylation confirms the results of acetylation. The ratios of the mobilities of the acetylated and native forms are dependent on pH over the range 6.0–7.5 (phosphate buffers) and 8.0–10.4 (glycine buffers). The shape of these curves suggests the presence of modifiable charged groups with a pK_a in the region of pH 6.8, and that these groups make a greater contribution to the net charge of the intermediate isoenzyme in the pH range 6–8 than to that of the other two variants. Further groups with pK_a values about 9.0, which can be acetylated, are indicated, and the slow variant appears to possess a greater proportion of these groups than the fast and intermediate isoenzymes, accounting for its lower mobility and the more pronounced effect of acetylation.

Although these observations account for some of the electrophoretic differences between the three placental phosphatase isoenzymes, small residual differences in the mobilities of the acetylated forms suggest that further variations in structure also exist involving groups that are not acetylated. The groups in the protein or carbohydrate portions of the enzyme molecule that are modified by acetylation cannot be identified from these data alone, but extension of the method to include reagents of greater specificity may permit this, provided that such reagents do not also abolish enzymic activity.

Moss, D. W. (1970). *Enzymologia* (in the Press).

Robson, E. B. & Harris, H. (1967). *Ann. hum. Genet.* **30**, 219.

Thomas, D. M. (1970). Ph.D. Thesis: University of London.

Characterization of an Alkaline Phosphatase from Rat Heart

By EDITH MÜLLER, A. G. E. PEARSE and D. W. MOSS. (*Departments of Histochemistry and Chemical Pathology, Royal Postgraduate Medical School, London, W.12, U.K.*)

An alkaline phosphatase has been detected histochemically in sections of rat heart, activity being particularly marked in the tissues of the right atrium. The amount of this enzyme is increased after injections of catecholamines or isoprenaline

(Müller & Pearse, 1969). The enzyme activity has now been extracted from the atria of isoprenaline-treated and control male rats and certain of its properties have been investigated.

Homogenization of atria in sucrose solution or extraction of the tissue with butan-1-ol or autolysis give essentially equal recoveries of phosphatase activity. The activity in the left atrium of unstressed control animals averages about $0.003 \mu\text{mol}$ of *p*-nitrophenyl phosphate hydrolysed/min per mg of tissue by the method of Bowers & McComb (1966), compared with $0.007 \mu\text{mol}/\text{min}$ per mg in the right atrium. After subcutaneous injection of isoprenaline (40 mg/kg body wt.), right-atrial activity rises from three- to ten-fold, depending on the age of the animals, to values averaging $0.025 \mu\text{mol}/\text{min}$ per mg, but that of the left atrium reaches only $0.006 \mu\text{mol}/\text{min}$ per mg.

The atrial phosphatase has a pH optimum in the region of 9.7 at 5 mM substrate concentration and is activated by Mg^{2+} ions. It is not inhibited by L-phenylalanine (Fishman, Green & Inglis, 1962) and is thus of non-intestinal type by this criterion. Starch-gel electrophoresis shows only one major zone of alkaline phosphatase activity in each of the right and left atria. The zones in extracts of the two tissues have similar electrophoretic mobilities. The increase in activity after injection of isoprenaline does not appear to be accompanied by the appearance of additional enzyme zones.

Bowers, G. N. & McComb, R. B. (1966). *Clin. Chem.* **12**, 70.
Fishman, W. H., Green, S. & Inglis, N. R. (1962). *Biochim. biophys. Acta*, **62**, 363.

Müller, E. & Pearse, A. G. E. (1969). *Cardiovasc. Res.* **3**, 391.

Protease and Ribonuclease Activities in Bovine Pituitary Lobes

By J. C. PICKUP and D. B. HOPE. (*Department of Pharmacology, University of Oxford, Oxford OX1 3QT, U.K.*)

The acid and alkaline proteinase activity first described by Adams & Smith (1951) in bovine anterior and posterior pituitary lobes has been reinvestigated by an assay in which the trichloroacetic acid-soluble digestion products of denatured haemoglobin are estimated by measuring both the extinction at 280 nm and the colour produced by Folin-Ciocalteu reagent. The pH-dependence curve for fresh and acetone-dried posterior, and fresh anterior, bovine pituitary lobes confirms the presence of a cathepsin with a pH optimum at 3.8, but an optimum between pH 8.3 and 9.7 is given by u.v.-absorbing material that is Folin-negative and therefore not the result of proteolytic activity.

Controls showed that the u.v.-absorbing products