mevalonate (20 µg/ml) the concentrations of radioactive bactoprenol were again found to be in the same ratio: 6750 c.p.m./mg of protein in the mesosomes and 7700 c.p.m./mg of protein in the plasma membrane. This showed that bactoprenol is not initially synthesized in the mesosome membrane and then transferred to the plasma membrane. The same phenomenon is true for the fatty acids and phospholipids of the mesosome and plasma membranes of *Micrococcus lysodeikticus* and *Bacillus megaterium* (T. D. Thomas & D. J. Ellar, unpublished work; M. J. Daniels, unpublished work).

Barker, D. C. & Thorne, K. J. I. (1970). *J. Cell Sci.* **7**, 755.
Reaveley, D. A. & Rogers, H. J. (1969). *Biochem. J.* **113**, 67.

Thorne, K. J. I. & Kodicek, E. (1962). *Biochim. biophys. Acta*, **59**, 280.

Thorne, K. J. I. & Kodicek, E. (1966). *Biochem. J.* **99**, 123.

Control of Respiration in *Azotobacter vinelandii* Membranes

By C. W. JONES, S. K. ERICKSON and B. A. C. ACKRELL. (*Department of Biochemistry, School of Biological Sciences, University of Leicester, Leicester LE1 7RH, U.K.*)

In spite of many previous reports that bacterial respiratory membranes catalyse oxidative phosphorylation (see Gel'man, Lukyanova & Ostrovskii, 1967), only a very few of these membrane preparations (see John & Hamilton, 1970; Pandit-Hovenkamp, 1971; Eilermann, Pandit-Hovenkamp & Kolk, 1970; Jones & Ackrell, 1971) exhibit significant ADP + P_i-dependent respiratory control as defined by Chance & Williams (1956).

As part of a general study of the energy-conservation system of the nitrogen-fixing aerobe *Azotobacter vinelandii* (Ackrell & Jones, 1969, 1971a,b) we have investigated in detail various parameters that affect the ability of partly coupled (P/O, NADH ≤ 1.10) respiratory membranes from this organism to exhibit respiratory control. We have concluded that significant respiratory control is possible only if an energy-coupling site and the rate-limiting step in electron transfer from substrate to oxygen are coincident. If they are naturally coincident, or are made coincident experimentally (see Pandit-Hovenkamp, 1971; Jones & Ackrell, 1971), the respiratory control index is directly proportional to: (1) the efficiency of energy coupling at the controlling site (assayed as the P/2e ratio) and (2) the electron pressure at the controlling site (determined from the electron-transfer rates over different segments of the respiratory chain).

Each of these parameters was varied experimentally either *in vivo* (by using membranes prepared from cells harvested at different stages of the growth curve) or *in vitro* (by exposing the isolated respiratory membranes to a variety of electron donors and acceptors, or electron-transfer inhibitors and uncouplers). By means of such experiments respiratory control was readily demonstrated at phosphorylation sites I and II, and the respiratory control index could be reproducibly manipulated to yield values between 1.00 and 1.65.

These experiments will be described in detail and the conclusions will be discussed in terms of the special role of respiration in the aerobic growth of this organism on atmospheric nitrogen.

This work was supported by the Science Research Council (Grant no. B/SR/5063) and the University of Leicester.

Ackrell, B. A. C. & Jones, C. W. (1969). *Biochem. J.* **116**, 21P.

Ackrell, B. A. C. & Jones, C. W. (1971a). *Eur. J. Biochem.* (in the Press).

Ackrell, B. A. C. & Jones, C. W. (1971b). *Eur. J. Biochem.* (in the Press).

Chance, B. & Williams, G. R. (1956). *Adv. Enzymol.* **17**, 65.
Eilerman, L. J. M., Pandit-Hovenkamp, H. G. & Kolk, A. H. J. (1970). *Biochim. biophys. Acta*, **197**, 25.

Gel'man, N. S., Lukyanova, M. A. & Ostrovskii, D. N. (1967). *Respiration and Phosphorylation of Bacteria*, p. 171. New York: Plenum Press.

John, P. & Hamilton, W. A. (1970). *FEBS Lett.* **10**, 246.

Jones, C. W. & Ackrell, B. A. C. (1971). In *Energy Transduction in Respiration and Photosynthesis*. Bari: Adriatica Editrice (in the Press).

Pandit-Hovenkamp, H. G. (1971). In *Energy Transduction in Respiration and Photosynthesis*. Bari: Adriatica Editrice (in the Press).

Membrane-Bound Enzymes in Adrenal Medulla: An Adenosine Triphosphatase Characteristic of the Golgi Apparatus

By FRANÇOISE DUBOIS, I. BENEDECZKY and A. D. SMITH. (*Department of Pharmacology, University of Oxford, Oxford OX1 3QT, U.K.*)

The subcellular localization of some phosphohydrolases in the bovine adrenal medulla has been studied by centrifugation techniques and by histochemical analysis with the electron microscope.

The following enzymes were mainly recovered in the microsomal fraction of homogenates: Mg²⁺-stimulated adenosine monophosphatase, adenosine diphosphatase and uridine diphosphatase; Ca²⁺-stimulated inosine diphosphatase and adenosine triphosphatase. The presence of a Ca²⁺-stimulated adenosine triphosphatase in the microsomal fraction