R-factor had not been introduced, was incubated under the same conditions with the irradiated recipient. Since the donor strain is unable to utilize thymine and the chromosomal DNA of the recipient is unable to replicate, the incorporation excess is ascribed to the synthesis of the DNA of the R-factor after transfer to recipient.

Although incorporation of [^3H]thymine during incubation of the irradiated recipient strain with the donor strain lacking the R-factor was less than that observed during transfer of the R-factor, some incorporation did occur. In contrast, much less incorporation occurred when the two strains were incubated separately with [^3H]thymine. It was found that incubation of the irradiated recipient strain with [^3H]thymine gave rise to [^3H]dTMP in the medium. Moreover, this material, when added to a culture of the donor strain, was incorporated efficiently in trichloroacetic acid-insoluble material. It is believed that the conversion of thymine into dTMP followed by incorporation of this compound into the DNA of the donor bacterial accounts for the incorporation of radioactivity into DNA in these experiments in the absence of transfer of R-factor.

Financial support from the Medical Research Council, Science Research Council and Imperial Chemical Industries Ltd. (Pharmaceuticals Division) is gratefully acknowledged.

- Freifelder, D. & Freifelder, D. R. (1968). *J. molec. Biol.* **32**, 15.
Meynell, E. & Datta, N. (1966). *Genet. Res.* **7**, 134.
Silver, R. P. & Flakow, S. (1970). *J. Bact.* **104**, 331.

Genetic Expression of Sex-Factor after Transfer into Irradiated Cells of *Escherichia coli* *uvrA*⁻

By G. R. BARKER and T. M. TWOSE. (*Department of Biological Chemistry, University of Manchester, Manchester M13 9PL, U.K.*)

It has been demonstrated that, after transfer of the episome into cells of *Escherichia coli* previously killed by u.v. irradiation, synthesis of DNA of the F-merogenote (Freifelder & Freifelder, 1968) and of the R-factor Rldrd19 (Silver & Falkow, 1970; Barker, Hardman & Twose, 1971) takes place even though chromosomal DNA synthesis is suppressed in the recipient cells. Experiments have shown (N. Hardman & T. M. Twose, unpublished work) that cells subjected to doses of approx. 7000–8000 erg/mm² are non-viable and exhibit no measurable synthesis of protein or increase in cell mass. The present experiments are concerned with the ability of irradiated cells to support the normal functions of sex-factors.

The F'-lac episome was transferred to a suitable mutant of *E. coli*, which was then subjected to appropriate conditions for induction of the lac operon. It was found that induction of β -galactosidase activity was completely blocked by u.v. irradiation of the recipient with a dose of 7400 erg/mm² before introduction of the episome. At lower doses partial inhibition of the induction was observed. Analogous experiments to investigate the possible formation of enzymes following transfer of R-factor present special difficulties and have not yet been carried out.

The presence of sex-factor in a viable cell confers the ability to act as donor in conjugation with a second recipient. U.v. irradiation of recipient cells with a dose of 7400 erg/mm² before acceptance of either of the episomes Rldrd19 and F'-lac completely prevented the ability subsequently to act as donor in conjugation. With lower doses of irradiation partial inhibition was observed. It is concluded that the episomes do enter the u.v.-irradiated cells, since synthesis of DNA takes place after incubation of mixed cultures under conditions suitable for conjugation (Barker *et al.* 1971), but that expression of episomal genes probably does not take place.

Financial support from the Science Research Council and Imperial Chemical Industries Ltd. is gratefully acknowledged.

- Barker, G. R., Hardman, N. & Twose, T. M. (1971). *Biochem. J.* **122**, 48P.
Freifelder, D. & Freifelder, D. R. (1968). *J. molec. Biol.* **32**, 15.
Silver, R. P. & Falkow, S. (1970). *J. Bact.* **104**, 331.

Protein Content of Myelin

By W. L. G. GENT, N. A. GREGSON, CAROL A. LOVELIDGE and A. F. WINDER. (*Department of Biochemistry, Guy's Hospital Medical School, London S.E.1, U.K.*)

Determination of the protein content of membrane preparations is difficult because of the large amounts of lipids not readily dissociable from the proteins. Systematic study of the problem has been made by several methods to find a valid procedure for comparison of protein contents of central-nervous-system myelins from different animal species and of membranes from different tissues.

Removal of lipids by a variety of solvents from freeze-dried rat central-nervous-system myelin and gravimetric measurement of the protein residue yields results dependent on solvent (Winder & Gent, 1970a), but suggests that protein does not exceed 20% (w/w).

By taking the nitrogen content of myelin to be