PROCEEDINGS OF THE BIOCHEMICAL SOCIETY

The 513th Meeting of the Society was held at the Lord Rank Research Centre, High Wycombe, Buckinghamshire, on Thursday, 18 March 1971, when the following papers were presented.

COLLOQUIUM ON 'CEREAL AND VEGETABLE STARCHES IN USE: THEIR CHEMISTRY AND BIOCHEMISTRY'

Aspects of the Chemistry of Vegetable Starches

By C. T. GREENWOOD. (*Flour Milling and Baking Research Association, Chorleywood, Herts. WD3 5SH, U.K.*)

The physical and chemical characteristics of a starch depend uniquely on the botanical sources, and consequently a wide diversity of starches is found in vegetables. The fundamental properties of typical vegetable starches will be outlined, including the amylose/amylopectin ratio, the changes that occur during growth of the plant and the susceptibility of the granule to enzymic attack, and the ease with which dispersion of the resultant starch components will be discussed. The features of these typical vegetable starches, which differ from the cereal starches, will be outlined.

The Structures of Amylopectins and Glycogens

By W. J. WHELAN. (*Department of Biochemistry, University of Miami, Miami, Fla. 33152, U.S.A.*)

An enzyme (isoamylase) capable of the total debranching of amylopectin and glycogen has been discovered in a species of *Cytophaga* (Gunja-Smith, Marshall, Smith & Whelan, 1970). An examination of amylopectin and glycogen after sequential treatment with phosphorylase, β -amylase and isoamylase revealed facts at variance with the regularly debranched version of the structure proposed by Meyer & Bernfield (1940). A new arrangement of the unit chains of both macromolecules has therefore been proposed (Gunja-Smith, Marshall, Mercier, Smith & Whelan, 1970). The structure will be discussed in terms of its compatibility with the action patterns of the enzymes of metabolism.

Consideration will also be given to the roles of hydrolytic enzymes in starch and glycogen synthesis. Such enzymes may make important contributions to the polysaccharide structure when there is a lesion involving one of the normal enzymes of synthesis.

A fuller account of this work has been published by Whelan (1971).

Gunja-Smith, Z., Marshall, J. J., Mercier, C., Smith, E. E. & Whelan, W. J. (1970). *FEBS Lett.* **12**, 101.

Gunja-Smith, Z., Marshall, J. J., Smith, E. E. & Whelan, W. J. (1970). *FEBS Lett.* **12**, 96.

Meyer, K. H. & Bernfield, P. (1940). *Helv. chim. Acta*, **23**, 875.

Whelan, W. J. (1971). *Biochem. J.* **122**, 609.

Some Industrial Aspects of the Enzymic Degradation of Starch

By D. J. MANNERS. (*Department of Brewing and Biological Sciences, Heriot-Watt University, Edinburgh EH1 1HX, U.K.*)

Although the use of amylolytic enzymes in the food and fermentation industries is steadily increasing (see, e.g., Saletan, 1968), the present paper is confined to a discussion of only two developments. These are the production of glucose and glucose syrups from starch, and the replacement of malt by barley and enzyme preparations in the brewing process.

Starch may be quantitatively converted into D-glucose by amyloglucosidase (glucoamylase) preparations (for a review see Fleming, 1968). The substrate is attacked from the non-reducing end and the stepwise hydrolysis of successive glucosidic linkages gives glucose as the primary product. Amyloglucosidases are produced by many fungi, e.g. *Aspergillus* sp., *Rhizopus* sp., *Endomyces* sp., and several examples have been selected for large-scale production, particularly by the Japanese fermentation industry.

Amyloglucosidases have only limited action on starch granules, and a preliminary treatment of these is necessary. In the 'acid-enzyme' conversion process the starch is treated initially with hot mineral acid under pressure, to effect gelatinization and solubilization, and to bring about some degradation of polysaccharide material. This 'thinning' action is then followed by enzymic saccharification of the neutralized starch hydrolysate. More recently, preliminary treatment of the starch with a relatively thermostable bacterial α -amylase has been used in an 'enzyme-enzyme' conversion (Denault &