hydroxylases, nitro reductase, glucuronyltransferase and cytochrome *P*-450 (Parke & Rahman, 1970). Induction of these enzymes by pretreatment of rats with other anutrients, such as phenobarbitone, 3-methylcholanthrene or β -ionone (Parke & Rahman, 1969), gives rise also to an increase of microsomal cytochrome *b*₅. Induction by pretreatment with safrole and isosafrole (125mg/day per kg) gives rise to an increase of cytochrome *b*₅ and also to the appearance of a new redox difference absorption spectrum, which progressively increases over 3 days of continued treatment with these anutrients, and may represent the biosynthesis of a new hepatic microsomal haemoprotein. Whereas the redox difference absorption spectrum of cytochrome *b*₅ (NADH) has a maximum at 425nm and that of cytochrome *P*-450 (NADPH) occurs at 445nm, the absorption maximum for the new haemoprotein occurs at 455nm and is manifest after reduction with NADH or NADPH. This new absorption peak is not therefore attributable to the increase of cytochrome *P*-450.

The new redox difference absorption spectrum does not appear when the safrole pretreatment of rats is accompanied by simultaneous administration of thioacetamide to inhibit nuclear-directed protein synthesis (Adams & Busch, 1963), or of 3-amino-1,2,4-triazole to inhibit the synthesis of haem (Baron & Tephly, 1969). Solubilization of the hepatic microsomal pellet from safrole-treated rats, by treatment with 0.5% deoxycholate, results in the disappearance of both cytochrome *b*₅ (425nm) and the new haemoprotein (455nm) from the 100000g deposit. The cytochrome *b*₅ absorption maximum (425nm) reappears quantitatively in the 100000g supernatant fraction, whereas the absorption maximum at 455nm, although present, is much diminished. Addition of safrole or isosafrole to the 100000g microsomal pellet obtained from liver homogenates from normal rats does not result in the appearance of the absorption maximum of this new haemoprotein, nor to any new absorption maximum other than that due to cytochrome *b*₅. From these observations it is considered that the new redox difference absorption spectrum may be attributed to the formation of a new microsomal haemoprotein, and is not merely an adduct of cytochrome *b*₅ with safrole, isosafrole or any of their metabolites.

Adams, H. R. & Busch, H. (1963). *Cancer Res.* **23**, 576.
 Baron, J. & Tephly, T. R. (1969). *Molec. Pharmac.* **5**, 10.
 Parke, D. V. & Rahman, H. (1969). *Biochem. J.* **113**, 12P.
 Parke, D. V. & Rahman, H. (1970). *Biochem. J.* **119**, 53P.

The Deacetylation of *N*-Arylacetamides *in vitro*

By J. W. GORROD. [Department of Pharmacy, Chelsea College of Science and Technology (University of London), London S.W.3., U.K.]

The administration of certain aromatic amines or their acetylated derivatives to experimental animals leads to the development of cancer (Clayson, 1962). However, as dogs fed on diets containing aromatic amines also develop tumours, although being unable to acetylate the toxicant (Poirier, Miller & Miller, 1963), it is possible that deacetylation may be a prerequisite for the initiation of the carcinogenic process resulting from exposure to *N*-arylacetamides.

Further, whereas 2-naphthylamine and 4-aminobiphenyl are carcinogenic, 1-naphthylamine and 2-aminobiphenyl are inactive (Weisburger & Weisburger, 1966). The carcinogenic compounds show considerable variation in their ability to produce tumours in different species. As the metabolism of an acetamido compound may differ from that of the parent amine (Booth & Boyland, 1964) the deacetylase activity of various species was examined *in vitro*.

Deacetylase activity was measured by determining the formation of aromatic amines from the corresponding acetamido compound by a g.l.c. technique (Gorrod, 1968). A deacetylase activity active towards 2- and 4-acetamidobiphenyl, 1- and 2-acetamidonaphthalene and 2-acetamidofluorene was found localized in the hepatic microsomal fraction of various species examined; optimum activity occurred between pH 8 and 9. The deacetylase activity was strongly inhibited by sodium fluoride (1mM) but was not inhibited by SKF 525A (2-diethylaminoethyl-2,2-diphenylvalerate hydrochloride) (2mM).

Hydrolysis of acetamido compounds was observed with hepatic microsomal preparations obtained from rat, cat, dog, mouse, hamster, cavy and human; 1-acetamidonaphthalene and 2-acetamidobiphenyl were the most rapidly hydrolysed substrates, whereas 2-acetamidofluorene was the poorest. The carcinogenic substrates were generally hydrolysed to their greatest extent in the species in which they were able to produce tumours, although only rarely did the rate of hydrolysis exceed that of the non-carcinogenic counterpart. The ease of hydrolysis of 2-acetamidobiphenyl is in accord with the failure of Gorrod & Carey (1970) to detect this compound as a metabolite of 2-aminobiphenyl.

N-(1-Naphthylthio)acetamide was completely resistant to this deacetylase activity in either rat or cavy preparations, which were easily able to hydrolyse the oxygen analogue.

The results reported do not support the idea of a direct relationship between deacetylation and

carcinogenic activity. It may be that oxidation of the nitrogen enhances the ease of hydrolysis, as suggested by Irving (1966), thereby allowing release of the *N*-hydroxylated carcinogen. Acetylation of an aromatic amine usually directs ring hydroxylation away from the *ortho*-position to give rise to inactive metabolites. It is therefore surprising to find such a high rate of hydrolysis associated with the isomers having the lower toxicity.

This work was carried out while the author was a Research Fellow of The Royal Commission for the Exhibition of 1851.

- Booth, J. & Boyland, E. (1964). *Biochem. J.* **91**, 362.
 Clayson, D. B. (1962). *Chemical Carcinogenesis*. London: J. and A. Churchill Ltd.
 Gorrod, J. W. (1968). *Tetrahedron Lett.* no. 59, p. 6155.
 Gorrod, J. W. & Carey, M. J. (1970). *Biochem. J.* **119**, 52P.
 Irving, C. C. (1966). *Cancer Res.* **26**, 1390.
 Poirier, L. A., Miller, J. A. & Miller, E. C. (1963). *Cancer Res.* **23**, 790.
 Weisburger, J. H. & Weisburger, E. K. (1966). *Chem. Engng News*, **43**, 124.

A New Form of *N*-Acetyl- β -glucosaminidase Present in Pregnancy Serum

By J. L. STIRLING. (*Department of Biochemistry, Queen Elizabeth College, Campden Hill, London W.8, U.K.*)

N-Acetyl- β -glucosaminidase activity in human serum has been shown to increase during pregnancy (Walker, Woollen & Pugh, 1960). By histochemical investigation they showed that there were high *N*-acetyl- β -glucosaminidase activities in the decidua cells and the chorion, and they suggested that these sites might be the origin of the elevated serum activity.

Two main forms of *N*-acetyl- β -hexosaminidase (A and B) are known to exist in human tissues and these can be differentiated by starch-gel electrophoresis (Robinson & Stirling, 1968; Okada & O'Brien, 1969). Serum shows only *N*-acetyl- β -glucosaminidase A when examined in this way, but when sera collected at various stages of pregnancy (1 month to full term) are examined two electrophoretically distinct forms can be seen. The elevation of total activity is due to an increased amount of form A and to the appearance of a new species of enzyme designated *N*-acetyl- β -glucosaminidase P. This pregnancy form of the enzyme has a slow anodic mobility in electrophoresis at pH 7.0 and can be distinguished from *N*-acetyl- β -glucosaminidase A, which migrates rapidly towards the anode. It can also be distinguished from form B, found in tissues, which moves towards the cathode.

N-Acetyl- β -glucosaminidases A and P can be

separated by DEAE-cellulose chromatography at pH 7.0 in 10mM-sodium phosphate buffer by application of a sodium chloride gradient, peak P appearing at a salt concentration of 0.12M and peak A at 0.19M. There was no separation of A and P activities on gel filtration through Sephadex G-200.

The thermal stability of enzyme P is intermediate between that of form A, which is rapidly denatured at pH 5.0 and 50°C, and that of form B, which is almost completely stable under these conditions. K_m values for *N*-acetyl- β -glucosaminidases A and P were 0.65 and 0.55mM respectively, but both had optimum pH 4.5 in 0.1M-sodium citrate buffer. *N*-Acetyl- β -glucosaminidases A and P could not be distinguished by inhibition due to acetate, *N*-acetyl-galactosamine or *p*-chloromercuribenzoate, and both were unaffected by dithiothreitol and *N*-ethylmaleimide.

When extracts of placenta and foetal membranes were investigated by starch-gel electrophoresis and DEAE-cellulose chromatography both tissues had A and B activities but only a trace of P activity, which was possibly contributed by the maternal blood in the organ. All the *N*-acetyl- β -glucosaminidase in cord serum at birth is of the A type.

This work was carried out at Makerere University, Uganda, and supported by a grant from the Publications and Research Grants Committee.

- Okada, S. & O'Brien, J. S. (1969). *Science, N.Y.*, **165**, 698.
 Robinson, D. & Stirling, J. L. (1968). *Biochem. J.* **107**, 321.
 Walker, P. G., Woollen, M. E. & Pugh, D. (1960). *J. clin. Path.* **13**, 353.

Activation of Coagulation Factor X by Insoluble Trypsin

By R. M. HOWELL and R. J. DUFE. (*Department of Biochemistry, Queen Elizabeth College, Campden Hill, London W.8, U.K.*)

Factor X occupies an important position in the final stage of coagulation leading to the activation of prothrombin. The parent zymogen (Factor X) is converted into the active form (Factor Xa) by agents that precede it in the clotting sequence, and additionally by trypsin (Hanahan & Papahadjopoulos, 1965) and a component of Russell's viper venom (Williams & Esnouf, 1962).

The present work is a continuation of an investigation on clot production *in vivo* in which non-proteolytic activation of Factor X was examined with various surface-active agents (Howell & Scott, 1967). For comparison a reliable protease activator is required, but prolonged contact with soluble trypsin leads to rapid loss of clotting activity because proteolysis cannot be terminated. Russell's viper venom is unsuitable as an activator for systems