gut bacteria responsible for the conversion of cyclamate into cyclohexylamine in rat faeces are different from those responsible for the conversion in human or rabbit faeces. It appears that the organisms responsible are clostridia in rat faeces, enterococci in human faeces and enterobacteria in rabbit faeces. Clostridia are uncommon in rabbit faeces (Smith, 1965).

All the colonies of enterococci from the faeces of the human convertor metabolized cyclamate to cyclohexylamine. Relevant to this are the observations that the numbers of enterococci in faeces from Japanese subjects are considerably higher (10^8 /g) than in faeces from British subjects (10^5 /g) (Hill *et al.* 1971) and that all Japanese subjects examined excreted cyclohexylamine after oral doses of cyclamate (Kojima & Ichibagase, 1966).

This work was supported in part by a grant from the International Sugar Research Foundation.

- Drasar, B. S. (1967). *J. Path. Bact.* **94**, 417.
 Golberg, L., Parekh, C., Patti, A. & Soike, K. (1969). *Toxic. appl. Pharmac.* **14**, 654.
 Hill, M. J., Crowther, J. S., Drasar, B. S., Hawksworth, G., Aries, V. & Williams, R. E. O. (1971). *Lancet*, i, 95.
 Kojima, S. & Ichibagase, H. (1966). *Chem. pharm. Bull., Tokyo*, **14**, 971.
 Renwick, A. G. & Williams, R. T. (1969). *Biochem. J.* **114**, 78 p.
 Renwick, A. G. & Williams, R. T. (1970). *Proc. 1st int. Sugar Res. Conf., Brussels*, p. 3. Bethesda, Md.: International Sugar Research Foundation Inc.
 Schaedler, R. W., Dubos, R. & Costello, R. (1965). *J. exp. Med.* **122**, 59.
 Smith, H. W. (1965). *J. Path. Bact.* **89**, 95.

The Metabolism of Methamphetamine in the Rat and Guinea Pig

By J. CALDWELL, L. G. DRING and R. T. WILLIAMS.
 (Department of Biochemistry, St Mary's Hospital Medical School, London W.2, U.K.)

Methamphetamine (2-methylamino-1-phenylpropane) is a dangerous drug from the point of view of drug-dependence, but little is known about its metabolism. It is partly excreted as such and as amphetamine in man (Cartoni & di Stefano, 1963; Beckett & Rowland, 1965), and largely demethylated to amphetamine in the dog (Axelrod, 1954).

[^{14}C]Methamphetamine [(±)-2-methylamino-1-phenyl[^{14}C]propane] was synthesized and its fate in rats and guinea pigs examined (dose 45 mg/kg). A marked species difference was found. In the rat the ^{14}C after oral administration was excreted mainly in the urine (about 80% in 3 days) with little (3%) in the faeces, but in the guinea pig about

70% appeared in the urine and 18% in the faeces after intraperitoneal injection. The main reaction in the rat was aromatic hydroxylation, the metabolites being 4-hydroxy-*N*-methylamphetamine (31%), 4-hydroxynorephedrine (16%) and 4-hydroxyamphetamine (6%), which were excreted as glucuronic acid conjugates. Unchanged methamphetamine (11% of the dose) and amphetamine (3%) were also found in the urine. In the guinea pig no aromatic hydroxylation occurred, for the main metabolite was benzoic acid (32%; mainly conjugated) followed in order by norephedrine (20%), amphetamine (13%) and methamphetamine (4%). These metabolites were found in the urine and faeces except benzoic acid and its conjugates, which were found only in the urine.

The main metabolic reaction of methamphetamine in the rat was thus aromatic hydroxylation and in the guinea pig *N*-demethylation, but both species were able to hydroxylate the α -carbon atom of the side chain since appreciable amounts of norephedrine (2-amino-1-phenylpropan-1-ol) were excreted by the guinea pig and of 4-hydroxynorephedrine by the rat. It has been suggested (Fischer, Horst & Kopin, 1965; Thoenen, Hürliman, Gey & Haefely, 1966) that norephedrine and 4-hydroxynorephedrine may act as false transmitters at nerve endings and may be involved in the undesirable effects of the chronic intake of the amphetamines. An earlier study on amphetamine (Dring, Smith & Williams, 1970) showed that little if any 4-hydroxynorephedrine was formed from amphetamine in the rat and norephedrine was not detected as a metabolite of amphetamine in the guinea pig. Kosman & Unna (1968) have reported that the continuous administration of methamphetamine to the rat and guinea pig causes depression, but with amphetamine given similarly the animals become excited. These latter observations may be correlated with the significant formation of norephedrine or its hydroxy derivative from methamphetamine but not from amphetamine.

- Axelrod, J. (1954). *J. Pharmac. exp. Ther.* **110**, 315.
 Beckett, A. H. & Rowland, M. R. (1965). *J. Pharm. Pharmac.* **17**, 1095.
 Cartoni, G. P. & di Stefano, S. (1963). *G. Biochim.* **8**, 298.
 Dring, L. G., Smith, R. L. & Williams, R. T. (1970). *Biochem. J.* **116**, 425.
 Fischer, J. E., Horst, W. D. & Kopin, I. J. (1965). *Br. J. Pharmac.* **24**, 477.
 Kosman, M. E. & Unna, K. R. (1968). *Clin. pharmac. Ther.* **9**, 240.
 Thoenen, H., Hürliman, A., Gey, K. F. & Haefely, W. (1966). *Life Sci.* **5**, 1715.