- Beary, M. E. (1967). *Biochem. J.* **103**, 54 P.
Conney, A. H. (1967). *Pharmac. Rev.* **19**, 317.
Dorfman, R. I. & Ungar, F. (1965). *Metabolism of Steroid Hormones*. New York: Academic Press Inc.
Sasame, H. A., Castro, J. A. & Gillette, J. R. (1968). *Biochem. Pharmac.* **17**, 1759.

An Intact Muscle Preparation Suitable for Studies of the Metabolic Effects of Electrical Stimulation

By E. B. FERN, V. M. PAIN and K. L. MANCHESTER.*
(Department of Biochemistry, University College London, Gower Street, London WC1E 6BT, U.K.)

Studies of the metabolic effects of stimulation on mammalian muscle have usually been carried out on whole hind limbs, either stimulated *in situ* or in perfused preparations. Changes in concentrations of metabolites in the plasma or perfusate from such systems can only indicate the response of a group of muscles, so that information on individual muscles is limited to that obtained by excision and analysis after stimulation. We have designed a system whereby an isolated muscle preparation may be electrically stimulated while allowing the incubation mixture to be readily sampled.

The system is basically that described by Pain & Manchester (1970) with modifications, and was developed for studies of the metabolism of the extensor digitorum longus muscle from young female rats. Each muscle is attached to an electrode and incubated at 37°C in 2 ml of Krebs-Ringer bicarbonate buffer continuously gassed with O₂ + CO₂ (95:5). The other electrode is immersed in the medium. Constant tension is applied to the muscle during the passage of electric stimuli, the voltage, duration and frequency of which may be varied to suit a particular type of muscle. Additions to the medium or withdrawal of samples can easily be carried out throughout the incubation period.

When extensor digitorum longus muscles of 30 mg or less were incubated in this system for 60 min, tissue weights were unaltered, and the concentration of K⁺ in the intracellular fluid was maintained at 88% (resting) and 76% (stimulated) of that in zero-time controls. With larger muscles we observed increases in tissue weight and extracellular fluid and decreases in K⁺ and ATP concentrations, these changes being particularly pronounced in stimulated muscles. Probably the O₂ supply becomes critical in muscles of this size, and it is likely that this would also be the case for

muscles with a higher proportion of aerobic red fibres.

E. B. F. is grateful to CIBA Laboratories, Horsham, Sussex, U.K., for financial support.

Pain, V. M. & Manchester, K. L. (1970). *Biochem. J.* **118**, 209.

The Physiological Role of Taurine

By L. K. KACZMAREK and A. N. DAVISON. (*Medical Research Council Membrane Biology Research Group, Department of Biochemistry, Charing Cross Hospital Medical School, London WC2N 4HH, U.K.*)

In a previous communication (Kaczmarek, Agrawal & Davison, 1970) we showed that taurine is present in nerve-ending particles isolated from rat brain, together with its synthesizing enzyme, cysteinesulphinate decarboxylase. In both distribution and properties this enzyme closely resembles glutamate decarboxylase, which is responsible for the synthesis of γ -aminobutyrate, a compound believed to act as a neuroinhibitor (Davison, 1956; Agrawal, Davison & Kaczmarek, 1971). We have now extended the work to a study of the uptake of taurine by cortical tissue, as well as its release into extracellular space on stimulation. Rat brain-cortex slices were first incubated for 30 min in the presence of radioactive taurine. The slices were transferred to cells that were perfused with oxygenated taurine-free Krebs-Ringer medium. After an initial washout of taurine from the extracellular space a slow loss of radioactivity from the tissue was observed. On the application of rectangular electrical pulses to the tissue, the efflux of radioactivity was considerably enhanced, a result that has been found for γ -aminobutyrate by Srinivasan, Neal & Mitchell (1969). If such a process is of physiological significance and taurine is released from nerve endings to exert an inhibitory influence on the postsynaptic membrane, it should be accompanied by a mechanism for the termination of its action. This may be by re-uptake into the tissue, as has been proposed for the cessation of γ -aminobutyrate action (Iversen & Neal, 1968). We have found that slices of rat brain cortex will efficiently take up radioactive taurine at low concentrations (0.5 μ M) from Krebs phosphate medium. This work adds support to the suggestion from iontophoretic evidence (Curtis & Watkins, 1965) that taurine may be considered as a possible neurotransmitter, but much further work will be necessary to verify this hypothesis.

Thanks are due to Dr J. F. Mitchell and Mr B. Rayner of the Department of Pharmacology, University of Cambridge for their invaluable help in the release experi-

* Present address: Department of Biochemistry, University of the West Indies, Mona, Kingston 7, Jamaica.