- Owen, W. J., Rogers, L. J. & Hayes, J. D. (1970). *Biochem. J.* **121**, 6P.
 Yamashita, T. & Butler, W. (1968). *Pl. Physiol., Lancaster*, **43**, 1978.
 Yamashita, T. & Butler, W. (1969). *Pl. Physiol., Lancaster*, **44**, 435.

Partial Purification from Rat Liver of an Enzyme Catalysing the Sulphurylation of Dehydroepiandrosterone

By R. RYAN and J. CARROLL. (*Department of Biochemistry, Trinity College, Dublin 2, Irish Republic*)

The sulphotransferases of mammalian tissues catalyse the transfer of the sulphuryl group of adenosine 3'-phosphate 5'-sulphatophosphate to a variety of acceptors, which include phenol, dehydroepiandrosterone, oestrone, L-tyrosine methyl ester and serotonin. None of these sulphotransferases has been obtained in a pure state, but Banerjee & Roy (1966) described the partial separation from guinea-pig liver of a number of sulphotransferases. They achieved a 20- and 70-fold increase in the specific sulphurylating activities towards dehydroepiandrosterone and oestrone respectively, but both of these fractions were active towards phenol. Mattock & Jones (1970) purified from female rat liver an enzyme that catalysed the sulphurylation of L-tyrosine methyl ester, but this purified preparation also catalysed the synthesis of *p*-nitrophenyl sulphate. McEvoy & Carroll (1970) described the isolation from male rat liver of an enzyme that catalysed the sulphurylation of phenols but had no activity towards steroids, L-tyrosine methyl ester or serotonin. Adams & Poulos (1967) isolated from bovine adrenals an oestrogen sulphotransferase that lacked activity towards 3 β -hydroxy steroids and *p*-nitrophenol. The present communication describes the partial purification from rat liver of a steroid sulphotransferase with high activity towards dehydroepiandrosterone and weak or no activity towards other acceptors.

Livers from female Wistar rats were homogenized in 0.15M-potassium chloride with a Potter-Elvehjem homogenizer, and centrifuged at 100 000g for 1 h. The resulting supernatant was made 1 mM with respect to 2-mercaptoethanol and diluted to a protein concentration of 10 mg/ml. The sulphotransferase was precipitated by the addition of cold ethanol to a concentration of 25% (v/v). The precipitated protein was dissolved in 5 mM-tris-HCl buffer, pH 8.0, and stirred at 2°C for 1 h. After centrifugation at 25 000g for 15 min, the supernatant was collected and adjusted to 55% saturation with respect to ammonium sulphate. The precipitated protein was dissolved in 5 mM-tris-HCl buffer,

pH 8.0, and dialysed against large volumes of the same buffer. During the dialysis inactive protein was precipitated and was discarded. Further purification was achieved by chromatography on columns of DEAE-cellulose and phosphocellulose P-11, and gel filtration on Sephadex G-200. The specific activity towards dehydroepiandrosterone of this partially purified sulphotransferase was 60-fold higher than that of the crude supernatant, and the enzyme had negligible activity towards *p*-nitrophenol, L-tyrosine methyl ester and serotonin.

Adams, J. B. & Poulos, A. (1967). *Biochim. biophys. Acta*, **146**, 493.

Banerjee, R. K. & Roy, A. B. (1966). *Molec. Pharmac.* **2**, 56.

McEvoy, F. A. & Carroll, J. (1970). *Biochem. J.* **119**, 26P.

Mattock, P. & Jones, J. G. (1970). *Biochem. J.* **116**, 797.

A Heparan Sulphate Sulphotransferase of Ox Lung

By T. FOLEY and J. R. BAKER. (*Department of Biochemistry, Trinity College, Dublin 2, Irish Republic*)

Suzuki & Strominger (1960) first showed that a preparation from hen oviduct could catalyse the sulphation of heparan sulphate by adenosine 3'-phosphate 5'-sulphatophosphate. Suzuki, Trenn & Strominger (1961) reported the partial purification and stability of the enzyme. Using similar methods, we have extended their studies. In seeking a more convenient source of the enzyme, a study was made of its distribution in rat tissues. Marked activity was detected in lung and brain homogenates only. For purposes of purification, ox lung was the sole source of the enzyme.

Experiments were designed to determine whether the enzyme was responsible for synthesizing *N*-sulphate or *O*-sulphate groups in heparan sulphate. [Both types of sulphate ester are present in this glycosaminoglycan (Brown, 1957).] Chemically de-*N*-sulphated heparan sulphate is a better substrate for the enzyme than is heparan sulphate, suggesting that enzymic *N*-sulphation is more likely. [³⁵S]Sulphate was enzymically transferred from adenosine 3'-phosphate 5' [³⁵S]-sulphatophosphate to de-*N*-sulphated heparan sulphate. The product was purified, and unsubstituted amino groups were acetylated, before incubation with a crude 'heparinase' from *Flavobacterium heparinum*. A paper-chromatographic separation of the resulting incubation mixture in an isobutyric acid-ammonia solvent system (Suzuki & Strominger, 1960) revealed only two radioactive areas. The major ³⁵S-labelled component had the same *R_f*