are being currently tested with regard to binding to DNA, effects on transcription by the rat liver RNA polymerases and binding of [^3H]cortisol.

Independent of the existence of secondary control cycles, we consider that the initial reaction to the hormonal induction, which triggers the whole series of consequent events, is the stereospecific binding of the hormone to a non-histone regulatory protein on the DNA in a way analogous to the binding of the bacterial inducers to their specific repressors. Because of the complexity of the genome of higher organisms, the development of systems *in vitro* testing this hypothesis is still a long way off. However, the current rate of progress in the molecular biology of eukaryotes lends hope to such prospects.

This work was supported by the Deutsche Forschungsgemeinschaft.

Beato, M., Homoki, J., Lukacs, I. & Sekeris, C. E. (1968). *Hoppe-Seyler's Z. physiol. Chem.* **340**, 1099.

- Beato, M., Schmid, W., Braendle, W. & Sekeris, C. E. (1971). In *Advances in the Biosciences*, vol. 7. Ed. by Raspé, J. New York: Pergamon Press Inc. (in the Press).
- Beato, M., Seifart, K. H. & Sekeris, C. E. (1970). *Archs Biochem. Biophys.* **138**, 272.
- Dahmus, M. E. & Bonner, J. (1965). *Proc. natn. Acad. Sci. U.S.A.* **54**, 1370.
- Karlson, P. & Sekeris, C. E. (1966). *Acta endocr., Copnh.*, **53**, 505.
- Paul, J. & Gilmour, R. S. (1968). *J. molec. Biol.* **34**, 305.
- Roeder, R. G. & Rutter, W. J. (1969). *Nature, Lond.*, **224**, 234.
- Seifart, K. H. & Sekeris, C. E. (1969). *Eur. J. Biochem.* **7**, 408.
- Sekeris, C. E. (1969). In *Progress in Endocrinology*, p. 7. Ed. by Gual, C. Amsterdam: Excerpta Medica Foundation.
- Sekeris, C. E., Homoki, J., Beato, M., Gallwitz, D., Seifart, K. H. & Lukacs, I. (1968). In *Advances in the Biosciences*, vol. 2, p. 222. Ed. by Raspé, G. New York: Pergamon Press Inc.

COMMUNICATIONS

Transfer Ribonucleic Acid Methylases in Normal Tissues and in Tumours

By V. M. CRADDOCK. (*Medical Research Council Laboratories Toxicology Unit, Woodmansterne Road, Carshalton, Surrey, U.K.*)

Much suggestive evidence has accumulated to support the idea, first put forward by Srinivasan & Borek (1963), that tRNA methylases may be involved in carcinogenesis (Craddock, 1970). Much of this evidence was based on work with tumours caused by viruses, and enzyme changes could have been related to viral induction rather than to malignancy. The present experiments show that a change in tRNA methylases occurs in tumours caused by chemicals, and that certain tRNA methylases are affected more than others.

The enzymes were studied in the high-speed supernatant fraction of homogenates by incubation with *S*-adenosyl[$\text{Me-}^{14}\text{C}$]methionine and a limiting amount of *Escherichia coli* tRNA. Enzyme preparations from normal tissues methylate tRNA to a low limited extent and then stop. According to current concepts, methylation of certain sites in the tRNA molecule takes place, the sites being determined by the specificity of the enzymes, and the 'saturation curve' reaches a plateau where an increase in the amount of enzyme present does not cause further methylation to take place. However, there are other possible explanations of this effect. Preparations from tumours of liver or kidney that

had been produced by treatment of animals with nitrosamines methylate *E. coli* tRNA more extensively than do preparations from normal tissues. When tRNA that had been incubated with methylase preparations and *S*-adenosyl[$\text{Me-}^{14}\text{C}$]methionine was analysed after hydrolysis by column chromatography, the major methylated bases formed were 5/3-methylcytosine, di- N^2 -methylguanine, N^2 -methylguanine and 1-methyladenine. There was a definite difference in the relative amounts of the different methylated bases produced depending on whether the methylase preparation was from tumour or from normal tissue. This difference was found to occur also between methylases prepared from a liver tumour compared with the preparation from 'normal' liver tissue adjacent to the tumour.

Liver taken from animals 24h after partial hepatectomy gave a 'saturation level' of methylation similar to that given by normal liver, and the relative amounts of the different methylated bases formed were similar to those given by preparations from normal liver. The differences between tumours and normal liver may therefore be related not simply to growth but to altered growth.

The author thanks Mr N. Howe and Mr J. Fletcher for skilled technical assistance.

- Craddock, V. M. (1970). *Nature, Lond.*, **228**, 1264.
- Srinivasan, P. R. & Borek, E. (1963). *Proc. natn. Acad. Sci. U.S.A.* **49**, 529.