Analysis of the trichloroacetic acid-soluble fraction from [$U-^{14}C$]glutamate showed that radioactivity had been incorporated into glutathione, ophthalmic acid, proline and pyrrolidone-5-carboxylic acid. Aspartate was not labelled. Most of the radioactivity in the trichloroacetic acid-soluble extract of the incubation with [$U-^{14}C$]aspartate was found in glutamate and glutathione. Alanine was also labelled. Most of the radioactivity from [$U-^{14}C$]alanine was recovered as unchanged alanine, but significant amounts of labelled lactate, glutamate, glutathione and aspartate were present.

These studies indicate that the role of the citric acid cycle in lens metabolism is greater than has been believed hitherto.

- Kuck, J. F. R. (1970). In *Biochemistry of the Eye*, p. 267. Ed. by Graymore, C. N. New York and London: Academic Press.
- Trayhurn, P. & van Heyningen, R. (1971). *Exptl Eye Res.* **12**, (in the Press)
- van Heyningen, R. (1965). *Biochem. J.* **96**, 419.
- van Heyningen, R. (1969). In *The Eye*, 2nd ed., vol. 1, p. 393. Ed. by Davson, H. New York and London: Academic Press.
- Waley, S. G. (1964). *Biochem. J.* **91**, 576.

Aromatization of Androst-4-ene-3,17-dione by Human Intestinal Bacteria

By P. GODDARD and M. J. HILL. (*Department of Bacteriology, St Mary's Hospital Medical School, London W.2, U.K.*)

As part of a wider study of cancer of the colon (Hill *et al.* 1971) we are investigating the nuclear dehydrogenation of steroids by intestinal bacteria. We report here the formation from androst-4-ene-3,17-dione of a novel steroid, tentatively identified as 17-methoxyoestratrien-1,3,5(10)-3-ol.

A culture of a strain of *Clostridium paraputrificum* (CC493) grown anaerobically at 37°C for 72h in Todd-Hewitt broth (Oxoid) containing 0.25mm-substrate was adjusted to pH 8 and shaken aerobically at 37°C for 96h after the addition of a further 0.25mm-substrate. The steroids were extracted with chloroform and t.l.c. analysis revealed the formation in moderate yield of a product that reacted with Folin reagent and with ferric chloride-ferricyanide reagent (both of which indicate the presence of a phenolic hydroxyl group). The product did not correspond to oestrone or oestradiol and behaved as a less-polar steroid.

The u.v.-absorption spectrum (maximum at 282nm, ϵ 2300) and the fluorescence spectrum (excitation maximum at 283nm with a corresponding emission maximum at 330nm) both indicated the presence of an aromatic ring A and a phenolic hydroxyl group, presumably still at C-3. The absence of a second absorption maximum at

about 300nm indicated that the product was not a seco-steroid produced by ring B fission.

Mass spectroscopy showed that the molecular weight of the product was 286, corresponding to methyloestradiol, the result of a migration of the 10-methyl group followed by aromatization of ring A. Such methyl migrations occur chemically to give 1-methyloestradiol or 1,17-dihydroxy-4-methyloestra-1,3,5(10)-triene (Dreiding, Plummer & Tomasewski, 1953). Both of these were synthesized and shown to differ from the bacterial product in t.l.c. behaviour and i.r. spectra. The trifluoroacetyl derivative was prepared and shown by mass spectroscopy to be the trifluoroacetate of oestradiol (molecular weight 464), indicating that the methyl group was lost during formation of the derivative. This suggested that the methyl group was not nuclear but at the 17-oxygen position.

Oestradiol was methylated with dimethyl sulphate, as was the bacterial product. Both yielded derivatives that were identical on t.l.c., and presumably corresponded to 3,17-dimethyloestradiol.

This result suggests that the bacterial product was the result of a migration of the methyl group from C-10 to the 17-oxygen function, 17-methoxyoestra-1,3,5(10)-trien-3-ol. Both androst-4-ene-3,17-dione and 17-hydroxyandrost-4-en-3-one were substrates for this novel migration. The reaction was carried out by a range of opalescent-negative clostridia, but no strains of the other genera tested (e.g. *Escherichia coli*, *Bacteroides* spp., *Bifidobacterium* spp.).

Financial support from the Cancer Research Campaign is gratefully acknowledged.

- Dreiding, A., Plummer, W. & Tomasewski, A. (1953). *J. Am. chem. Soc.* **75**, 3159.
- Hill, M. J., Crowther, J. S., Drasar, B. S., Hawksworth, G., Aries, V. & Williams, R. E. O. (1971). *Lancet*, *i*, 95.

Spectral Changes of a Cationic Dye after Interaction with Endotoxins

By J. JANDA. (*Department of Biochemistry, Imperial College of Science and Technology, London S.W.7, U.K.*)

The use of a carbocyanine dye for detection of undegraded endotoxins as defined by Nowotny (1969) was examined. Spectral shifts resulting from interactions of various compounds with the dye were observed by using a modified procedure of a method estimating polysaccharides (Edstrom, 1969).

New absorption maxima were found in the range 430–485nm. They occurred with certain endotoxin preparations and long-chain aliphatic sulphates. Under various conditions the shifts could serve as