Activity and Regulation of an Acidic Amino Acid Permease of *Aspergillus nidulans*

By J. H. ROBINSON, C. ANTHONY and W. T. DRABBLE. (*Department of Physiology and Biochemistry, University of Southampton, Southampton SO9 5NH, U.K.*)

An acidic amino acid permease is formed during germination of *Aspergillus nidulans* conidia. This permease is inhibited by sodium azide (95% at 10 mM), is inactive at 2°C and exhibits both pH and temperature optima. Uptake of glutamate occurs when the intracellular glutamate concentration is 100 times the extracellular concentration. Basic and neutral amino acids do not competitively inhibit the permease. L-Aspartate and L-cysteic acid are competitive inhibitors of glutamate uptake, and conversely glutamate is a competitive inhibitor of aspartate uptake. The K_m for glutamate at 30°C is 180 μ M; the K_i for glutamate inhibition of aspartate uptake is 140 μ M. There is similar agreement between the values for aspartate (K_m , 100 μ M; K_i for inhibition of glutamate uptake, 140 μ M). The K_m for glutamate is the same both before and some hours after germ-tube emergence. Mycelium germinated on glutamate as sole nitrogen source has the same K_m for glutamate as mycelium germinated on ammonia.

Aspergillus conidia take up glutamate at very low rates. Permease activity increases 100-fold during germination when one of the following compounds is the sole nitrogen source: L-glutamate, L-aspartate, L-cysteic acid, glycine, L-lysine. The increase is 50-fold with nitrate as sole nitrogen source, but less than eightfold with ammonia, methylamine, urea or asparagine. The increase in permease activity resulting from germination in a glutamate medium is prevented by the simultaneous presence of ammonia. This effect of ammonia is less marked in a mutant that is resistant to methylamine; some of the enzymes of this mutant are resistant to repression by ammonia (Arst & Cove, 1969). When conidia germinated with ammonia as nitrogen source are transferred to a glutamate medium the permease activity increases tenfold within 4 h. This change in activity is prevented by ammonia (25 mM) or cycloheximide. These results suggest that formation of the permease is regulated by a mechanism involving repression by ammonia, or by a product of ammonia metabolism.

Further, there is some evidence that ammonia, or a metabolite of ammonia, inhibits the permease. Preincubation for 30 min with ammonia (25 mM) leads to about 40% inhibition of the rate of glutamate uptake, but no inhibition occurs when ammonia is added 1 min before glutamate (0.1 mM).

We thank Rank Hovis McDougall (Research) Ltd. for financial assistance.

Arst, H. N. & Cove, D. J. (1969). *J. Bact.* **98**, 1284.

A Cytochrome-Deficient Mutant of *Pseudomonas* AM1

By C. ANTHONY and P. DUNSTAN. (*Department of Physiology and Biochemistry, University of Southampton, Southampton SO9 5NH, U.K.*)

Two mutants of *Pseudomonas* AM1 have been isolated that have lost the ability to oxidize methanol and ethanol. Except for these alcohols, mutant PCT-29 grows on the same substrates as wild-type *Pseudomonas* AM1 and is able to oxidize formaldehyde, formate, methylamine, succinate and malate. Mutant PCT-76 is superficially similar, except that methylamine does not support growth and whole cells do not oxidize this substrate. Mutant PCT-76 may be a double mutant; its properties with respect to methylamine are therefore not further discussed in this communication.

Ultrasonic extracts of mutant PCT-29 do not have the alcohol dehydrogenase described by Anthony & Zatman (1965), showing that this enzyme is essential for growth of *Pseudomonas* AM1 both on methanol and on ethanol.

Ultrasonic extracts of mutant PCT-76, however, have wild-type activities of the alcohol dehydrogenase. The failure of this mutant to oxidize alcohols can be explained by the absence of cytochrome *c*, which is the only soluble cytochrome found in wild-type *Pseudomonas* AM1 (Anthony, 1970). Ultrasonic extracts of mutant PCT-76 contain no soluble cytochromes and, further, no cytochrome *c* can be observed in whole cells of this mutant.

Membrane fragments from mutant PCT-76 contain all the cytochromes and all the carotenoid pigment of the mutant, and appear to be identical with membrane fragments from the wild-type organisms. Results obtained with pigment mutants of *Pseudomonas* AM1 indicate that the carotenoid pigments are not involved in electron transport in this organism. The *a*-type cytochrome, which is found together with *b*-type cytochrome in membrane fragments from mutant PCT-76 and wild-type organisms, is probably the cyanide-sensitive component involved in the oxidation of all substrates by *Pseudomonas* AM1.

The results presented in this communication demonstrate an unexpected complexity in the electron-transport system in *Pseudomonas* AM1. The cytochrome *c* of this organism can interact with the systems for electron transport from all substrates that can be oxidized (Anthony, 1970).