Nucleotide Sequence Analysis with Polynucleotide Kinase and Nucleotide 'Mapping' Methods

5'-TERMINAL SEQUENCES OF DEOXYRIBONUCLEIC ACID FROM BACTERIOPHAGES λ AND 424

By KENNETH MURRAY

Department of Molecular Biology, University of Edinburgh, Edinburgh EH9 3JR, U.K.

(Received 31 July 1972)

The polynucleotide kinase reaction was used in analyses of complex mixtures of oligodeoxynucleotides which were fractionated by various two-dimensional nucleotide 'mapping' procedures. Parallel ionophoretic analyses on DEAE-cellulose paper, pH2, and AE-cellulose paper, pH3.5, of venom phosphodiesterase partial digests of 5'-terminally labelled oligonucleotides enabled the sequence of the nucleotides to be deduced uniquely. A 'diagonal ionophoresis' method has been used with mixtures of nucleotides. Application of these methods to 5'-terminally labelled DNA from bacteriophage λ gave the terminal sequences pA-G-G-T-C-G and pG-G-G-C-G. Identical 5'-terminal sequences were found with DNA from bacteriophage 424.

Polynucleotide kinase, which transfers the γ -phosphorus of ATP to the 5'-hydroxyl group of nucleotides, provides an invaluable approach to the sequence analysis of nucleic acids and oligonucleotides (Novogrodsky & Hurwitz, 1965, 1966; Richardson, 1965). In addition to its use for labelling DNA molecules or large fragments of them at their 5'-termini (Richardson, 1966), this reaction enables work to be done on an ultramicro scale with digests of unlabelled nucleic acids. The polynucleotide kinase reaction has been used to identify the 5'-terminal nucleotides of bacteriophage T7 DNA (Richardson, 1966; Richardson & Weiss, 1967) and of DNA from bacteriophage λ and related bacteriophages, which are the nucleotides at the cohesive ends of these molecules (Wu & Kaiser, 1967). When used in conjunction with the high-resolution nucleotide 'mapping' methods that have been developed for radioactive nucleotides (Sanger *et al.*, 1965; Murray, 1970), the reaction facilitates analysis of complex mixtures of nucleotides such as those resulting from endonuclease digests of nucleic acids (Murray, 1968; Murray & Wu, 1968). Although only the 5'-terminus of a nucleotide becomes labelled in the reaction, sequence analysis is still possible by measurement of differences in ionophoretic mobility on ion-exchange papers (Sanger *et al.*, 1965) between the components of a partial exonuclease (venom phosphodiesterase) digest of the 5'-labelled nucleotide or from analyses of the components of an alternate exonuclease (spleen phosphodiesterase) partial digest which have been labelled after the partial digestion. The polynucleotide kinase reaction has been applied previously to endonuclease digests of DNA to provide a source

of labelled nucleotides for the preparation of α - ^{32}P -labelled deoxynucleoside 5'-triphosphates of very high specific radioactivity (Murray & Wu, 1968). Szekely & Sanger (1969) and Southern (1970) have also used the reaction to analyse pyrimidine nucleotides derived from DNA by the Burton degradation (Burton & Petersen, 1960).

The present paper describes applications of the polynucleotide kinase reaction to the analysis of endonuclease (pancreatic deoxyribonuclease) digests of DNA and the use of these digests in direct determination of 5'-terminal nucleotide sequences of DNA from bacteriophages λ and 424.

Materials and Methods

Abbreviation and notation of nucleotides

Since only deoxynucleotides are discussed in this paper, the usual prefix 'd' has been omitted throughout. All the nucleotides referred to in the text, Plates, Figures and Tables have a 5'-phosphate, unless stated otherwise, which has also been omitted throughout. Note that the numbers of nucleotides in a given Plate or Figure and Table do not necessarily correspond with those in another Plate or Figure and Table, except for Plates 4 and 5 where they do correspond. Similarly, a given nucleotide may carry a different number in the different Plates or Figures and their corresponding Tables.

Materials

Cellulose acetate (Oxoid) strips for electrophoresis were obtained from Oxo Ltd., London E.C.4, U.K.

Ion-exchange papers used were Whatman Chromedia AE-81 and DE-81. The dyes used for electrophoretic markers (Xylene Cyanol FF, Orange G, and acid Fuchsin) were obtained from G. T. Gurr Ltd., London S.W.6, U.K. CsCl used in the preparation of bacteriophage was the AnalaR grade from British Drug Houses Ltd., Poole, Dorset, U.K. Streptomycin sulphate was generously given by Glaxo Laboratories Ltd., Ulverston, Lancs., U.K. DNA from bacteriophages λ and 424 was prepared by the procedure of Kaiser & Hogness (1960). Phenol (AnalaR grade) was distilled under N_2 and collected under air-free water; it was usually used within 24 h and was equilibrated with 0.5 M-Tris-HCl buffer (pH 7.5 or 8.0) immediately before use. Bacteriophage T7 DNA was kindly provided by Dr. R. S. Hayward. Chicken erythrocyte DNA was prepared from saline-washed deoxyribonucleoprotein (Murray *et al.*, 1968) by titration to pH 0.8 at -1°C (Murray, 1969). Calf thymus DNA and salmon sperm DNA were purchased from Sigma (London) Chemical Co. Ltd., Kingston-upon-Thames, Surrey, U.K.

Bacteriophage λ . Several methods were used for the preparation of this bacteriophage. In the earlier stages of the work, the heat-inducible lysogen *Escherichia coli* C600 (λ c1857) was used both in liquid culture (usually in batches of 6×500 ml in 1-litre Roux bottles) and on large plates ($24\text{ cm} \times 24\text{ cm}$) of agar (10 g of Difco tryptone, 5 g of NaCl, 10 g of Difco agar per litre adjusted to pH 7.5). In either case, bacteriophage were usually recovered from the lysates (which were titred on *E. coli* W1485 and normally contained about 2×10^{10} – 6×10^{10} plaque-forming units/ml) by centrifugation at 21 000 rev./min in an MSE 10×100 ml rotor in an SS50 or SS65 centrifuge for 2½ h. The supernatant normally contained less than 1% of the bacteriophage and was discarded. The pellets were resuspended by gentle shaking overnight at 4°C in about 1/25 the original lysate volume of λ -dil medium (Kaiser & Hogness, 1960). Bacteriophage suspensions were clarified by centrifugation at 10 000 rev./min in a MSE 8×50 ml rotor in an HS 18 centrifuge for 10 min and incubated for 1–3 h at 25°C with pancreatic ribonuclease (10 $\mu\text{g}/\text{ml}$) and pancreatic deoxyribonuclease (10 $\mu\text{g}/\text{ml}$). The bacteriophage were then sedimented, resuspended, and the suspensions clarified as before. CsCl was added to 41.5% (w/w) and the bacteriophage were recovered by isopycnic centrifugation (Kaiser & Hogness, 1960). The bacteriophage band was removed with a bent, sterile Pasteur pipette, diluted with 41.5% (w/w) CsCl in λ -dil medium and again centrifuged as above. The bacteriophage band was then collected via a needle through the bottom of the tube and bacteriophage preparations were stored as suspensions in CsCl solution at 4°C until required.

Preparation of DNA. The bacteriophage were

diluted to a titre of about 5×10^{12} with water and dialysed against 1 mM-EDTA buffered at pH 8.0 with 0.01 M-Tris-HCl buffer. Greatly improved yields of the bacteriophage were obtained by the use of a heat-inducible lysogen of the lysis-defective strain λ S7; two such strains of *E. coli* were used: 803(λ S7c1857) and W1485(λ S7c1857). Cultures in L broth or M9 medium (Adams, 1959) supplemented with: casein hydrolysate (Oxoid), 1.5%; glucose, 0.4%; NaCl, 0.05% and MgSO_4 , 0.02%, were grown at 32°C to a cell density of about $5 \times 10^8/\text{ml}$, induced at 45°C for 10–15 min, and growth continued at 37°C for 2–3 h. The pH was monitored and maintained above 7 by addition of small quantities of 1 M-Tris-HCl buffer, pH 8. Cells were harvested by centrifugation (5 min at 5000 rev./min, MSE HS 18 centrifuge), resuspended in a convenient volume of λ -dil medium, cooled to 0°C and lysed by addition of a few drops of chloroform. After about 1 h, during which the suspension was shaken intermittently, the lysate was clarified by centrifugation (30 min, 10 000 rev./min, MSE HS 18 centrifuge, 8×50 ml or 6×250 ml rotor) and titred on *E. coli* strain Y mel (an *suIII*⁺ strain). Bacteriophage were purified from the lysate as described above for preparations from the normal lysogen. This procedure was particularly useful for the preparations of radioactively labelled bacteriophage. In such preparations, the radioactive isotope was added after induction, and for preparations of ^{32}P -labelled bacteriophage a low-phosphate medium was used.

Bacteriophage 424. A spontaneous clear-plaque mutant was picked from a lysate made by heat induction of the lysogen *E. coli* C600 (424cI^+) and a liquid lysate was prepared from it. A 2.5 litre culture of *E. coli* strain C600 at a cell density of $5 \times 10^8/\text{ml}$ was infected with this lysate at a multiplicity of 1 and growth continued until the E_{650} reached a minimum. The culture was then lysed with chloroform and clarified by centrifugation at 10 000 rev./min (MSE HS 18 centrifuge, 6×250 ml rotor). Bacteriophage were recovered from the lysate and purified as described for bacteriophage λ .

Enzymes. The following were purchased from Worthington Biochemical Corp., Freehold, New Jersey, U.S.A.: pancreatic deoxyribonuclease, snake venom phosphodiesterase, spleen phosphodiesterase, and bacterial alkaline phosphatase. Bacterial alkaline phosphatase was also obtained from Whatman Biochemicals Ltd., Maidstone, Kent, U.K. Preparations of bacterial alkaline phosphatase were diluted to a concentration of 1 mg/ml in 0.1 M- NH_4HCO_3 and heated at 100°C for 5 min before use (sometimes it was necessary to clarify the solution after heating by centrifugation). The preparations were examined for endonuclease contamination by incubation with radioactively labelled bacteriophage λ DNA which was then analysed by sedimentation through a sucrose

gradient as described for polynucleotide kinase: the preparations obtained from Whatman Biochemicals Ltd. were essentially free from deoxyribonucleases. Glyceraldehyde phosphate dehydrogenase (rabbit muscle) and 3-phosphoglycerate kinase (yeast), were purchased from Sigma (London) Chemical Co. Ltd. Semen phosphomonoesterase was prepared by the method of Wittenberg & Kornberg (1953). Endonuclease I from *E. coli* prepared as described by Lehman *et al.* (1962) was generously provided by Dr. P. R. Brown.

Polynucleotide kinase. This was prepared from *E. coli* strain B infected at a cell density of 2×10^9 /ml with bacteriophage T4 at a multiplicity of 4. Cells were chilled for 20 min after infection and harvested at once. Frozen cells were ground with twice their weight of alumina, mixed with 0.05 M-Tris-HCl buffer, pH 7.4 (0.001 M in 2-mercaptoethanol) (7.5 ml/g of cells), and centrifuged (MSE HS18 centrifuge) at 10000g for 30 min at about 2°C. The supernatant was centrifuged (MSE SS50 centrifuge) at 70000g for 2 h at about 2°C and the residue was discarded. The enzyme was prepared from the supernatant solution by the procedure of Richardson (1965), except that the DEAE-cellulose chromatography step was repeated (as described by Wu & Kaiser, 1967, except that eluate fractions were not collected into ATP solution) to ensure removal of nucleotides. To minimize contamination with phosphatase and endonuclease activities, only the most-active fractions of the eluate from DEAE-cellulose were retained for chromatography on phosphocellulose. To test for the presence of endonuclease activity, samples of the phosphocellulose eluate fractions (10 μ l) were incubated (37°C, 30 min) with radioactively labelled DNA from bacteriophage λ and sedimented through 6–20% (w/v) sucrose gradients (MSE SuperSpeed centrifuge, 3 $\times$ 3 ml SW rotor, 50000 rev./min at 20°C for 125 min). Some phosphocellulose eluate fractions of polynucleotide kinase have been stored at 0°C for over a year without appreciable loss of activity. In later preparations, conversion of a dinucleoside monophosphate into its dinucleotide was used as a test for enzyme activity. This was particularly useful when many samples (e.g. fractions from column chromatography) were to be analysed, for about 20 samples could be examined simultaneously, easily and economically, and with relatively little expenditure of time. For example, 1 μ l of the sample was incubated in a capillary tube (37°C, 30 min) with 10 μ l of solution containing Tris-HCl buffer, pH 7.4 (500 nmol), MgCl₂ (50 nmol), 2-mercaptoethanol (125 nmol), C-U (10 nmol), and [γ -³²P]-ATP (0.025 nmol; about 10^5 c.p.m. of ³²P) and the reaction mixture was then applied to AE-cellulose paper for ionophoresis at pH 3.5. The mobility of pC-U with respect to the blue marker (Xylene Cyanol FF) is about 0.9 whereas ATP hardly moves

from the origin even on prolonged ionophoresis under these conditions. The paper was then scanned with a hand monitor (Mini Instruments Ltd., London W.C.2, U.K.) and where desirable a more precise observation was made by radioautography or counting with a strip scanner (Packard radiochromatogram scanner). Parallel experiments with a range of C-U concentrations were used to estimate enzyme concentration.

[γ -³²P]ATP. This was prepared by the method of Glynn & Chappell (1964), adapted as follows for the small-scale production of a product of very high specific radioactivity. A solution of [³²P]P_i (20 mCi) was evaporated to dryness in its vial in a vacuum desiccator over NaOH pellets and conc. H₂SO₄. Evaporation was repeated after addition of water (1 ml). The following reaction mixture was then transferred to the vial: 10 μ mol of Tris buffer, pH 8.5; 1 μ mol of MgCl₂; 2 μ mol of ATP; 0.4 μ mol of 3-phosphoglycerate; 2 μ mol of 2-mercaptoethanol; 2 μ l of phosphoglycerate kinase; 5 μ l of 3-phosphoglyceraldehyde dehydrogenase; total volume 0.2–0.3 ml. (In some cases, larger quantities of the enzyme were used.) The mixture was maintained at room temperature (20°C) for about 30 min and then applied in a 15 cm streak to a sheet of Whatman 3 MM paper for ionophoresis at pH 3.5 or pH 4.4 in 1 M-pyridine-acetate buffer. The ATP band was detected by radioautography (about 1 s exposure) and eluted with water (usually in a total volume of about 2 ml). The solutions were stored frozen. In calculating the specific radioactivity of the product it was assumed that the quantity of ATP recovered was equal to that put into the reaction mixture as ATP: this was checked by counting successive elute fractions from the paper. All operations were carried out behind screens of 12.7 mm ($\frac{1}{2}$ in) Perspex (Lucite) and papers carrying radioactive material were handled sandwiched between Perspex sheets. These procedures were convenient when particularly high specific radioactivities of ATP in essentially salt-free solutions were required. Yields of ATP varied between 10 and 80%, the reason for which has not been established. Control experiments with small quantities of radioactive phosphate were usually very efficient (at least 75% of the phosphate being transferred to ATP), but the same mixture with the total batch of [³²P]P_i frequently gave a much lower yield. Control experiments and samples from preparative reactions were analysed by descending chromatography (about 2–3 h) on DEAE-cellulose paper in 0.3 M-ammonium formate (A. A. Travers, unpublished work) with a reference sample of ATP that could be detected under u.v. light. Variations in time and temperature of the reaction were relatively unimportant as shown in Table 1. It was important to use a freshly prepared reaction mixture, for the yields of ATP from reactions at 0°C (30 min) with reaction mixtures (containing

enzymes, but no radioactive phosphate) that had been previously kept at 0°C for 1, 2½ and 12 h were, respectively, 70, 71 and 38 %. [γ - ^{32}P]ATP of very high specific radioactivity is now produced on a monthly schedule, and at a reasonable price, by The Radiochemical Centre, Amersham, Bucks., U.K. These preparations have been used in recent experiments.

Methods

Dephosphorylation of pancreatic deoxyribonuclease digests of DNA. Solutions of DNA (0.005–0.1 ml; E_{259} 0.5–2.0) were digested with pancreatic deoxyribonuclease, usually at a concentration of 0.1 mg/ml in 0.01 M-sodium acetate buffer, pH 5.0, at 37°C for at least 1 h and often overnight. The solutions were then incubated with bacterial alkaline phosphatase (0.1 vol. of a 0.5 mg/ml solution in 0.1 M- NH_4HCO_3) at 37°C for 2 h. In some experiments treatment with deoxyribonuclease and phosphatase was effected simultaneously, usually for 3 h at 37°C. Some digests were dephosphorylated with semen phosphomonoesterase; the digest was incubated with 0.1–0.05 vol. of the enzyme preparation in 0.1 M-sodium acetate buffer, pH 6.0 (Wittenberg & Kornberg, 1953). After the digestion, bacterial alkaline phosphatase was inactivated by treatment with EDTA (0.02 M) at pH 2 or less (10 min, 20°C; Dahlberg, 1968) and the solution was neutralized after the addition of Tris-HCl buffer (0.1 M), pH 7.5. Semen phosphomonoesterase was inactivated by heating at 100°C for 2 min. Reactions were usually done on a micro scale (about 20 μl of DNA solution) to keep volumes low for the polynucleotide kinase reaction and its subsequent analysis; this is particularly desirable if the reaction products are to be analysed with a cellulose acetate/DEAE-cellulose paper system rather than with ion-exchange papers alone.

Phosphorylation with polynucleotide kinase. Reaction conditions were essentially as described by Richardson (1965). For the phosphorylation of reaction products in which phosphatase had been inactivated by treatment with EDTA at low pH it was necessary, obviously, to ensure that an excess of Mg^{2+} was added. Small-scale reactions (about 35 μl after digestion and dephosphorylation of 20 μl of DNA solution) were incubated at 37°C with 20 μl of a solution containing 2 μmol of Tris-HCl buffer, pH 7.5, 0.2–1.0 μmol of MgCl_2 , 0.01 μmol of ATP (γ - ^{32}P -labelled), 0.5 μmol of 2-mercaptoethanol, and 5 μl of polynucleotide kinase (phosphocellulose fraction, usually >2000 units/ml). Similar proportions were used with larger reaction volumes. Reaction mixtures were usually incubated for about 1 h at 37°C and sometimes overnight (at either 37°C or room temperature). Ionophoresis on AE-cellulose paper at pH 3.5 was used to check the reaction before analysis of the reaction products by a two-dimensional

ionophoretic method. Where necessary, reaction mixtures were evaporated to dryness in a vacuum desiccator and dissolved in a very small volume of buffer (about 5 μl) for ionophoresis.

Phosphorylation of DNA. Carefully purified DNA from bacteriophage λ was heated (60°C, 10 min, then cooled in ice) to break Hershey circles and concatamers, treated with phosphatase, extracted three times with freshly distilled phenol, and phosphorylated with polynucleotide kinase as described by Wu & Kaiser (1967). In some instances the kinase reaction was stopped by heating (100°C, 2 min; the solution usually became somewhat turbid) before proceeding to further operations. Labelled DNA from these reactions was recovered by acid precipitation in 3% (w/v) HClO_4 (15 min, 0°C) after addition of carrier DNA. The precipitate was dissolved in 0.1 M-NaOH (0.1 ml) and again precipitated with 3% HClO_4 and the precipitate was washed with 80% (v/v) aq. ethanol, before being dissolved (overnight at room temperature) in 0.1 M- NH_4HCO_3 , evaporated to dryness in a vacuum desiccator and then digested with pancreatic deoxyribonuclease as described above, or with *E. coli* endonuclease I (Lehman *et al.*, 1962). Gel filtration on Sephadex G-100 was also used for recovery of the terminally labelled DNA and is preferable to acid

Table 1. *Effect of time and temperature on the conversion of radioactive phosphate into radioactive ATP*

A reaction mixture consisting of 1 M-Tris-HCl buffer, pH 8.5 (10 μl), 0.1 M- MgCl_2 (10 μl), 0.1 M-2-mercaptoethanol (18 μl), 0.1 M-ATP (20 μl), 0.075 M-sodium 3-phosphoglycerate (5 μl), glyceraldehyde 3-phosphate dehydrogenase (5 μl), 3-phosphoglycerate kinase (2 μl) and water (50 μl) was mixed with $^{32}\text{P}[\text{P}_i]$ (10 μCi) that had been evaporated to dryness in a glass tube. The mixture was divided into three equal parts which were incubated at 0°, 25° and 37°C. Samples (about 10 μl) were withdrawn from each tube after 5, 15 and 60 min and applied to DEAE-cellulose paper for descending chromatography in 0.3 M-ammonium formate. A reference sample of ATP was included on the paper. Radioactive compounds were detected on the dried paper by radioautography and the appropriate zones of the paper were cut out and their radioactivity determined by counting in a liquid-scintillation spectrometer.

Time of reaction Temperature (°C)	% radioactivity as ATP		
	5 min	15 min	60 min
0	84	89	90
25	84	85	81
37	80	82	75

precipitation, but the method eventually adopted as a routine was sedimentation through sucrose gradients [4.2 ml of 6–20% (w/v) sucrose in 0.01 M-Tris-HCl buffer, pH 8.0; 3 × 5 ml SW rotor, MSE SS50 centrifuge; 40000 rev./min, 150 min, 20°C]. Fractions (6 drops) were analysed by Čerenkov counting (Clausen, 1967) and the labelled DNA was found at or near the bottom of the centrifuge tubes and appropriate fractions were dialysed, concentrated in a vacuum desiccator (in 'Repelcoated' tubes) and the solutions digested with pancreatic deoxyribonuclease or endonuclease I from *E. coli* (Lehman *et al.*, 1962). DNA recovered by centrifugation or gel filtration had essentially its native structure and was far more readily digested by the endonucleases than were preparations that had been recovered by acid precipitation. With these samples, the conditions of digestion could be adjusted fairly easily and reproducibly to give larger nucleotides than were obtained from the acid-precipitated DNA samples, which always required high enzyme concentrations (0.2–5 mg/ml) and seldom yielded 5'-terminal fragments bigger than tetranucleotides. The concentrated DNA solutions recovered from sucrose gradients were digested in 0.02 M-Tris-HCl buffer, pH 7.5, 5 mM-MgCl₂ at room temperature (23°C) or at 37°C for periods varying from 30 min to several hours and pancreatic deoxyribonuclease concentrations between 0.1–0.005 mg/ml. Nucleotides from these digests were hydrolysed further, with or without prior fractionation, as described here and previously (Murray, 1970).

Ionophoresis and radioautography. Details of these operations were as described previously (Sanger *et al.*, 1965; Murray, 1970). In some cases, fractionation was achieved by ionophoresis on AE-cellulose paper at pH 3.5 followed by second-dimension ionophoresis on DEAE-cellulose paper at pH 2.

Analysis of nucleotides. (i) Complete and partial digestions with venom phosphodiesterase. These were conducted as described by Murray (1970). In some cases, partial digests were divided into three parts for ionophoresis on (a) DEAE-cellulose paper at pH 2, (b) AE-cellulose paper at pH 2, and (c) AE-cellulose paper at pH 3.5. Complete digests were analysed by ionophoresis on AE-cellulose (AE-81) paper at pH 3.5, because in this system ATP remains close to the origin whereas the four mononucleotides are quite satisfactorily resolved when the blue dye (Xylene Cyanol FF) has migrated 25 cm from the origin. Nucleotides were detected by radioautography and, where necessary, quantitated by liquid-scintillation counting for radioactivity with 0.4% (w/v) 2,5-bis-(5-*t*-butylbenzoxazol-2-yl)thiophen (BBOT) in toluene. When very small quantities of radioactivity were involved, reference markers (10 mg/ml; 10 μ l) were run coincidentally with the samples and detected under u.v. light ('chromatVue';

Ultraviolet Products Inc., San Gabriel, Calif., U.S.A.); appropriate zones of the paper were then cut out and counted for radioactivity with an automatic low background, anticoincidence, gas-flow counter (Nuclear-Chicago Ltd., High Wycombe, Bucks., U.K.).

(ii) Analysis via phosphorylation of partial spleen phosphodiesterase digests. Nucleotides were dephosphorylated, digested with spleen phosphodiesterase, rephosphorylated with [γ -³²P]ATP and polynucleotide kinase, and the components were separated by ionophoresis and analysed by digestion with venom phosphodiesterase. The sequence of operations was done on a micro scale as follows. The nucleotide was dissolved in water (about 20 μ l), incubated with bacterial alkaline phosphatase solution (0.5 mg/ml, 2 μ l; 37°C, 30 min), mixed with 0.1 M-sodium acetate buffer, pH 5.1 (containing EDTA and Tween 80) (2 μ l), spleen phosphodiesterase solution (about 1 mg/ml; 2 μ l) and kept at 20°C for about 45 min. EDTA (0.2 M, pH 7.4; 2 μ l) and 1 M-HCl (1 μ l) were then added and after 10 min at room temperature the solution was neutralized with 1 M-Tris-HCl buffer, pH 7.5 (5 μ l), and 1 M-NaOH (1 μ l). Polynucleotide kinase reaction mixture was then added [e.g. 10 μ l of the following mixture: 20 μ l of 1 M-MgCl₂; 25 μ l of 1 M-Tris-HCl buffer, pH 7.5; 50 μ l of 0.1 M-2-mercaptoethanol; 2 μ l of [γ -³²P]ATP of very high specific radioactivity (1–20 mCi/ μ mol); 25 μ l of polynucleotide kinase (fraction VI)] and incubated at 37°C for about 1–4 h. The reaction mixture was divided into two parts for ionophoretic fractionation of the labelled nucleotides on DEAE-cellulose paper (pH 2) and AE-cellulose paper (pH 3.5); the latter was usually the more useful because the excess of ATP from the reaction remained near the origin. Radioactive nucleotides were detected by radioautography, eluted with 30% (v/v) triethylamine carbonate solution, and analysed by digestion with venom phosphodiesterase; normally only complete digests were necessary.

(iii) Analysis by 'diagonal ionophoresis'. In some instances digestion with venom phosphodiesterase was done on the paper (Murray, 1970). On DEAE-cellulose paper a venom phosphodiesterase concentration of about 0.1 mg/ml (about 7 h at 37°C) was usually satisfactory, but on AE-cellulose paper a concentration of about 0.25 mg/ml was preferable. After incubation in a humid atmosphere the paper strips were dried and sewn on to a sheet of the same type of paper for ionophoresis at right angles to that used in the first separation.

Results and Discussion

Routine determination of nucleotide sequences via a terminal-labelling reaction and nucleotide 'mapping' methods depends upon the fulfilment of two

conditions. One is that the sequence of the terminally labelled nucleotides must be obtainable uniquely from mobility differences between the products of a single exonuclease digest. This required the development of new systems giving mobility shifts characteristic of G and T residues, the key to which was provided by previous analyses of nucleotide mixtures on AE-cellulose paper (Murray, 1970). The other is that the terminal-labelling reaction must be quite general, and reproducible. This appeared highly probable from the work of Richardson (1965) and Novogrodsky & Hurwitz (1966), and was established through a comparison of nucleotide 'maps' of pancreatic deoxyribonuclease digests of two DNA preparations from bacteriophage λ , one of which was made from uniformly ^{32}P -labelled bacteriophage and the other from non-radioactive bacteriophage, the digest being dephosphorylated with bacterial alkaline phosphatase and then labelled by means of the polynucleotide kinase reaction. These nucleotide 'maps' are shown in Plate 1, and analyses of nucleotides from one of the 'maps' of terminally labelled nucleotides (Plate 1*d*) are given in Table 2. The general features of the nucleotide 'maps' of the terminally labelled and uniformly labelled digests are qualitatively very similar; the relatively higher proportion of larger, slow-moving nucleotides in the uniformly labelled digest is partly an illusion arising from the higher proportion of labelled phosphorus atoms in the nucleotides in this digest, and partly the result of less-efficient digestion of the DNA with deoxyribonuclease in this particular experiment.

This result showed that nucleotide 'maps' made with the polynucleotide kinase reaction would be representative of the nucleotides present in the digest and so might provide a basis for 'mapping' nucleic acids from sources such as higher plants and animals which cannot be labelled *in vivo* as readily as prokaryotic systems. This is a much-simpler, more-useful, and far more-sensitive method for comparative work than the neutron-activation procedure used to compare nucleotide 'maps' of DNA from calf thymus and *E. coli* (Murray & Offord, 1966). Virtually identical nucleotide 'maps' of the type shown in Fig. 1 and Plate 2 were obtained with terminally labelled digests (pancreatic deoxyribonuclease) of DNA from bacteriophages λ and T_7 and from calf thymus and chicken erythrocyte.

The nucleotides in a terminally labelled pancreatic deoxyribonuclease digest of bacteriophage λ DNA were used to extend investigations of nucleotide 'maps' by two-dimensional ionophoresis on DEAE-cellulose paper (pH9.7 followed by pH2.0). This system was used because of its high resolving power and capacity. Complete digestion with venom phosphodiesterase gave the 5'-terminus of the nucleotides, showing that several of the nucleotides were mixtures. Ionophoretic analysis of partial digests with venom

phosphodiesterase was used for sequence analysis on these nucleotides. Mobility shifts (Sanger *et al.*, 1965) between related nucleotides in a partial digest (particularly where separations were effected in parallel on DEAE-cellulose and AE-cellulose papers, as discussed further below), together with the identification of the radioactive mononucleotide present, gave the results listed in Table 3 (and Fig. 1 and Plate 2). These results show the clear resolution of isomers obtainable in many instances with this particular two-dimensional system. Similar procedures were used to identify terminally labelled nucleotides from the same pancreatic deoxyribonuclease reaction mixture, but fractionated by ionophoresis on AE-cellulose paper at pH3.5, followed by a second ionophoresis on DEAE-cellulose paper at pH2. This system (Fig. 2) was used because no dinucleotide moves ahead of the blue marker on AE-cellulose paper at pH3.5 and the nucleotides C and G have a greater retarding effect on ionophoretic mobility on AE-cellulose paper than on DEAE-cellulose paper; it is particularly useful for the fractionation of mixtures of small nucleotides (for example in analyses of alkaline hydrolysates of RNA for 2'-*O*-methyl nucleotides), almost all the dinucleotides, for example, being resolved. Analyses of nucleotides separated in this way are given in Table 4.

Deduction of the sequence of terminally labelled nucleotides from ionophoretic analyses of their products on partial hydrolysis proved to be more reliable than with uniformly labelled nucleotides. The success of this analysis rests on the use of both AE-cellulose and DEAE-cellulose papers for ionophoretic fractionation of the partial digests. (Ionophoresis on AE-cellulose paper at pH3.5 frequently showed that nucleotides obtained from endonuclease digests that had been fractionated by two-dimensional ionophoresis on cellulose acetate and DEAE-cellulose paper, or DEAE-cellulose paper at two different pH values, were mixtures, often of isomers.) On DEAE-cellulose paper at pH2, Sanger *et al.* (1965) showed that loss of a C residue from an oligoribonucleotide gives a value for M (defined as the ratio of the distance between the two nucleotides to the distance moved from the origin by the slower, i.e. larger of the two nucleotides) of 0.1–0.3, loss of an A residue gives a value for M of 0.5–1.0, and loss of either a G or a U residue (T behaves as U) gives a value in the range of about 1.5–2.5. These values have proved quite reliable, in general, and apply in the deoxy series as with ribonucleotides. In some cases, however, abnormally low values have been found with C residues and sometimes nucleotides differing in size by one C residue are not resolved; also the ambiguity between G and T residues imposes a serious limitation on the analysis of nucleotides in, for example, pancreatic deoxyribonuclease digests of DNA (Murray, 1970). The relative effect of C and G residues on ionophoretic behaviour

EXPLANATION OF PLATE I

Nucleotide 'maps' of pancreatic deoxyribonuclease digests of bacteriophage λ DNA

(a) and (b) were made with uniformly labelled DNA and (c) and (d) with digests that had been labelled by the polynucleotide kinase reaction. Ionophoresis on cellulose acetate, pH 3.5, followed by DEAE-cellulose paper, pH 2, was used for fractionation of digests shown in (a) and (c), and cellulose acetate, pH 3.5, followed by AE-cellulose paper, pH 2, was used for those shown in (b) and (d). The line drawing (dii) is a key to (di). Analyses of nucleotides from (d) are given in Table 2.

EXPLANATION OF PLATE 2

Terminally labelled nucleotides from a pancreatic deoxyribonuclease digest of bacteriophage λ DNA resolved by two-dimensional ionophoresis on DEAE-cellulose paper

The same ionophoresis system as that in Fig. 1 was used, but it was run further in the second dimension to enhance resolution. Analyses of these nucleotides are in Table 3. The line drawing (ii) is a key to (i).

K. MURRAY

EXPLANATION OF PLATE 3

Diagonal ionophoresis method for partial digests with venom phosphodiesterase

Patterns from (a) a homogeneous heptanucleotide and (b) a mixture of two pentanucleotides with the same 5'-terminus. (c) and (d) are patterns from a pancreatic deoxyribonuclease digest of 5'-terminally labelled DNA from bacteriophage λ . In (a) and (b) the partial exonuclease digest was fractionated on DEAE-cellulose paper, pH 2, and after a second partial digestion with venom phosphodiesterase had been done on the paper, the strip of paper was sewn on to another sheet of DEAE-cellulose paper for ionophoresis in a direction at right angles to the first ionophoresis. In (c) and (d) the pancreatic deoxyribonuclease digest was fractionated by ionophoresis on (c) AE-cellulose paper, pH 3.5, and (d) DEAE-cellulose paper, pH 2. Each strip of paper was then treated with venom phosphodiesterase (details are in the text) and sewn on to another sheet of the same paper for the second ionophoresis, which was at right angles to the first. The line drawings (c ii) and (d ii) are the keys for (c i) and (d i) respectively.

EXPLANATION OF PLATE 4

Nucleotide 'maps', made by two-dimensional ionophoresis on DEAE-cellulose paper (pH 9.7 followed by pH 2), of pancreatic deoxyribonuclease digests of 5'-terminally labelled DNA from (a) bacteriophage λ and (b) bacteriophage 424.

For details see the text. The line drawings (a ii) and (b ii) are the keys of (a i) and (b i) respectively.

For details see the text. The line drawings (a ii) and (b ii) are the keys for (a i) and (b i) respectively.

K. MURRAY

Table 2. 5'-Terminus and base sequence in nucleotides resolved by two-dimensional ionophoresis on cellulose acetate (pH3.5) and AE-cellulose paper (pH2) (see Plate 1d)

Nucleotides eluted from the ion-exchange paper with triethylamine carbonate were divided into two parts approximately in the ratio 1:5. The smaller part was used for complete digestion with venom phosphodiesterase (0.1 mg of enzyme/ml) and the resulting 5'-mononucleotides were separated and identified by ionophoresis on Whatman no. 52 paper or AE-cellulose paper at pH3.5 followed by radioautography. The larger portion was used for partial digests with venom phosphodiesterase (0.01 mg of enzyme/ml). One-half of each digest was fractionated on AE-cellulose paper, pH3.5, and the other half on DEAE-cellulose paper, pH2, and the papers were radioautographed. Nucleotide sequences were then deduced from *M* values measured on the two papers. Underlining indicates that this nucleotide was the major component in mixtures.

Nucleotide no.	5'-Terminus	Nucleotide sequence	Nucleotide no.	5'-Terminus	Nucleotide sequence
3	G, <u>T</u>		40	A	
5	A		41	A	A-G-A
6	T	T-G-G-G	42	C, G, <u>T</u>	<u>T-G-C</u>
7	G, T	G-G-T-T			<u>T-C-G</u>
8	G, T				G-C-T
9	G, T	G-T-T-T	43	G	G-G
10	<u>T</u>	T-T-T-T	44	C	
12	G, T		45	A	A-T-A
13	<u>G</u>		46	A, <u>T</u>	T-A-A
15	C, A		47	C	C-G-C-C
16	G, T		48	G	
17	<u>G</u> , T	G-G-C-T (+isomers)	49	C, G, <u>T</u>	G-C-A
18	A, G, <u>T</u>	T-G-A-G	50	<u>A</u> , <u>T</u>	A-G-C
19	G	<u>G-T-A-T</u>			A-C-G
		<u>G-A-T-T</u>	51	T	T-C-T
20	T	T-A-T-T	52	G, <u>T</u>	<u>T-G</u>
21	G	G-G-C-A			<u>G-T</u>
22	C, G, <u>T</u>	T-G-C-G	54	C	
		G-C(G,T)	55	G	
23	G, T	G-G-T	56	<u>A</u> , T	A-C-T
		T-G-G			A-T-C
24	C	Mixture	57	G	T-C-A
25	G	G-G-C-C	58	T	G-A
26	A	A-G-G	59	C, <u>A</u>	T-T
27	G	G-G-A	60	C	C-G-C
28	T	T-G-T	61	A	
29	T	T-T-T	62	C	C-G
30	C, A		63	G	G-C
31	C, <u>A</u> , G		64	A	A-T
32	<u>C</u> , G	C-G-G	65	T	T-A
33	G	<u>G-C-G</u>	66	C	
34	A	A-T-G	67	C	C-A
35	A	A-G-T	68	C	C-C
36	T	T-G-A	70	<u>C</u> , T	<u>C-T</u>
37	A, T	<u>A-T-T</u>			<u>T-C</u>
		T-T-A	71	C	
38	C, A, G		73	A, <u>T</u>	
39	<u>A</u>		74	<u>C</u> , <u>A</u>	

phosphodiesterase digests of 5'-labelled nucleotides thus gives appreciable information about the base sequence of the nucleotides and usually, particularly with smaller nucleotides, the sequence can be deduced unequivocally. In other cases, and especially with some of the larger oligonucleotides (such as may be obtained from an alleged mRNA, for the method is equally applicable to oligoribonucleotides; Szekely & Sanger, 1969) further evidence is necessary to determine whether the nucleotide under examination is homogeneous or is a mixture of two or more nucleotides having a common 5'-terminus. Two methods can be used for this purpose. The simpler involves further partial digestion with venom phosphodiesterase of the products of the first digestion and examination of the products from each component of the first digest: this may be achieved simply by carrying out the second digest on the strip of DEAE-cellulose paper (Murray, 1970) followed by ionophoresis in a direction at right angles to the first ionophoresis after stitching the strip of DEAE-cellulose (or AE-cellulose) paper to another sheet of the same paper. The grid-diagonal patterns that would be produced from a homogeneous heptanucleotide and from a mixture of two pentanucleotides giving similar one-dimensional patterns are represented in Plate 3(a) and 3(b). Each of the products from a homogeneous nucleotide will give all of the faster-migrating nucleotides on further partial digestion with the exonuclease, so that the pattern of spots seen on a radioautograph form a complete 'pyramid' (Plate 3a). Each product from a partial digest of a mixture of nucleotides will not produce all of the progressively smaller nucleotides already present when digested further with the exonuclease so that the pattern of spots does not form the complete 'pyramid', but leaves a number of gaps (Plate 3b). The example shown in Plate 3(b) is nucleotide number 24 from the 'map' shown in Fig. 1 and Plate 2, which proved to be a mixture of the pentanucleotides A-T-A-A-A and A-G-A-A-A; the sequence deduced for a homogeneous heptanucleotide that would have given the pattern shown in Plate 3(a) would have been A-T-C-A-C-A-A. Pancreatic deoxyribonuclease digests of 5'-terminally labelled DNA from bacteriophage λ provide a useful and interesting example for the application of the diagonal method because they comprise two different families of nucleotides, each having progressively overlapping sequences from the 5'-terminus (this is discussed in more detail below). These digests are thus equivalent to a partial exonuclease (venom) digest of a mixture of two 5'-terminally labelled oligonucleotides. Plate 3(c) and 3(d) shows the patterns obtained when pancreatic deoxyribonuclease digests of terminally labelled DNA from bacteriophage λ were separated on AE-cellulose paper (Plate 3c) and DEAE-cellulose paper (Plate 3d), treated with venom phosphodiesterase on the paper, and ionophoresis

was then repeated as in the first separation, but in a direction at right angles. The presence of nucleotides from two different families is readily apparent from the pattern of spots moving off the diagonal [the patterns are of the type shown in Plate 3(b) and not of the type in Plate 3(a)], and mobility differences between related nucleotides when calculated from the two experiments (i.e. AE-cellulose paper and DEAE-cellulose paper) enabled the sequences illustrated to be determined. Careful examination of the diagonal on AE-cellulose paper (Plate 3c), for example, shows the nucleotide A-G-G appearing as a digestion product (i.e. as a product of the venom phosphodiesterase digest carried out on the paper after the first ionophoretic separation) in three positions on the second dimension. These positions correspond with nucleotides separated on the first dimension with mobility changes (*M* values) of 0.5 and 3.0 and, therefore, differing in sequence successively by C and G residues. They correspond with the sequences for their parent nucleotides of A-G-G-T, A-G-G-T-C and A-G-G-T-C-G. The second procedure for studying the homogeneity of 5'-terminally labelled nucleotides involves dephosphorylation of the nucleotide followed by partial digestion with spleen phosphodiesterase to furnish a mixture of nucleotides, each of which is a substrate for polynucleotide kinase since it has a free 5'-hydroxyl group. After rephosphorylation with polynucleotide kinase the mixture of nucleotides is fractionated by ionophoresis and analysed by hydrolysis of its constituents with venom phosphodiesterase. This method is outlined in Scheme 1. Since all the nucleotides present form an overlapping series the sequence of the parent nucleotide, if homogeneous, can be confirmed in all positions except the 3'-terminus, or the presence of a mixture will usually be apparent. These procedures have been applied successfully to some small oligonucleotides.

The more interesting applications of the polynucleotide kinase reaction involve the determination of nucleotide sequences in regions of DNA molecules which can be related to some function of biological significance. Examples of nucleotide sequences that are recognized by specific proteins include the regions involved in integration of a prophage into its host chromosome (lysogenization), host-controlled modification and restriction, and the cohesive ends of DNA from bacteriophage λ and related bacteriophages (Hershey, 1971). Nucleotide sequences at the ends of bacteriophage λ DNA have been studied through repair reactions with DNA polymerase by Wu & Kaiser (1968). This approach, which required, *inter alia*, very highly purified deoxynucleoside triphosphates and a precise knowledge of their specific radioactivity, suggested that the 'left-hand' end of the DNA molecule has the sequence pG-G-G-G-C followed by a somewhat ambiguous sequence of three G and two C residues, and that the 'right-hand'

Table 3. 5'-Terminus and base sequence in nucleotides resolved by two-dimensional ionophoresis on DEAE-cellulose paper (pH9.7 followed by pH2) as illustrated in Fig. 1 and Plate 2

For details see Table 2.

Nucleotide no.	5'-Terminus	Nucleotide sequence	Nucleotide no.	5'-Terminus	Nucleotide sequence
4	T	T-G-C-G (+others)	49	<u>A</u> , G, T	A-T-C-G (+isomers)
7	T	T-T-T	50	G	G-T-C-A
10	G	G-T-C-G (+others)	51	C, <u>G</u> , T	
13	G	G-G-G	53	T	T-C-T-C
14	G, T	G-G-T	54	T	T-T-C-C
15	<u>G</u> , T	T-G-G	55	A, <u>T</u>	T-A-A-A
16	T	T-T-T-C	56	T	T-C-T
17	T	T-T-G	58	C, T	
18	G	G-T-T	59	G	G-C-T-C
19	T	T-G-T	60	G	G-C-C-T
23	A		61	C	C-G-T-C
24	A	A-G-A-A-A A-T-A-A-A	62	T	<u>T-G-C-C</u>
24A	A		62A	A, G	
25	G		63	G	G-G-C
25A	A, G	G-A-T-A	64	G	G-C-T
26	A, <u>T</u>	T-A-T-A	65	C, G	C-G-T
26A	A, <u>T</u>	T-A-A-G A-A-T-G	66	T	G-T-C
27	T	T-G-A-A <u>T-A-G-A</u>	67	G	T-G-C
28	<u>A</u> , T	A-A-T-T <u>T-T-A-A</u> T-A-A-T	69	G	G-T
32	A		70	C, G	
32A	G, <u>T</u>	T-T-C-A <u>G-A-C-T</u> G-T(C,A)	72	G	G-C-G
35	A, G, T		73	C	C-G-G
36	C, <u>T</u>	T-C-A-T <u>T-A-C-T</u> C-A-T-T	74	G, T	T-C-G
36A	T	T-A-T-C	75	G	G-G
37	G	G-G-C-A	76	T	T-G
38	T	T-C-G-A (+isomers)	79	A	A-A-G-C
39	<u>T</u>	<u>T-G-C-A</u>	85	<u>A</u> , G	
39A	A	A-G-G	86	<u>A</u>	A-A-T-C
40	G	G-G-A	89	C	
41	G	G-T-A	90	A	A-G-C-A
41A	A	A-T-G	91	G	G-A-A
42	A	A-G-T	92	A	A-A-T
43	T	T-G-A	93	<u>A</u> , G	A-G-A-C
44	A, T	A-T-T	94	G	G-C-A-A
45	T	T-T-A	95	C, <u>T</u>	G-A-C-A
		T-A-T	95A	C, <u>T</u>	T-A-C-A
			96	<u>A</u>	C-A-T-A
			97	T	A-A-G
			100	A	T-A-A
			101	A	A-T-C-A
			102	C, T	A-G-A
			103	<u>A</u>	
			103A	A	A-T-A
			106	G	A-G-C-C
			107	C	G-A-C-C
					C-A-T-C
					C-A-C-T

Table 3 (continued)

Nucleotide no.	5'-Terminus	Nucleotide sequence	Nucleotide no.	5'-Terminus	Nucleotide sequence
107A	A	A-A-A-A	130	C	
108	T	T-A-C-C	131	C	
109	A	A-G-C	132	G	
110	A	A-C-C-G	133	A, C	
111	A, G	A-C-G-C	134	C	
		G-C-C-A	135	A	
112	C, G		136	T	T-C-C
113	C	C-G-C-A	138	T	T-A
113A	C, G		138A	A	A-T
114	T	T-C-C-A	139	G	G-C
115	A	A-T-C-C	146	T	T-T-C
117	C, A		147	C	
118	C	C-A-G	150	C	C-C-G
119	C		151	G	G-C-C
121	G	G-C-A	152	C	C-G-C
122	A	A-C-G	154	A	
123	C	C-G-A	155	C	C-G
124	T	T-C-A	156	C	C-A-A
125	A	A-T-C	161	A	A-C-A
126	G	G-A	164	A	A-C-C
127	C		200	C	C-C-A
128	A	A-G	201	C	C-A
129	T		202	C	C-C-C

5'-end of the DNA molecule is terminated by the sequence pA-Y. These conclusions, however, depended on the specific radioactivity and purity of the deoxynucleoside triphosphates used in the polymerase reactions. Subsequent work where the radioactively labelled product of the polymerase repair reaction was analysed in more detail (Wu, 1970) showed this result to be appreciably in error (possibly because the nucleoside triphosphates were insufficiently pure and their specific radioactivities were incorrect), and has extended the sequence determined to the whole of the cohesive ends of the bacteriophage λ DNA molecule (Wu & Taylor, 1971).

Initial attempts to study the 5'-terminal sequences of bacteriophage λ DNA were made with [γ - ^{32}P]ATP of specific radioactivity of about 10^3 mCi/mmol and gave the dinucleotides pA-G and pG-G. To extend these sequences and to assess the utility of the combination of 5'-terminal labelling and nucleotide 'mapping' as a means for the determination of sequences directly (as opposed to polymerase repair reactions), DNA preparations from bacteriophage λ were dephosphorylated and labelled at their 5'-termini with ATP of very high specific radioactivity (about 10^4 mCi/mmol). In experiments of this sort a completely general 5'-terminal-labelling reaction is being used to look specifically for sequences at the ends of molecules with a chain length of about

5×10^4 residues. If the ATP has a specific radioactivity of about 10^4 mCi/mmol, then about 3×10^4 c.p.m./ $50 \mu\text{g}$ of DNA should be obtained at each 5'-terminus of bacteriophage λ DNA. Any contaminating oligonucleotides (fragments of bacteriophage or host DNA, or host RNA) will be labelled also, and if they have a low chain-length then even at very low concentrations they may well contribute more labelled nucleotides to the reaction products than will the DNA being examined. For example, in a bacteriophage λ DNA sample containing as little as 0.1% by weight of oligonucleotides with a chain-length of about 50, the oligonucleotides would contribute terminally labelled nucleotides quantitatively equivalent to those of the DNA. Obviously it is essential that the ATP and enzyme preparations used, as well as the DNA to be labelled, be purified as far as possible, and further purification of the terminally labelled DNA after the reaction is usually necessary. For this purpose, sedimentation through sucrose gradients has proved very satisfactory.

The terminally labelled DNA preparations were digested with pancreatic deoxyribonuclease or endonuclease I from *E. coli*. Similar experiments were done with DNA from bacteriophage 424. Nucleotide 'maps' of such digests made by two-dimensional ionophoresis on DEAE-cellulose paper are shown

Fig. 2. Two-dimensional ionophoretic separation (AE-cellulose paper, pH 3.5, followed by DEAE-cellulose paper, pH2) of terminally labelled nucleotides from a pancreatic deoxyribonuclease digest of bacteriophage λ DNA

Analyses of these nucleotides are given in Table 4. The drawing is a composite of several experiments.

in Plate 4. Analyses of partial digests with venom phosphodiesterase (Plate 5) of some of the nucleotides from these 'maps' proved that corresponding nucleotides from the two DNA preparations are identical. These examples provide a useful illustration of the use of the two ionophoretic systems (DEAE-cellulose paper at pH2 and AE-cellulose paper at pH3.5) to solve sequences uniquely. Nucleotides 8 and 10 are not differentiated on DEAE-cellulose paper (Plate 5b) and the first product from each behaves identically with nucleotide 15 and has a mobility (M) value of 2.1 and therefore involves the loss of a G or T residue from the parent nucleotide. The next product from nucleotides 8 and 10, and the

first product from nucleotide 15, has an M value of 1.7 (and the mobility of A-G) and so loss of a G or T residue is again involved, and in each case the mononucleotide located is an A residue. On AE-cellulose paper, however, nucleotides 8 and 10 are clearly separated (Plate 5a) and the first product from nucleotide 8 is identical with nucleotide 10 and has an M value of 0.65 indicating loss of a C residue from nucleotide 8. The second product from nucleotide 8 and the first from nucleotide 10 are identical, with M values of 1.4 corresponding to the loss of a T or A residue, and the next product in each case (not visible in the illustration) has an M value of 2.5 (and the mobility of A-G), which can only arise from the loss of a G residue. Thus the sequence of nucleotide 8 can only be A-G-G-T-C and that of nucleotide 10 is A-G-G-T. The other sequences shown in Plates 4 and 5 were deduced in the same way.

Provided that the preparations of DNA, [γ - 32 P]-ATP and polynucleotide kinase used in this type of experiment had been sufficiently purified, all the radioactive nucleotides derived from an endonuclease digest of 5'-terminally labelled DNA must belong to one of two families representing the two terminal sequences. The nucleotides therefore provide two series of progressively overlapping sequences and since their separation had been effected by two-dimensional ionophoresis on DEAE-cellulose paper at pH9.7 followed by pH2, one set of M values has been obtained in the separation process if it is known how the nucleotides are grouped, i.e. if their 5'-terminus is known. [In the case of DNA from bacteriophages λ and 424 (and others) this is easily found because the two 5'-terminal nucleotides differ; in some other cases the terminal nucleotides are the same which obviously complicates the analysis.] It is then only necessary to analyse the partial digestion products of the nucleotides on AE-cellulose paper at pH3.5 to deduce the sequences uniquely. This is the procedure now followed and has been used to determine terminal sequences of DNA from several other bacteriophages (K. Murray and N. E. Murray, unpublished work), and is particularly useful with small quantities of radioactivity (e.g. <100 c.p.m. per nucleotide).

The results obtained for the 5'-terminal sequences of bacteriophage λ DNA by this direct method are the same as those obtained by repair syntheses with DNA polymerase (Wu & Taylor, 1971). Bacteriophage 424 DNA has 5'-terminal sequences identical with those of bacteriophage λ , at least to the extent of 11 out of the 12 nucleotides in the cohesive ends of bacteriophage λ DNA. Baldwin *et al.* (1966) have shown that DNA from bacteriophages λ and 424 have similar cohesive sites because they will form mixed dimers. The terminal sequences of these two DNA molecules would therefore be expected to be similar, but it is interesting to see that these sequences, which lie

Table 4. 5'-Terminus and base sequence of nucleotides resolved by two-dimensional ionophoresis on AE-cellulose (pH3.5) and DEAE-cellulose (pH2.0) papers (see Fig. 2)

For details see Table 2.

Nucleotide no.	5'-Terminus	Nucleotide sequence	Nucleotide no.	5'-Terminus	Nucleotide sequence
1	G	G-G-T	27	A, T	A-C-T
2	G, T	G-T-T			A-T-C
		T(T,G)			T-C-A
3	T	T-T-T	28	A	A-G
4	G	G-G-A	29	G	G-A
5	G	G-T-A	30	G	
6	A, T	T-A-G	31	C	
		T-G-A	32	A	
		A-G-T	33	G	G-C-C
		A-T-G	34	C	
7	T	T-A-T	35	C	C-G-C
		T-T-A	36	A	A-T
8	G	G-C-G	37	T	T-A
		G-G-C	38	A	
9	G		39	G	G-C
10	C, A		40	C	C-G
11	G	G-C-T	41	C, A	
12	G	G-G	42	C, A	
13	T	T-C-T	43	C	C-A-A
14	G	G-T	44	A	A(C,A)
15	T	T-G	45	A	A(C,A)
16	A, T		46	A	A-A
17	A, G, T		47	C, T	C-T
18	A		48	C, T	T-C
19	A	A-A-G	49	C	
20	A	A-A-T	50	C, A	A-C-C
21	T	T-A-A			C(C,A)
22	T	T-T	51	C	C-C-C
23	A	A-G-A	52	C	C-A
24	C, G		53	C	C-C
25	C, T				
26	G	G-C-A			

Table 5. *M* values for 5'-deoxynucleotides on AE-cellulose paper at pH3.5

For details see the text.

3'-Terminal nucleotide	Range of <i>M</i> values
A	0.9-1.5
C	0.5-0.7
G	2.0-3.0
T	1.0-1.5

within the breaks produced by the action of the *ter* gene product (Mousset & Thomas, 1969) are almost, if not completely, identical.

The extent of sequence that can be determined by

the procedures described here is limited by the ionophoretic mobilities of the labelled nucleotides, and by the number of nucleotides that can be separated from partial endonuclease digests of terminally labelled DNA preparations. The DNA from bacteriophages λ and 424 gives a particularly unfavourable series of 5'-terminal nucleotides in that each terminus comprises an exceptionally high proportion of G and/or T residues. These impart very low mobilities to relatively small oligonucleotides, so that even with very prolonged ionophoresis (runs of up to 3 days have been used) of partial exonuclease digests the nucleotides do not move far enough to give reliable *M* values. A higher proportion of C or A residues near the 5'-termini would certainly have permitted the determination of longer sequences.

Scheme 1. Scheme for confirmatory analysis of a 5'-terminally labelled oligonucleotide

For details see the text. * denotes ^{32}P . W, X, Y and Z represent unspecified nucleosides.

For separation of some of the larger oligonucleotides derived from terminally labelled DNA, the two-dimensional procedures involving ionophoresis on cellulose acetate and chromatography on thin layers of cellulose impregnated with polyethyleneimine described by Southern & Mitchell (1971) have proved very useful, although analysis of some of these larger nucleotides is not trivial.

I am indebted to Dr. A. D. Kaiser and Dr. P. Berg for initiating and encouraging my interest in studies of the structure of bacteriophage λ DNA and applications of the polynucleotide kinase reaction, to the William Waldorf Astor Foundation for the award of a research travel grant (1966), to Mrs. J. S. Brown and Mr. T. B. Bruce for technical help, to Mr. S. G. Hughes for preparations of bacteriophage and bacteria, and to Dr. Noreen E. Murray for help with preparations of bacteriophages λ and 424. The bacteriophage strains were kindly provided by Dr. A. D. Kaiser and Dr. R. L. Baldwin. The work described was supported by a grant from the Science Research Council.

References

- Adams, M. M. (1959) *Bacteriophages*, Interscience Publishers, New York
- Baldwin, R. L., Barrand, P., Fritsch, A., Goldthwait, D. A. & Jacob, F. (1966) *J. Mol. Biol.* **17**, 343–357
- Burton, K. & Petersen, G. B. (1960) *Biochem. J.* **75**, 17–27
- Clausen, T. (1967) *Anal. Biochem.* **22**, 70–73
- Dahlberg, J. E. (1968) *Nature (London)* **220**, 548–552
- Glynn, I. M. & Chappell, J. B. (1964) *Biochem. J.* **90**, 147–149
- Hershey, A. D. (ed.) (1971) *The Bacteriophage Lambda*, Cold Spring Harbor Laboratories, New York
- Kaiser, A. D. & Hogness, D. S. (1960) *J. Mol. Biol.* **2**, 392–415
- Lehman, I. R., Roussos, G. G. & Pratt, E. A. (1962) *J. Biol. Chem.* **237**, 819–828
- Mousset, S. & Thomas, R. (1969) *Nature (London)* **221**, 242–244
- Murray, K. (1968) *Abstr. Chem. Soc. (London) Autumn Meet.* Abstract no. G5
- Murray, K. (1969) *J. Mol. Biol.* **39**, 125–144
- Murray, K. (1970) *Biochem. J.* **118**, 831–841
- Murray, K. & Offord, R. E. (1966) *Nature (London)* **211**, 376–378
- Murray, K. & Wu, R. (1968) *Biochem. J.* **108**, 42P
- Murray, K., Vidali, G. & Neelin, J. M. (1968) *Biochem. J.* **107**, 207–215
- Novogrodsky, A. & Hurwitz, J. (1965) *Fed. Proc. Fed. Am. Soc. Exp. Biol.* **24**, 602
- Novogrodsky, A. & Hurwitz, J. (1966) *J. Biol. Chem.* **241**, 2923–2932
- Richardson, C. C. (1965) *Proc. Nat. Acad. Sci. U.S.A.* **54**, 158–165
- Richardson, C. C. (1966) *J. Mol. Biol.* **15**, 49–61
- Richardson, C. C. & Weiss, B. (1967) *J. Mol. Biol.* **23**, 405–415
- Sanger, F., Brownlee, G. G. & Barrell, B. G. (1965) *J. Mol. Biol.* **13**, 373–398
- Southern, E. M. (1970) *Nature (London)* **227**, 794–798
- Southern, E. M. & Mitchell, A. R. (1971) *Biochem. J.* **123**, 613–617
- Szekely, M. & Sanger, F. (1969) *J. Mol. Biol.* **43**, 607–617
- Wittenberg, J. & Kornberg, A. (1953) *J. Biol. Chem.* **202**, 431–444
- Wu, R. (1970) *J. Mol. Biol.* **51**, 501–521
- Wu, R. & Kaiser, A. D. (1967) *Proc. Nat. Acad. Sci. U.S.A.* **57**, 170–177
- Wu, R. & Kaiser, A. D. (1968) *J. Mol. Biol.* **35**, 523–537
- Wu, R. & Taylor, E. (1971) *J. Mol. Biol.* **57**, 491–511