Relationship between Structure and the Insulin-Like Activity of the Neurohypophyseal Peptide Hormones

By E. SCHILLINGER, E. SCHRÖDER, E. KLIEGER and K. LUBKE. (*Schering AG, 11-Berlin 65, W. Germany*)

Little is known about the relationship between structure and insulin-like activity of the neurohypophyseal hormones. We therefore synthesized a number of analogues of vasopressin by using conventional methods or solid-phase peptide synthesis.

Among others, analogues with changes at the N-terminal amino group, with changes in the amino acid at position 3 and with changes in the amino acid and changes of the functional group at position 8 were synthesized. Since these positions of the molecule have a special influence on the vasopressor and oxytocic activities, their importance with respect to the insulin-like activity was of special interest.

All compounds were tested for their ability to stimulate the oxidation of [$1\text{-}^{14}\text{C}$]glucose to $^{14}\text{CO}_2$ in isolated fat-cells of the rat. The compounds tested were divided into groups of highly active, active and inactive substances.

Evidence for a structure-activity relationship was obtained, but, in comparison with the well-known relationship between structure and the vasopressor and oxytocic activities, only a few analogies could be seen. On the other hand, the differences show that a dissociation of the insulin-like activity from the vasopressor, antidiuretic and oxytocic activities is possible.

A Comparative Study of the Distribution and Metabolism of Monocomponent Human Insulin and Porcine Proinsulin in Man

By C. V. TOMPKINS, M. C. SRIVASTAVA, P. H. SÖNKSEN and J. D. N. NABARRO. (*Institute of Nuclear Medicine, The Middlesex Hospital Medical School, London W1N 7RL, U.K.*)

Insulin and proinsulin were infused (on separate occasions) by the 'priming-dose-constant-infusion' technique. The infusion rates were adjusted to produce steady-state concentrations of hormone in the blood that extended over a wide range. Proinsulin was infused at an equimolar rate to insulin. Plasma disappearance rate ($t_{1/2}$) was measured at the end of the infusion. Plasma glucose and serum proinsulin and growth hormone were measured every 5 min throughout the 3 h of study. Serum proinsulin and growth hormone were measured by a double-antibody radioimmunoassay. The pro-

insulin assay was not influenced by endogenous insulin or proinsulin.

The metabolic clearance rate, $t_{1/2}$ and apparent 'distribution space' were calculated by conventional techniques. The results for the group of normal subjects, based on the metabolic clearance rate measured at the highest infusion rate immediately before the measurement of $t_{1/2}$, were as follows: for insulin (nine experiments), metabolic clearance rate 13.3 (range 9.7–17.8) ml/min per kg body wt., $t_{1/2}$ 4.39 (range 3.00–5.75) min and 'distribution space' 83.0 (range 66.5–111.8) ml/kg body wt.; for pro-insulin (ten experiments), metabolic clearance rate 3.15 (range 2.52–4.31) ml/min per kg body wt., $t_{1/2}$ 25.6 (range 17.8–38.0) min and 'distribution space' 105.5 (range 62.7–147.6) ml/kg body wt.; P values <0.001 , <0.001 and $<0.1 > 0.05$ respectively. Thus for insulin the metabolic clearance rate fell progressively with increasing steady-state insulin concentrations in the blood, whereas for proinsulin the metabolic clearance rate remained constant throughout a wide range of blood hormone concentrations (3–100 ng/ml).

The biological potency of proinsulin, as measured by the fall in plasma glucose concentration, was approx. 5% that of insulin. It is clear that, although insulin and proinsulin appear to be distributed in approximately the same 'space', the metabolism of the two hormones is different.

The Importance of 'Distribution' in the Metabolism of Growth Hormone in Man

By M. C. SRIVASTAVA, D. OWENS, C. V. TOMPKINS, P. H. SÖNKSEN and J. D. N. NABARRO. (*Institute of Nuclear Medicine, The Middlesex Hospital Medical School, London W1N 7RL, U.K.*)

The metabolic clearance rate of human growth hormone has been studied by using a 'priming-dose-constant-infusion' technique. The metabolic clearance rate has been shown to remain constant at 3.1 ml/min per kg body wt. at blood concentrations in the range 3–50 ng/ml (Srivastava, Tompkins, Sönsken & Nabarro, 1970). The disappearance rate of growth hormone ($t_{1/2}$) from the blood was measured after discontinuation of the infusion; it was found to be monoexponential over the 60 min in which it was measured. Metabolic clearance rate was calculated as:

$$\frac{\text{Infusion rate of growth hormone (ng/min)}}{\text{Steady-state concn. of blood growth hormone (ng/ml)}}$$