Specific precipitation of polyribosomes, particularly from the 200S region of the gradient, was achieved in TMK buffer with F(ab')₂ fragments of the antibodies. Pretreatment with puromycin resulted in decreased precipitation of polyribosomes by specific F(ab')₂ fragments.

A rapidly labelled RNA was extracted from the precipitated polyribosome-antibody complexes. Its sedimentation coefficient was 11-12S and its base composition was 48mol of G+C/100mol. The size and base composition of this RNA are consistent with its being the mRNA for the immunoglobulin light chain synthesized by MOPC 149 plasmacytoma.

S.L.B. and T.L.D. held National Research Council of Canada Postgraduate Studentships. The Medical Research Council of Canada provided financial support.

Davis, B. K., Delovitch, T. L. & Schon, A. H. (1969). *Nature, Lond.*, **222**, 172.

Hartlieb, R. & Koningsberger, V. V. (1968). *Biochim. biophys. Acta*, **166**, 512.

Williamson, A. R. & Askonas, B. A. (1967). *J. molec. Biol.* **23**, 201.

The Occurrence of 5.8S Ribonucleic Acid in Plants

By P. I. PAYNE and T. A. DYER. (*Agricultural Research Council Unit of Developmental Botany, 181A Huntingdon Road, Cambridge CB3 0DY, U.K.*)

A 5.8S RNA species has been shown to be present in several flowering plants, a fern, a club-moss and a green alga, and has several properties in common with the corresponding molecule (also described as 7S, 25S A and 'I' RNA) from animal cells (Pene, Knight & Darnell, 1968; Sy & McCarty, 1970; King & Gould, 1970). First, plant and animal 5.8S RNA species are very similar in size. We found that the plant RNA species varied in molecular weight between 49800 (*Tussilago farfara*) and 51600 (*Selaginella kraussiana*) as determined by polyacrylamide-gel electrophoresis, the mean value being 50400. Secondly, it is associated with the largest (25S) rRNA in equimolar amounts. Thirdly, this complex may be dissociated by conditions that disrupt hydrogen bonds. With the broad-bean RNA we have shown that dissociation by heat occurs within a narrow temperature range (37-52°C in a buffer containing no Mg²⁺ and of low ionic strength). Similar findings have been reported for the animal RNA (Sy & McCarty, 1970; King & Gould, 1970). It is therefore probable that 5.8S RNA is a universal component of cytoplasmic 80S ribosomes.

The 5.8S RNA was recovered in equimolar amounts with 5S RNA from broad-bean roots, but in green leaves it was found in submolar amounts. The relative proportions of 5S and 5.8S RNA in

leaves of dark-grown plants were similar to those in roots, but when such plants were exposed to light, which stimulated the synthesis of chloroplast ribosomes, the proportion of 5.8S RNA progressively decreased until the relative amounts were comparable with those in leaves of light-grown plants. These observations suggested that chloroplast 70S ribosomes do not contain 5.8S RNA, and this finding was confirmed by heating and cooling the RNA fraction from purified chloroplasts prepared from broad-bean and spinach leaves.

As it has also been reported that 5.8S RNA could not be detected in *Escherichia coli* preparations (Pene *et al.* 1968; Sy & McCarty, 1970), we examined two other prokaryotes, the blue-green algae *Nostoc* and *Anabaena flosaquae*, and similarly could not detect this component. The 5.8S RNA is therefore probably not a component of the 70S class of ribosomes, but is restricted to eukaryotic 80S ribosomes. Our findings demonstrate a further similarity in chloroplast and prokaryotic ribosome structure and provide additional support for the endosymbiont theory of the origin of chloroplasts (Margulis, 1970).

King, H. W. S. & Gould, H. (1970). *J. molec. Biol.* **51**, 687.

Margulis, L. (1970). In *Origin of Eukaryotic Cells*. London: Yale University Press.

Pene, J. J., Knight, E. & Darnell, J. E. (1968). *J. molec. Biol.* **33**, 609.

Sy, J. & McCarty, K. S. (1970). *Biochim. biophys. Acta*, **199**, 86.

Mutagenicity of 'K-Region' Epoxides of Polycyclic Hydrocarbons towards Bacteriophage T₂

By M. J. COOKSON, P. L. GROVER and P. SIMS. (*Chester Beatty Research Institute, Institute of Cancer Research: Royal Cancer Hospital, Fulham Road, London SW3 6JB, U.K.*)

The microsomal metabolism of polycyclic hydrocarbons is now known to involve the formation of epoxide intermediates that are probably 'K-region' derivatives (Sims, Hewer & Grover, 1971). Synthetically prepared 'K-region' epoxides have been found to be alkylating agents that react with nucleic acid either *in vitro* (Grover & Sims, 1970) or in cells in culture (Grover, Forrester & Sims, 1971). Polycyclic hydrocarbon epoxides are therefore particularly interesting as potential mutagens, and we have now tested a series of these compounds for activity in producing plaque-type and host-range mutations of bacteriophage T₂ (Loveless, 1959).

In the host-range-mutagenesis assay, where buffered suspensions of bacteriophage T₂ were