Proteins that Bind Carcinogen Metabolites

By B. KETTERER (*Courtauld Institute of Biochemistry, The Middlesex Hospital Medical School, London W1P 5PR, U.K.*)

During the course of their metabolism some carcinogens have been shown to give rise to alkylating agents that react with proteins in the target tissue (Magee & Hultin, 1962; Miller & Miller, 1969). In the case of the amino-azo-dye carcinogens and rat liver, proteins in all subcellular fractions may become arylated, but those of the endoplasmic reticulum and soluble supernatant fraction account for most of the covalently bound amino-azo-dye. Proteins of the soluble supernatant have been studied most intensively, and here it is found that only a limited number of protein species bind amino-azo-dyes with a high degree of affinity (Sorof, 1969). In our laboratory three main proteins are found, these being referred to as amino-azo-dye-binding proteins A, B and C, in order of increasing molecular weight.

The arylation that occurs involves amino acid side chains, and to date methionyl, cysteinyl and tyrosyl residues have been shown to be arylated by amino-azo-dyes *in vivo* (Miller & Miller, 1969; Ketterer & Christodoulides, 1969). Present evidence suggests that different amino acid residues are arylated in different proteins. For example amino-azo-dye-binding protein A, which has been highly purified (Ketterer *et al.*, 1967), is a protein in which methionyl residues but not cysteinyl residues are arylated (Ketterer & Christodoulides, 1969). This protein has a molecular weight of 14000 and an unusual amino acid composition, and exists as three isomers with pI 5.2, 6.0 and 7.4. The more acidic forms tend to become converted into the pI 7.4 isomer during prolonged isoelectric focusing, suggesting the loss of reversibly bound anions.

Binding of organic anions is a well-established property of amino-azo-dye-binding protein B, which has also been isolated and purified (Ketterer *et al.*, 1967; Litwack & Morey, 1970). In this protein amino-azo-dye metabolites arylate cysteinyl residues (Ketterer & Christodoulides, 1969). This protein is obviously an important component of the body. It accounts for approx. 4% of the total liver soluble protein and also occurs in the kidney, small intestine (Baldwin *et al.*, 1968; Fleischer *et al.*, 1971) and skin. It binds reversibly a diversity of organic anions, and for this reason it has recently been named ligandin (Litwack *et al.*, 1971a). For example it binds anionic metabolites of corticosteroids (Morey & Litwack, 1969) and even more strongly anionic metabolites of oestradiol (Litwack *et al.*, 1971b); it binds bilirubin and cholecystographic agents (Levi *et al.*, 1969) and also an amino-azo-dye metabolite-glutathione conjugate. It has a molecular weight of 45000 and is

composed of subunits of similar molecular weights. On prolonged isoelectric focusing it has pI approx. 9.3.

Amino-azo-dye-binding protein C, although it accounts for a large part of the bound amino-azo-dye, has yet to be purified.

An understanding of the significance of amino-azo-dye-binding proteins in carcinogenesis would be aided by a knowledge of their physiological function. At present one protein still has to be isolated and another has been isolated and characterized chemically, but nothing as yet is known of the function of either in the cell. However, some progress has been made with the third protein, ligandin. It binds many substances that either can be or have been metabolized by the endoplasmic reticulum, and it is induced by the same substances that induce the drug-metabolizing enzymes of this system (Reyes *et al.*, 1971). The function may be to transport substrates to the endoplasmic reticulum or metabolites away to other parts of the cell. If this is so its arylation by amino-azo-dye metabolites would not be surprising, nor would the reversible binding of an amino-azo-dye-glutathione conjugate.

It is important to note that ligandin shows specificity of binding. For example it binds anionic steroid hormone metabolites, but not bile acids. Further, it binds the various anionic steroid hormone metabolites with different degrees of affinity. Susceptibility to hepatocarcinogenesis in the rat is sex-dependent, and it is therefore interesting that anionic oestradiol metabolites (including a dihydroxyoestradiol-glutathione conjugate) and amino-azo-dye-glutathione conjugate are both bound by ligandin. This presents an interesting situation in which this carcinogen may compete with a steroid hormone for ligandin and then influence the course of carcinogenesis.

- Baldwin, R. W., Barker, C. R. & Moore, M. (1968) *Brit. J. Cancer* **22**, 776
- Fleischer, G., Robbins, J. & Arias, I. M. (1971) *J. Exp. Med.* in the press
- Ketterer, B. & Christodoulides, L. (1969) *Chem.-Biol. Interactions* **1**, 173
- Ketterer, B., Ross-Mansell, P. & Whitehead, J. K. (1967) *Biochem. J.* **103**, 316
- Levi, A. J., Gatmaitan, Z. & Arias, I. M. (1969) *J. Clin. Invest.* **48**, 2156
- Litwack, G. & Morey, K. S. (1970) *Biochem. Biophys. Res. Commun.* **38**, 1141
- Litwack, G., Ketterer, B. & Arias, I. M. (1971a) *Nature (London)* in the press
- Litwack, G., Morey, K. S. & Ketterer, B. (1971b) in *Drugs and Cellular Control Mechanisms* (Rabin, B. R., ed.), Macmillan, London, in the press
- Magee, P. N. & Hultin, T. (1962) *Biochem. J.* **83**, 106
- Miller, E. C. & Miller, J. A. (1969) *Ann. N.Y. Acad. Sci.* **163**, 731
- Morey, K. S. & Litwack, G. (1969) *Biochemistry* **8**, 4813

- Reyes, H., Levi, A. J., Gatmaitan, Z. & Arias, I. M. (1971) *J. Clin. Invest.* in the press
- Sorof, S. (1969) in *Jerusalem Symposia on Quantum Chemistry*, vol. 1: *Physicochemical Mechanisms of Carcinogenesis* (Bergmann, E. D. & Pullmann, B., eds.), p. 108, Israel Academy of Sciences and Arts,

The Binding of Polypeptides by Neurophysins

By D. B. HOPE (*Department of Pharmacology, University of Oxford, South Parks Road, Oxford OX1 3QT, U.K.*)

COMMUNICATIONS

A Requirement for Zinc Ions for Deoxyribonucleic Acid Synthesis by Lymphocytes Stimulated by Phytohaemagglutinin

By J. K. CHESTERS (*Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB, U.K.*)

Alford (1970) reported that the transformation of human peripheral-blood lymphocytes by phytohaemagglutinin required a bivalent cation in addition to Ca^{2+} . His results suggested that this requirement was most efficiently met by Zn^{2+} .

Lymphocytes from pig blood obtained at slaughter were used for the present investigations. The incorporation of [^3H]thymidine into the DNA of the lymphocytes was inhibited by 80–90% by the addition to the medium of 0.6mM-EDTA or 0.6mM-diethylenetriamine-*NNN'N'N'*-penta-acetate but not by 0.6mM-nitrilotriacetic acid. The addition of Zn^{2+} as ZnSO_4 reversed the inhibition by EDTA or diethylenetriamine-*NNN'N'N'*-penta-acetate. With 0.6mM-EDTA complete reversal of the inhibition required 0.05mM- Zn^{2+} ; concentrations less than this caused only partial reversal, whereas higher concentrations again decreased the incorporation. The initial pattern of reactivation with increasing concentrations of Zn^{2+} suggested a competition between EDTA and a cellular chelator that, complexed with Zn^{2+} , was directly or indirectly involved in [^3H]thymidine incorporation. The competition between chelators for the metal ions present in the system was investigated by using a computer simulation based on the program described by Perrin & Sayce (1967). The simulation suggested that the cellular chelator had to have a stability constant of the order of 10^{12} to account for the observed competition with EDTA for the activating Zn^{2+} . The validity of the simulation was tested by using this value for the stability constant of the active site to calculate the proportion of the sites that had accepted Zn^{2+} over a range of added Zn^{2+} concentrations. These proportions were found to agree well over a range of suboptimal Zn^{2+} concentrations with the experimentally determined rates of thymidine incorporation.

The time-course of the requirement for Zn^{2+} was investigated by adding EDTA at various times after stimulation of the cells with phytohaemagglutinin.

It was further studied by adding EDTA before phytohaemagglutinin and subsequently attempting to reverse the inhibition by adding an optimal amount of Zn^{2+} . At least half the maximum rate of incorporation was still obtained provided EDTA was added not earlier than 12h before the incorporation was measured. If EDTA was added initially, Zn^{2+} had to be supplied at least 17h before the incorporation was measured if at least half the maximum activity was to be obtained.

The specificity of the reversal by Zn^{2+} of the inhibition caused by EDTA will be discussed together with the implications of an apparently high affinity constant for Zn^{2+} at the active site within the cells.

Alford, R. H. (1970) *J. Immunol.* **104**, 698

Perrin, D. D. & Sayce, I. G. (1967) *Talanta* **14**, 833

Electrophoretic Slowing of Sensitized Lymphocyte-Macrophage Mixtures: a Cellular Technique in Immunochemistry

By E. A. CASPARY and E. J. FIELD (*Medical Research Council Demyelinating Diseases Unit, Newcastle General Hospital, Westgate Road, Newcastle upon Tyne NE4 6BE, U.K.*)

Immunochemical studies have generally been confined to the interaction of substances with specific antibody prepared against them. The study of specific cellular immunity, however, is more difficult and not readily made quantitative. A very specific change in immunologically sensitized cells may, however, be measured by using cell electrophoresis in the presence and in the absence of the sensitizing antigen, when the mobility of the cell will be decreased by the antigen, and a number of studies of this kind have been reported over the last 5 years (Sundaram *et al.*, 1967a,b; Hartveit *et al.*, 1968; Deingdoh & Turk, 1968). An initial study of guinea-pig peritoneal-macrophage exudates from sensitized animals showed that the antigen slowing response depended first on the presence of lymphocytes and secondly that it could be inhibited by puromycin or other inhibitors of protein synthesis (Caspary, 1971). It was further