Steroid Binding and Stabilization of Glucocorticoid-Receptor Proteins from Rat Thymus

By P. A. BELL* and A. MUNCK (*Department of Physiology, Dartmouth Medical School, Hanover, N.H. 03755, U.S.A.*)

Equilibrium and kinetic studies of glucocorticoid binding to rat thymus cells *in vitro* have demonstrated an obligatory role for binding to specific receptors before initiation of the effects of active steroids on glucose transport (Munck & Brinck-Johnsen, 1968; Munck & Wira, 1971). Nuclear and cytosol receptors can be extracted from cells with or without cortisol pretreatment by disruption by hypo-osmotic shock in 1.5 mM-MgCl₂; these receptors are at least in part proteins (Wira & Munck, 1970; Munck & Wira, 1971). We now report further studies of the cytosol receptors isolated without cortisol pretreatment of cells, using a dextran-charcoal assay specific for saturable binding of glucocorticoids to receptors.

The time-constant for dissociation of [³H]cortisol from the cytosol receptor at 37°C, 2.3 min, is similar to that characteristic of specifically bound cortisol in intact cells. At 3°C dissociation is slower, with a time-constant of 65 min. Measurement of maximum binding at a range of cortisol concentrations gives an apparent association constant of $1.7 \times 10^7 \text{ M}^{-1}$ at 3°C. Cortisol, dexamethasone and triamcinolone acetonide all compete with cortisol for receptor sites; both dexamethasone and triamcinolone acetonide are bound more strongly than cortisol. Cortisone does not compete. The time-constants for dissociation at 3°C are 30 min for cortisol, 322 min for dexamethasone and more than 12 h for triamcinolone acetonide.

Association of 0.02 μM -[³H]cortisol with the cytosol receptor at 3°C reaches a maximum by 90 min, but declines to 29% of the maximum by 8 h. With saturating concentrations of [³H]cortisol maximum binding is still retained at 8 h, and can be displaced by addition of an excess of non-radioactive cortisol. In the absence of steroid the receptor is rapidly inactivated, but can be partially stabilized by addition of 2.5 mM-EDTA.

With 0.02 μM -[³H]dexamethasone maximum binding is not attained until 3–4 h, and 66% of the maximum remains bound at 8 h; with 0.02 μM -[³H]triamcinolone acetonide 100% of maximum binding is retained at 8 h. We conclude that the steroid-receptor complex is stable, but that the receptor is rapidly inactivated in the absence of steroid, and that the apparent increased stabilization observed with dexamethasone and triamcinolone acetonide reflects their inability to respond by rapid

dissociation to disturbances of the equilibrium caused by inactivation of the free receptor.

P. A. B. thanks the Medical Research Council for leave of absence. This work was supported by U.S. Public Health Service Research Grant AM-03535.

Munck, A. & Brinck-Johnsen, T. (1968) *J. Biol. Chem.* **243**, 5556

Munck, A. & Wira, C. (1971) in *Schering Workshop on Steroid Hormone Receptors, Berlin, 1970: Advances in the Biosciences*, vol. 7, pp. 301–330, Pergamon Press-Vieweg, Oxford-Braunschweig

Wira, C. & Munck, A. (1970) *J. Biol. Chem.* **245**, 3436

Determination of Oestradiol-17 β and Oestrone in Human Plasma

By R. A. HAWKINS and R. E. OAKLEY (by invitation) (*Division of Steroid Endocrinology, Department of Chemical Pathology, School of Medicine, University of Leeds, Leeds LS2 9LN, U.K.*)

Many hormones are bound in their target tissues by specific proteins before the expression of biological activity. These proteins can be utilized as analytical tools. For example, an oestrogen-binding 'protein' from rabbit uterus has been used in the determination of oestradiol-17 β in human plasma (Korenman *et al.*, 1969; Corker & Exley, 1970).

We report here our experiences with this oestrogen-binding 'protein' for the determination of oestrone and oestradiol-17 β in human plasma.

The method comprises: (i) addition of [³H]oestrone and [³H]oestradiol-17 β to the plasma to permit determination of manipulative losses; (ii) extraction of the plasma with ether; (iii) separation of oestrone, oestradiol-17 β and other steroids by t.l.c. on silica gel; (iv) reduction of the isolated oestrone fraction to oestradiol-17 β and rechromatography; (v) the separate determination, by competitive protein-binding assay (Korenman *et al.*, 1969), of the oestradiol-17 β fraction from the plasma and that derived from oestrone by reduction.

Recovery of added [³H]oestradiol-17 β was $76.9 \pm 5\%$ (mean $\pm$ S.D., $n = 40$) and that of [³H]oestrone was $45.0 \pm 9.3\%$ ($n = 39$).

Blanks (5 ml of tap water) were $5.5 \pm 5.6 \text{ pg}$ ($n = 23$) for oestradiol-17 β and $5.4 \pm 7.1 \text{ pg}$ ($n = 17$) for oestrone.

Precision was determined on a pool of male plasma and on samples of female plasma. The coefficient of variation (Snedecor, 1952) at the male plasma concentration was 29% for oestradiol-17 β ($n = 18$) and 30% for oestrone ($n = 13$). At concentrations above 5 ng/100 ml of plasma the coefficient of variation was 13% for oestradiol-17 β ($n = 18$) and 16% for oestrone ($n = 7$).

* Present address: Dunn Nutritional Laboratory, Milton Road, Cambridge CB4 1XJ, U.K.

Accuracy was assessed by adding known quantities of each oestrogen (50–2500 pg) separately to 2.5 ml samples taken from a pool of male plasma and determining the masses of each oestrogen present. The equation of the regression line for the recovery of oestradiol-17 β was $y = 1.03x + 24$ (endogenous concentration, determined separately, 33 pg/2.5 ml) and for the recovery of oestrone was $y = 0.99x + 79$ (endogenous concentration 74 pg/2.5 ml). No interference by one oestrogen in the determination of the other was detected.

A high degree of specificity was achieved by a combination of the insensitivity of the cytosol binding 'protein' to non-oestrogenic steroids (Korenman, 1969; Corker & Exley, 1970; and our observations) and selective chromatography.

Application of the method to the determination of oestrone and oestradiol-17 β throughout the menstrual cycle will be presented.

We thank the Medical Research Council for financial support.

Corker, C. S. & Exley, D. (1970) *Steroids* **15**, 469

Korenman, S. G. (1969) *Steroids* **13**, 163

Korenman, S. G., Perrin, L. E. & McCallum, T. P. (1969) *J. Clin. Endocrinol.* **29**, 879

Snedecor, G. (1952) *Biometrics* **8**, 85

The Intracellular Distribution of Adrenal 3 β -Hydroxy Steroid Dehydrogenase and 3-Oxo Steroid Δ^4 – Δ^5 -Isomerase Studied with the Zonal Rotor

By R. A. COWAN, J. K. GRANT, C. A. GILES and W. BIDDLECOMBE (*University Department of Steroid Biochemistry, Royal Infirmary, Glasgow C.4, and Department of Physiology, University of Glasgow, Glasgow W.2, U.K.*)

The conversion of pregnenolone into progesterone during steroid biosynthesis in the adrenal cortex requires two enzymes, an NAD⁺-dependent 3 β -hydroxy steroid dehydrogenase and a 3-oxo steroid Δ^4 – Δ^5 -isomerase. It is well established that these enzymes are located in the microsomal fraction obtained by differential centrifugation after tissue homogenization (Beyer & Samuels, 1956). The observed association of dehydrogenase and isomerase activities with the mitochondrial fraction has been attributed to contamination by microsomal enzymes (for discussion see Hayano *et al.* 1956). This view has recently been questioned and the possibility of a bimodal distribution of these enzymes in the adrenal cell has been suggested (McCune *et al.*, 1970). Experiments involving rate-zonal centrifugation of a crude mitochondrial preparation on sucrose gradients in the HS zonal rotor (Measuring and

Scientific Equipment Ltd.) have now indicated that the dehydrogenase and isomerase are firmly bound to the mitochondria, as well as to the endoplasmic reticulum, of the bovine adrenal cortex.

The crude mitochondrial fraction prepared from 5g of bovine adrenal cortex by conventional differential centrifugation was loaded on to a continuous sucrose density gradient (10–55%, w/v) linear with rotor radius. Separation was allowed to proceed for 45 min at 7500g, after which 15 ml fractions were collected and centrifuged at 24000g for 15 min. The pellets were assayed for marker enzymes as follows: for mitochondria, cytochrome oxidase, succinate dehydrogenase, steroid 11 β -hydroxylase and cholesterol side-chain-cleavage enzyme system; for microsomal fraction, 5'-nucleotidase and steroid-21-hydroxylase; for lysosomes, acid phosphatase. In addition, the protein and cholesterol contents of each fraction were determined. Certain fractions were also examined by electron microscopy.

Biochemical and electron-microscopic examination of the fractions show the microsomal particles to be concentrated at the light end of the gradient, whereas the mitochondria sediment as a broad band through the gradient. The width of this band reflects considerable variation of mitochondrial size.

A comparison of the distribution of dehydrogenase and isomerase enzymes with that of the marker enzymes revealed that the enzymes responsible for the conversion of pregnenolone into progesterone are firmly associated with mitochondria and microsomal fraction. The significance of this observation in relation to pathways of steroid biosynthesis in the adrenal cortex will be discussed, together with the advantages of the zonal rotor for preparing mitochondria with minimal microsomal contamination.

R. A. C. acknowledges the receipt of a Scholarship for Training in Research Methods from the Medical Research Council.

Beyer, K. F. & Samuels, L. T. (1956) *J. Biol. Chem.* **219**, 69

Hayano, M., Saba, N., Dorfman, R. I. & Hechter, O. (1956) *Recent Progr. Horm. Res.* **12**, 79

McCune, R. W., Roberts, S. & Young, P. L. (1970) *J. Biol. Chem.* **245**, 3859

Phosphorylation of the Erythrocyte Membrane: Stimulation by Adenosine 3':5'-Cyclic Monophosphate

By M. J. DUFFY and V. SCHWARZ (*Department of Medical Biochemistry, University of Manchester, Manchester M13 9PL, U.K.*)

Cyclic AMP (adenosine 3':5'-cyclic monophosphate) is thought to be implicated in the control of cation permeability of membranes. It is possible