ions are believed to lead to substitution at position 8 (Lowe & Dean, 1971; Acheson, 1960; Weibel *et al.*, 1971). On the other hand the vicinal hydroxyl groups of the ribose rings (of NAD) are susceptible to periodate oxidation (Gilham, 1971) and to the titanous chloride method of Novais (1971). Further substitution of the pyrophosphate bridge (to give phosphonates or phosphate esters) would appear to be difficult to carry out selectively; this part of the NAD molecule may well be involved when NAD is coupled to polymers by using carbodi-imide-type reagents (H. G. Khorana, personal communication). Others have speculated that these reactions may involve the ribose hydroxyl groups (Larsson & Mosbach, 1971), and the amine of the adenine ring cannot, in our view, be entirely ruled out either. Reaction of the 2'-phosphate of NADP with carbodi-imide-type reagents on the other hand gives a more definite product with cellulose phosphate. Reactions such as the enzymic coupling of insolubilized nicotinamide analogues to ADP-ribose (Dixon & Webb, 1964; Beilmann, 1971) have yet to be successfully applied to the insolubilization of NAD(P).

The lack of reactivity of the nicotinamide nucleotides emphasises the fact that many of the above coupling reagents introduce an undesirable capacity for ion-exchange in the resulting polymer. Methods are required for distinguishing between specific and non-specific interactions between insolubilized cofactor and enzyme. Thus displacements of enzymes from columns with either unbound cofactor or substrate or both should at first sight be relatively insensitive to high salt concentrations. However, enzymes seem to possess non-catalytic cofactor-binding sites whose dissociation constants are only 5 times those of the catalytic sites (Weiner, 1969). Another complicating feature is shown in the following experiment. Glucose 6-phosphate dehydrogenase can be bound to a column of NADP-cellulose at low ionic strength. Additions of NADP plus substrate cause a steady elution of NADPH from the column. Increasing the salt concentration results in an eluate whose NADPH concentration increases with time, indicating that the enzyme is catalytically active when bound to the cofactor column. This interaction is salt-sensitive, and the enzyme can thus be displaced solely by salt-dependent effects.

We have shown that the CNBr activation technique introduces groups that can show marked ion-exchange properties, and, further, that the strength of the latter is dependent on the quantity of CNBr used in the activation process (C. R. Lowe & P. D. G. Dean, unpublished work). These groups arise by degradation of the activated complex, a process that may be accentuated by the poor reactivity of adenine nucleotides. This problem does not arise with more reactive ligands, since it can be completely

eliminated by treatment with 3-aminopropane-1,2-diol or D-glucosamine after sufficient time has been allowed for some nucleotide to react.

Further side reactions occur when nicotinamide nucleotides are coupled to extended 'spacers' by a water-soluble carbodi-imide reaction (Lowe & Dean, 1971); the properties of the resulting polymer are, under many conditions, reminiscent of those of the original CNBr-activated polymer.

We believe that the most satisfactory way of analysing these complications is to investigate substrate and cofactor displacements under wide ranges of salt concentrations, especially under the conditions of salt gradients.

NAD columns have been compared in properties with the following insolubilized fragments of the NAD molecule: ADP-ribose, ADP, AMP, adenosine and adenine. The binding of various dehydrogenases has been studied. A gradation of affinities has been observed that are characteristic for some groups of dehydrogenases.

Some more recent results from our investigations of the properties of these polymers will be discussed.

This work was supported in part by the Science Research Council.

- Acheson, R. M. (1960) *An Introduction to the Chemistry of Heterocyclic Compounds*, pp. 307-308, Interscience, New York
- Axén, R., Ernäck, J. & Porath, S. (1967) *Nature (London)* **215**, 491
- Beilmann, J. F. (1971) *Bull. Soc. Chim. Biol.* **50**, 151
- Dixon, M. & Webb, E. C. (1964) *Enzymes*, 2nd edn., p. 370, Longmans Green, London
- Gilham, P. T. (1971) *Methods Enzymol.* **21**, 191
- Larsson, P.-O. & Mosbach, K. (1971) *Biotechnol. Bioeng.* **13**, 393
- Lowe, C. R. & Dean, P. D. G. (1971) *FEBS Lett.* **14**, 313
- Novais, F. M. (1971) Ph.D. Thesis, University of Birmingham
- Porath, J. & Fornstedt, N. (1970) *J. Chromatogr.* **51**, 479
- Weibel, M. K., Weetall, H. H. & Bright, H. J. (1971) *Biochem. Biophys. Res. Commun.* **44**, 347
- Weiner, H. (1969) *Biochemistry* **8**, 552

General Ligands and (Co)Substrate Elution in Affinity Chromatography

By KLAUS MOSBACH, HUGH GUILFORD, ROLF OHLSSON and MARGARETHA SCOTT (*Biochemical Division, Chemical Centre, University of Lund, P.O. Box 740, S-220 07 Lund, Sweden*)

A hallmark of enzyme purification by affinity chromatography or biospecific absorption has been the high degree of selectivity of this technique, since it has been possible to bind a specific inhibitor for

precisely the protein under scrutiny. This very specificity, however, clearly limits the scope of applicability of many immobilized purification systems, the synthesis of which often involves elaborate chemistry. We in Lund have therefore focused our attention on the development of generalized affinity chromatography by using matrix-bound ligands with a broad 'enzyme spectrum' (Mosbach *et al.*, 1971).

The ligands we selected to bind to separate gel matrices were NAD^+ and an AMP analogue. For preparation of immobilized NAD^+ , 6-aminohexanoic acid has been coupled to Sepharose 4B by the CNBr method (Axén *et al.*, 1967), and subsequently dicyclohexylcarbodi-imide was used to attach NAD^+ , probably via ester linkages (Larsson & Mosbach, 1971). These preparations have been shown to retain some of the cozymic function of the free nicotinamide nucleotide (Mosbach, 1971; Larsson & Mosbach, 1971). The use of similarly immobilized cofactors in affinity chromatography has also been reported by other workers in this field (Lowe & Dean, 1971).

For preparation of the immobilized AMP analogue, N^6 -(6-aminoethyl)-AMP (H. Guilford, P.-O. Larsson & K. Mosbach, unpublished work) has been linked directly to activated Sepharose 4B. In both cases the 'spacing' of the cofactor and inhibitor moieties from the matrix by a hexacarbon 'chain' of length about 0.7 nm ($\text{\AA}$) was used to increase the steric availability of the ligands to the enzymes studied, e.g. glyceraldehyde 3-phosphate dehydrogenase (EC 1.2.1.12) and lactate dehydrogenase (EC 1.1.1.27).

On application of a mixture of bovine serum albumin, glyceraldehyde 3-phosphate dehydrogenase and lactate dehydrogenase to columns packed with either gel matrix, only the two enzymes were retained. A 0.15 mM solution of NAD^+ eluted only glyceraldehyde 3-phosphate dehydrogenase, and subsequent application of 0.15 mM-NADH eluted only lactate dehydrogenase. The observed elution pattern is consistent with the affinities of NAD^+ and NADH reported for the two dehydrogenases (McPherson, 1970, and references cited therein). Subsequent more detailed studies with the AMP-Sepharose preparation gave the following results.

(1) The enzymes could not be eluted by a 0.0–0.5 M-KCl gradient, indicating strong binding of the enzyme to the nucleotide.

(2) Elution of glyceraldehyde 3-phosphate dehydrogenase followed by lactate dehydrogenase was achieved by application of a gradient of 0.0–0.15 M-salicylate, which is known to be an inhibitor competing with NAD^+ for both enzymes (Hines & Smith, 1964).

(3) Batchwise elution studies of bound lactate dehydrogenase under specified conditions revealed the following approximate relative elution efficiencies

with different eluents: lactate+semicarbazide, 0; 1 mM- NAD^+ , 0; lactate+ NAD^+ , 17; lactate+semicarbazide+ NAD^+ , 21; NADH, 21. We ascribe the closely comparable efficiency of NADH and of lactate+semicarbazide+ NAD^+ to the enzymic formation of reduced cofactor taking place in the latter system under the experimental conditions. The rather high efficiency of lactate+ NAD^+ is surprising in view of the fact that the equilibrium is far to the side of the oxidized form of the cofactor. The cumulative effect in enzyme elution due to the presence of the substrate, lactate, will be discussed.

(4) Commercially available crude pyruvate kinase having no affinity for AMP (Reynard *et al.*, 1961), but highly contaminated with lactate dehydrogenase activity, was completely separated from the dehydrogenase in a single-step procedure: all the pyruvate kinase appeared in the early fractions, leaving the lactate dehydrogenase bound.

The general applicability to affinity chromatography of the bound nucleotide preparations described will be discussed together with aspects of selective (co)substrate elution.

Axén, R., Porath, J. & Ernback, S. (1967) *Nature (London)* **214**, 1302

Hines, W. J. W. & Smith, M. J. H. (1964) *Nature (London)* **201**, 192

Larsson, P.-O. & Mosbach, K. (1971) *Biotechnol. Bioeng.* **13**, 393

Lowe, C. R. & Dean, P. D. G. (1971) *FEBS Lett.* **14**, 313

McPherson, A., Jr. (1970) *J. Mol. Biol.* **51**, 39

Mosbach, K. (1971) *Sci. Amer.* **224**, 26

Mosbach, K., Guilford, H., Larsson, P.-O., Ohlsson, R. & Scott, M. (1971) *Biochem. J.* **125**, 20p

Reynard, A. M., Hass, L. F., Jacobsen, D. D. & Boyer, P. D. (1961) *J. Biol. Chem.* **236**, 2227

Molecular Properties of Immobilized Proteins

By D. GABEL and J. PORATH (*Institute of Biochemistry, University of Uppsala, Uppsala, Sweden*)

A method is presented that allows us to follow conformational changes in immobilized proteins (Gabel *et al.*, 1972). The method is based on the fact that a gel bed of an immobilized protein can be considered as a fluorescent surface. A special cell has been devised that fits into most commercial and home-made spectrofluorometers. As it can be operated as a small chromatographic column, it allows rapid dialysis and change of liquid.

It was established that gels of Sephadex and agarose represent an environment for tryptophan similar to that of water. Changes in the observed fluorescent properties of proteins can thus be attributed to changes in their conformation. The inherent limitations of the method are discussed.