studies indicated that, as with the related enzyme 2-hydroxybutyrate dehydrogenase, both the polar choline head group and the hydrocarbon chains are involved in the binding of the lipid to the protein.

These specific lipid-protein complexes are probably examples of allotropic interactions. This form of complexing may explain aspects of the behaviour of these enzymes *in vivo*.

Dodd, G. H. & Radda, G. K. (1969) *Biochem. J.* **114**, 409
Racker, E. (1967) *Fed. Proc. Fed. Amer. Soc. Exp. Biol.* **26**, 1335

Biological Activity of Double-Stranded Ribonucleic Acid Fragments

By V. G. EDY and MARIA SZEKELY (*Department of Biochemistry, Imperial College of Science and Technology, London SW7 2AY, U.K.*)

Fungal viruses containing double-stranded RNA have been detected in a number of *Penicillium* species, and all of them tested as yet have been found to possess the capacity to induce interferon in animals and to give protection against infection by different viruses (for review see Colby, 1971). The double-stranded RNA responsible for these effects has been studied with respect to the importance of molecular weight for biological activity.

Double-stranded RNA isolated from *Penicillium chrysogenum* virus (Lemke & Ness, 1970; Buck *et al.*, 1971) has been fragmented in three different ways: (a) by digestion with ribonuclease III; (b) by digestion with high concentrations of pancreatic ribonuclease A in high salt concentration; (c) by ultrasonic disintegration.

Degradation was followed by viscosity measurements, polyacrylamide-gel electrophoresis and sucrose-density-gradient analysis. With the exception of some special mild conditions, breakdown was always random, producing a heterogeneous mixture of fragments of a rather wide size range. Optimal conditions for obtaining relatively large fragments with no traces of intact material have been worked out. The 'melting' temperature, T_m , was practically unchanged after degradation, suggesting that the breakdown products were still double-stranded.

The capacity of the RNA to induce interferon was diminished even after mild degradation, though at the same time the toxicity increased as compared with that of the intact RNA. (These changes were observed in a virus-sensitive strain of mice, BSVS.) As similar results were obtained whichever method of degradation was used, but the results varied with the degree of degradation, it would seem that biological activity is dependent on molecular weight rather than on the intact structure of any particular site in the RNA molecule.

We thank Dr. F. Himmelweit for performing the assays of interferon-inducing capacity. This work forms part of a collaborative study on antiviral agents with Beecham Research Laboratories. V. G. E. thanks the Medical Research Council for a Research Studentship.

Buck, K. W., Chain, E. B. & Himmelweit, F. (1971) *J. Gen. Virol.* **12**, 131
Colby, C. (1971) *Progr. Nucleic Acid Res. Mol. Biol.* **11**, 1
Lemke, P. A. & Ness, T. M. (1970) *J. Virol.* **6**, 813

Resolution of Supercoiled Deoxyribonucleic Acid Structures by Light-Scattering

By A. M. CAMPBELL and D. J. JOLLY (*Institute of Biochemistry, University of Glasgow, Glasgow W.2, U.K.*)

There is, at present, considerable interest in and uncertainty about both the exact tertiary structure of supercoiled DNA and the possible transitions occurring on alteration of the secondary structure by, for example, dye titration (Bauer & Vinograd, 1968; Wang, 1969; Revet *et al.*, 1971; Dean & Lebowitz, 1971). Previous techniques used have been mainly hydrodynamic, requiring small amounts of material, but providing rather ambiguous answers to the problem of tertiary-structure determination.

Light-scattering requires larger amounts of material, but supplies a direct numerical parameter of tertiary structure, the root-mean-square radius, and also a complete simple interference curve on which further calculations can be made.

We have undertaken a light-scattering study of bacteriophage ϕ X174-RFI DNA in attempts to differentiate between different possible structures, such as (a) the straight interwound model, (b) the toroidal model (Gray, 1967) and (c) the interwound Y-shaped model as seen in electron micrographs.

The experiments were performed on pure bacteriophage ϕ X174-RFI DNA in BPES buffer (Krasna & Harpst, 1964), pH 6.8, at 25°C by using a refractive-index increment of 0.166 ml/g (Krasna, 1970).

Results were as follows. The number of superturns/molecule was -11.8 ± 1.6 ; in further calculations it was assumed to be an integer, 12. The molecular weight of bacteriophage ϕ X174-RFI DNA was $3.17 \times 10^6 \pm 0.19 \times 10^6$, significantly lower than the value of twice 1.7×10^6 previously reported for single-stranded bacteriophage ϕ X174 DNA (Sinsheimer, 1959). The root-mean-square radius was 103.5 ± 12 nm.

To extract further information from the scattering experiment, we predicted reciprocal particle scattering factors, $P(\theta)^{-1}$, for each of the models mentioned above, having dictated their dimensions with the experimental root-mean-square radius and molecular