of mitochondrial oxidative phosphorylation by bongkreikic acid was measured. Again, the presence of CoA considerably shortened the time required for bongkreikic acid to act.

A possible explanation for these observations is that bongkreikic acid must first be converted into a CoA derivative before it can inhibit adenine nucleotide translocation. Further work is aimed towards the isolation of such a derivative and identification of an enzyme (probably an acyl-CoA synthetase) responsible for its formation.

We thank Mrs. Cheryll Crosby for her excellent technical assistance.

Henderson, P. J. F. & Lardy, H. A. (1970) *J. Biol. Chem.* **245**, 1319

Henderson, P. J. F., Lardy, H. A. & Dorschner, E. (1970) *Biochemistry* **9**, 3453

Kemp, A., Out, T. A., Guiot, H. F. L. & Souverijn, J. H. M. (1970) *Biochim. Biophys. Acta* **223**, 460

Klingenberg, M., Grebe, K. & Heldt, H. W. (1970) *Biochem. Biophys. Res. Commun.* **39**, 363

Shug, A., Lerner, E., Elson, C. & Shrago, E. (1971) *Biochem. Biophys. Res. Commun.* **43**, 557

Ureolysis in the Ovine Rumen

By A. R. COOK (Department of Microbiology, Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB, U.K.)

The rumen contents of sheep and cattle on all diets possess ureolytic activity. Although this activity was shown to be of bacterial intracellular origin, so confirming earlier work (Gibbons & McCarthy, 1957), only a few ureolytic bacteria have so far been isolated and positively identified (Gibbons & Doetsch, 1959). In the present work only ten ureolytic strains out of over 1000 bacteria were isolated from a sheep fed on hay and dried grass.

The bacteria were isolated by two methods. In the first, 15 different media varying from the non-selective counting media to several selective media were used in the standard roll-tube method. The media should provide growth conditions for all types of rumen bacteria, but the only ureolytic bacteria isolated were five strains of *Staphylococcus* (group II), two strains of group VI and a strain of *Lactobacillus casei* var. *casei*. These were isolated from the 10^6 -dilution cultures.

In the second method, media that enriched the ureolytic bacteria of undiluted rumen fluid were used to enrich serial dilutions of rumen fluid, and *Klebsiella aerogenes* and a *Streptococcus* sp. (intermediate in properties between *Streptococcus faecium* and *Streptococcus faecalis*) were isolated from dilutions 10^5 and 10^7 respectively.

No ureolytic strains of obligate anaerobes were isolated from the rumen, and the results of this survey suggest that ureolytic activity is associated with bacteria present in comparatively small numbers and not with those bacteria present in high numbers and considered important in rumen function.

The *Streptococcus* sp. was ureolytic only under anaerobic conditions, and investigations of factors controlling the level of the activity per unit growth show that the activity would account for most of the ureolytic activity of the sheep rumen contents.

Gibbons, R. J. & Doetsch, R. N. (1959) *J. Bacteriol.* **77**, 417-428

Gibbons, R. J. & McCarthy, R. D. (1957) *Misc. Publ. Univ. Md. Agr. Exp. Sta.* **291**, 12-16

Factors Affecting the Inhibition of Mitochondrial Adenosine Triphosphatase by Trialkyltin Compounds

By M. J. SELWYN, S. J. DUNNETT, R. D. PHILO and A. P. DAWSON (School of Biological Sciences, University of East Anglia, Norwich NOR 88C, U.K.)

The 2,4-dinitrophenol-stimulated ATPase* of rat liver mitochondria is potently inhibited by trialkyltin compounds (Aldridge, 1958; Aldridge & Street, 1971). However, unless Mg^{2+} ions are present, at concentrations greater than or equal to the concentration of ATP, the apparent K_i for inhibition by trimethyltin is very much increased.

At pH 6.8 the effectiveness of trimethyltin as an inhibitor of the 2,4-dinitrophenol-stimulated ATPase of whole mitochondria is diminished by the presence of 10 mM- P_i . In whole mitochondria the situation is complicated by intramitochondrial pH changes resulting from the Cl^- -OH $^-$ exchange mediated by trialkyltins (Selwyn *et al.*, 1970). At pH 6.8 the effectiveness of trimethyltin as an inhibitor of the Mg^{2+} -stimulated ATPase of fragmented mitochondria is also diminished by the presence of P_i , but at pH 7.5 the effect of P_i is much less. Rose (1969) has reported that trialkyltins are complexed by P_i and that the affinity decreases above pH 7. In the absence of P_i the K_i for inhibition of the ATPase of broken mitochondria by trimethyltin is about one-quarter of the dissociation constant measured directly by Aldridge & Street (1970). However, Aldridge & Street (1970) measured the binding at pH 6.8 in the presence of 15 mM- P_i . When allowance is made for the effects of P_i there is little discrepancy between the two estimates of the dissociation constant.

The failure of atractyloside to inhibit the ATPase of broken mitochondria supports the view (Stockdale

* Abbreviation: ATPase, adenosine triphosphatase.