viscosity of which can be accounted for in terms of components A and B, two transitions in viscosity were observed as KCl concentration was increased. Increasing the salt concentration to 50mM-KCl resulted in a 50% decrease in viscosity, attributed to the polyelectrolyte charge-shielding effect. The second transition, indicated by a 25% decrease in viscosity between 0.5M- and 1.5M-KCl, was examined further. Although there was no variation in the molecular weight, nor any interconversion of mucoprotein components A and B, there was a marked change in the hydrodynamic properties of the mucoprotein. The s^0 values of mucoprotein components A and B decrease from $s_{25,w}^0$ 18.7 and 4.9S to 16.1 and 3.9S respectively when the concentration of KCl was raised from 0.2M to 2.5M. Values of intrinsic viscosity, of K_s (Creeth & Knight, 1965) and of the frictional ratios show that the mucoproteins A and B contracted from expanded hydrated spheres in 0.2M-KCl to asymmetric compact equivalent hydrodynamic ellipsoids in 2.5M-KCl. This transition is only partially reversible, and indicates the occurrence of definite structural changes within the mucoprotein.

The viscosity of component A was highly shear-dependent in 0.2M-KCl, this shear-dependence being almost absent in 2.5M-KCl. Further, at zero shear in 0.2M-KCl the intrinsic viscosity of component A showed an anomalously high increase in viscosity with concentration, reflecting the ability of the mucus to form a gel under physiological conditions. The absence of this anomalous increase in viscosity in 2.5M-KCl indicates that the expanded hydrated structure, as well as the highly polymerized form of the mucoprotein (component A), is necessary for gel formation. The concentration-dependence of shear suggests that gel formation does not involve strongly co-operative interactions (Rees, 1969).

The authors thank the Medical Research Council for financial support.

Creeth, J. H. & Knight, C. G. (1965) *Biochem. J.* **105**, 1135

Rees, D. (1969) *Advan. Carbohydr. Chem.* **24**, 267

Snary, D. & Allen, A. (1971) *Biochem. J.* **123**, 845

Snary, D., Allen, A. & Pain, R. H. (1970) *Biochem. Biophys. Res. Commun.* **40**, 844

Mechanisms of Colicinogeny

By K. G. HARDY and G. G. MEYNELL (*Guinness-Lister Research Unit, Lister Institute, Chelsea Bridge Road, London SW1W 8RH, U.K.*)

Nine colicin factors were examined in *Escherichia coli* K 12 with respect to the effect on colicin synthesis of the hosts *recA* (recombination-deficiency) allele and the relative amount of each colicin that is free or cell-bound. The results can be summarized as

follows: group 1 (colicin factors E1a-16, E1-K30, E2-P9 and K-235), *recA*⁺/*recA* ratio of bound colicin ≥ 16 and bound/free colicin ratio in *recA*⁺ host ≤ 1 ; group 2 (colicin factors Ia-CT4, Ib-P9, V-K30 and V-K94), *recA*⁺/*recA* ratio of bound colicin ≤ 2 and bound/free colicin ratio in *recA*⁺ host > 8 ; group 3 (colicin factor B-K98), *recA*⁺/*recA* ratio of bound colicin ≥ 16 and bound/free colicin in *recA*⁺ host > 8 .

Colicin synthesis may result from de-repression of the colicin operon in group 1, from conversion (i.e. constitutive synthesis) in group 2, and from conversion accompanied by a high rate of vegetative plasmid replication in group 3. A hypothetical group 4, with *recA*⁺/*recA* ratio of bound colicin ≤ 2 and bound/free colicin ratio in *recA*⁺ host, may be impossible.

Relationships between the Uptake of Uracil and Amino Acids by *Candida utilis*

By R. W. JONES and D. G. WILD (*Microbiology Unit, Department of Biochemistry, University of Oxford, Oxford OX1 3QU, U.K.*)

Candida utilis was grown at 30°C in a minimal medium with ammonia as the nitrogen source. When leucine was added to such a culture, there was no change in either the rate of protein synthesis, as measured by the incorporation of radioactive sulphate, or growth rate. However, when the medium also contained radioactive uracil, the rate of incorporation of this latter into RNA increased by about 50%. Chemical analyses showed that the addition of leucine had no effect on the protein and RNA contents of the cells. Instead, the contribution of exogenous uracil to the pyrimidine nucleotide pools increased from about 40 to 60% of the total, so that the RNA made after the amino acid was added was of greater specific radioactivity.

These findings suggested that leucine increases the rate of uptake of uracil by *C. utilis*. This was confirmed directly. The maximum velocity of uracil uptake was almost doubled in yeast grown with leucine present, whereas the Michaelis constant was unaltered. Experiments with cycloheximide showed that this apparent increase in the number of sites available for uracil uptake depended on protein synthesis. The kinetics of the appearance of the increased uptake of uracil, and also the decay of this activity when leucine was removed from cultures, support the interpretation that leucine acts through an inductive process.

These actions of leucine are restricted neither to this amino acid nor to the incorporation of uracil. Growth in the presence of leucine also stimulated the uptake of uridine, cytosine and adenine as well

as that of amino acids, including leucine and lysine. The effects of phenylalanine and aspartic acid were similar to those of leucine. With lysine in the medium, the capacity of the cells to take up uracil, leucine, phenylalanine and aspartic acid was unaffected, but the uptake of lysine itself was decreased. Similarly, growth with uracil led to a diminished capacity to take up this pyrimidine, but had no effect on the incorporation of the amino acids mentioned above.

Thus in *C. utilis* there is a complex series of relationships between mechanisms for the uptake of purines, pyrimidines and amino acids in which induction and repression seem to be involved.

Altered Purine Permease Activity in a Purine-Analogue-Resistant Mutant of *Saccharomyces cerevisiae*

By W. R. PICKERING and R. A. WOODS (*Department of Genetics, University of Sheffield, Sheffield S10 2TN, U.K.*)

Mutants of yeast selected for resistance to the purine analogue 4-aminopyrazolo[3,4-*d*]pyrimidine have been allocated to seven genes, *app1* to *app7* (Pickering & Woods, 1972). The mutant *app1-14* shows lower uptake of adenine and hypoxanthine and smaller purine pools than does the wild-type. Guanine uptake and pool size are unchanged in the mutant.

The wild-type is able to take up ^{14}C -labelled 4-aminopyrazolo[3,4-*d*]pyrimidine from the medium and to incorporate it into cold trichloroacetic acid-insoluble material, but there is no detectable uptake by mutant *app1-14*. Uptake of adenine and hypoxanthine obey simple enzyme kinetics in both strains. The K_m values in the wild-type are K_m (adenine) $1.2 \times 10^{-5} \text{ M}$ and K_m (hypoxanthine) $2.2 \times 10^{-5} \text{ M}$, and for mutant *app1-14* K_m (adenine) $2.6 \times 10^{-6} \text{ M}$ and K_m (hypoxanthine) $2.9 \times 10^{-5} \text{ M}$. Uptake of adenine and hypoxanthine is competitively inhibited by purines and purine analogues in wild-type, but not in the mutant. Uptake of guanine in both strains is inhibited by hypoxanthine, but not by adenine. Purine phosphoribosyltransferase activity with adenine, hypoxanthine, guanine and the purine analogue as substrates is 50–60% higher in mutant *app1-14* than in the wild-type.

These results suggest that the resistance of mutant *app1-14* to 4-aminopyrazolo[3,4-*d*]pyrimidine and other purine analogues is due to increased specificity of a purine permease that is primarily concerned with the uptake of adenine.

Pickering, W. R. & Woods, R. A. (1972) *Heredity* in the press

Specificity of the Proline Permease of *Escherichia coli*

By I. ROWLAND and H. TRISTRAM (*Department of Botany and Microbiology, University College London, London W.C.1, U.K.*)

An earlier study of the specificity of the proline permease of *Escherichia coli* (Tristram & Neale, 1968) has been extended. In general, the presence of both a carboxyl group at position 2 of the pyrrolidine ring and a secondary amine group were essential for uptake of a compound by the proline permease. Compounds in which replacement of the pyrrolidine ring by other five-membered rings, e.g. the thiazolidine ring, usually possessed high affinity for the permease. Azetidine-2-carboxylic acid, in which the ring is smaller than that of proline, was a substrate, whereas pipecolic acid, the higher homologue of proline, was completely inactive. On the other hand *cis*- and *trans*-3,4-methano-L-prolines, also six-membered ring structures, were both substrates. The difference between these latter compounds and pipecolic acid may reside in the fact that the methanoproline exist in the 'boat' configuration whereas pipecolic acid possesses the 'chair' configuration.

Substituted prolines with *cis* configuration possessed a higher affinity for the permease than did the corresponding *trans* isomers. Effectiveness of substituted prolines as substrates was influenced by the position, size and the polar or non-polar nature of the substituent group.

At 0°C proline accumulation is prevented, but a stoichiometric exchange between intracellular and extracellular proline occurs (Britten & McClure, 1962). Under these conditions many proline analogues with affinity for the permease exchanged with previously accumulated L-[^{14}C]proline. However, some analogues, notably *trans*-3-methyl- and *trans*-4-methyl-L-proline, though possessing high affinity for the permease, showed an almost total inability to exchange with intracellular proline. These observations substantiate previous suggestions that proline uptake and exchange may not be closely interrelated (Britten & McClure, 1962; Tristram & Neale, 1968).

Britten, R. J. & McClure, F. T. (1962) *Bacteriol. Rev.* **26**, 292

Tristram, H. & Neale, S. (1968) *J. Gen. Microbiol.* **50**, 121