as that of amino acids, including leucine and lysine. The effects of phenylalanine and aspartic acid were similar to those of leucine. With lysine in the medium, the capacity of the cells to take up uracil, leucine, phenylalanine and aspartic acid was unaffected, but the uptake of lysine itself was decreased. Similarly, growth with uracil led to a diminished capacity to take up this pyrimidine, but had no effect on the incorporation of the amino acids mentioned above.

Thus in *C. utilis* there is a complex series of relationships between mechanisms for the uptake of purines, pyrimidines and amino acids in which induction and repression seem to be involved.

Altered Purine Permease Activity in a Purine-Analogue-Resistant Mutant of *Saccharomyces cerevisiae*

By W. R. PICKERING and R. A. WOODS (*Department of Genetics, University of Sheffield, Sheffield S10 2TN, U.K.*)

Mutants of yeast selected for resistance to the purine analogue 4-aminopyrazolo[3,4-*d*]pyrimidine have been allocated to seven genes, *app1* to *app7* (Pickering & Woods, 1972). The mutant *app1-14* shows lower uptake of adenine and hypoxanthine and smaller purine pools than does the wild-type. Guanine uptake and pool size are unchanged in the mutant.

The wild-type is able to take up ^{14}C -labelled 4-aminopyrazolo[3,4-*d*]pyrimidine from the medium and to incorporate it into cold trichloroacetic acid-insoluble material, but there is no detectable uptake by mutant *app1-14*. Uptake of adenine and hypoxanthine obey simple enzyme kinetics in both strains. The K_m values in the wild-type are K_m (adenine) $1.2 \times 10^{-5} \text{ M}$ and K_m (hypoxanthine) $2.2 \times 10^{-5} \text{ M}$, and for mutant *app1-14* K_m (adenine) $2.6 \times 10^{-6} \text{ M}$ and K_m (hypoxanthine) $2.9 \times 10^{-5} \text{ M}$. Uptake of adenine and hypoxanthine is competitively inhibited by purines and purine analogues in wild-type, but not in the mutant. Uptake of guanine in both strains is inhibited by hypoxanthine, but not by adenine. Purine phosphoribosyltransferase activity with adenine, hypoxanthine, guanine and the purine analogue as substrates is 50–60% higher in mutant *app1-14* than in the wild-type.

These results suggest that the resistance of mutant *app1-14* to 4-aminopyrazolo[3,4-*d*]pyrimidine and other purine analogues is due to increased specificity of a purine permease that is primarily concerned with the uptake of adenine.

Pickering, W. R. & Woods, R. A. (1972) *Heredity* in the press

Specificity of the Proline Permease of *Escherichia coli*

By I. ROWLAND and H. TRISTRAM (*Department of Botany and Microbiology, University College London, London W.C.1, U.K.*)

An earlier study of the specificity of the proline permease of *Escherichia coli* (Tristram & Neale, 1968) has been extended. In general, the presence of both a carboxyl group at position 2 of the pyrrolidine ring and a secondary amine group were essential for uptake of a compound by the proline permease. Compounds in which replacement of the pyrrolidine ring by other five-membered rings, e.g. the thiazolidine ring, usually possessed high affinity for the permease. Azetidine-2-carboxylic acid, in which the ring is smaller than that of proline, was a substrate, whereas pipecolic acid, the higher homologue of proline, was completely inactive. On the other hand *cis*- and *trans*-3,4-methano-L-prolines, also six-membered ring structures, were both substrates. The difference between these latter compounds and pipecolic acid may reside in the fact that the methanoproline exist in the 'boat' configuration whereas pipecolic acid possesses the 'chair' configuration.

Substituted prolines with *cis* configuration possessed a higher affinity for the permease than did the corresponding *trans* isomers. Effectiveness of substituted prolines as substrates was influenced by the position, size and the polar or non-polar nature of the substituent group.

At 0°C proline accumulation is prevented, but a stoichiometric exchange between intracellular and extracellular proline occurs (Britten & McClure, 1962). Under these conditions many proline analogues with affinity for the permease exchanged with previously accumulated L-[^{14}C]proline. However, some analogues, notably *trans*-3-methyl- and *trans*-4-methyl-L-proline, though possessing high affinity for the permease, showed an almost total inability to exchange with intracellular proline. These observations substantiate previous suggestions that proline uptake and exchange may not be closely interrelated (Britten & McClure, 1962; Tristram & Neale, 1968).

Britten, R. J. & McClure, F. T. (1962) *Bacteriol. Rev.* **26**, 292

Tristram, H. & Neale, S. (1968) *J. Gen. Microbiol.* **50**, 121