exact localization within the immunoglobulin fragments and chains is therefore of importance. Recent work in co-operation with Dr M. Turner and Dr H. Bennich on the pFc' fragment has added to the further localization of some of these markers within the Fc fragment (M. W. Turner, J. B. Natvig & H. Bennich, unpublished work). Gm(a), 'non a' and Gm(b⁰) are present within the pFc' fragment of the respective subclasses. In contrast, Gm(g) 'non g', Gm(n) and Gm(b¹) were not detected in the pFc' fragment and were probably related to structures nearer to the N-terminal part of the Fc portion.

These markers are also employed in studying immunoglobulin genes. The structural homologies, and the very close genetic linkages found by using the Gm and Am markers indicate that the γ G and the γ A subclass cistrons have developed through relatively recent duplications, due to unequal crossing-over. If this is so, one would expect to find evidence of unequal crossing-over, giving rise to deletions, duplications and hybridizations of the γ -globulin heavy-chain genes. In fact a number of gene complexes as well as single genes have been described that may be explained as result of crossing-over. There are several examples of probable crossing-over events appearing in various population groups. Further, evidence for products of deletions and duplications has been found both in major Gm-Am gene complexes and in isolated polypeptide chains. One particular case of a hybrid γ G heavy chain consisting of γ G1 Fc and γ G3 Fab (Kunkel, Natvig & Joslin, 1969) is an example of a gene deletion and hybridization due to an unequal homologous crossing-over.

The markers mentioned above are probably all part of the constant portion of the heavy chain. Further, for the light chain no genetic markers have so far been definitely localized to the variable portion. However, some reaction patterns with human anti- γ -globulin antibodies of the 'anti-antibody' type are related to the Gm allotypes. These 'anti-antibodies' cannot be inhibited by large excess of native γ G-globulin or F(ab')₂ fragments thereof. In contrast, they react easily with whole γ G-globulin or F(ab')₂ fragments altered through certain immunological reactions, particularly anti-Rh reactions. Since the 'anti-antibody' specificity is related to heavy-chain Gm allotypes and also to the antibody activity of the reactant γ G molecules, the antigenic determinants are likely to be in the variable part of the Fd fragment. Thus allotypic markers related to the Gm system may exist in the variable part of the γ G heavy chain.

Kunkel, H. G., Smith, W. K., Joslin, F. G., Natvig, J. B. & Litwin, S. D. (1969). *Nature, Lond.*, **223**, 1247.
 Natvig, J. B., Kunkel, H. G. & Joslin, F. G. (1969). *J. Immun.* **102**, 611.
 Vyas, G. N. & Fudenberg, H. H. (1969). *Clin. Res.* **17**, 469.

Genetic Markers of Rabbit Immunoglobulins

By J. M. WILKINSON. (*Department of Biochemistry, University of Birmingham, Edgbaston, Birmingham 15, U.K.*)

The occurrence of allotypes in rabbit immunoglobulins was first demonstrated by Oudin in 1956, when he showed that immunization of rabbits with purified antibody from some, but not all, other rabbits gave rise to precipitating antisera and that the antigenic specificities so detected were inherited in a simple Mendelian fashion (see Oudin, 1966). Since that time a considerable body of work has demonstrated the presence of more than 20 allotypic specificities on both heavy and light chains and in all three major classes of rabbit immunoglobulins. The most surprising finding has been that of Todd (1963) and Feinstein (1963), who showed that the α group determinants A1, A2 and A3 of the γ -chain are also shared by the α - and μ -chains, implying that, if the specificities are determined by primary structure, all three chains must have some sequence in common.

Sequence studies on such a heterogeneous population of proteins have been slow to catch up with the antigenic analysis, but it has been clear for some time from the amino acid analysis data for the whole chains and their fragments (Koshland, 1967; Prahl & Porter, 1968) that the differences in allotypic specificity could be correlated with differences in composition. The sequences responsible for the α and β group specificities and the γ -chain alleles A11 and A12 (Mandy & Todd, 1968) have been further investigated. Of these only the A11-A12 pair show a simple relation to sequence, there being a good correlation between a methionine-for-threonine replacement at position 226 in the γ -chain and the specificities A11 and A12 respectively (Prahl, Mandy & Todd, 1969).

The C-terminal sequence of the κ -chains, which carry the β group specificities, has been investigated by Appella, Rejnek & Reisfeld (1969). They have shown several differences between the sequence of the last seven amino acids in chains carrying A4, A5 and A6 specificities, and this has been confirmed by Frangione (1969), who also reported differences in the composition of the peptides containing the intrachain disulphide bonds. There seem thus to be multiple amino acid differences between the chains, as suggested by the amino acid composition, and this is surprising in view of the constancy of the C-terminal half of the human light chains.

Frangione, B., Franklin, E. C., Fudenberg, H. & Koshland, M. E. (1966). *J. exp. Med.* **124**, 715.
 Kunkel, H. G., Natvig, J. B. & Joslin, F. G. (1969). *Proc. natn. Acad. Sci. U.S.A.* **62**, 144.

The *N*-terminal sequence of the γ -chains has been studied in some greater detail, particularly for the allotypes A1 and A3 (Wilkinson, 1969a). The sequence of the first 34 residues has shown that there are 13 differences between the two sequences, and there are also a considerable number of positions that vary within each sequence; it is thus difficult to pinpoint those residues that may be responsible for the allotypic specificity. The sequence of the *C*-terminal half of the Fd region is known (Fruchter, Jackson, Mole & Porter, 1970), and as the serum pool used in this work was mainly of A1 specificity it may be taken as the A1 sequence; the corresponding sequence from A3 Fd fragment has been investigated in part and these data, taken with the peptide analyses of Koshland (1967), show that there is no allotype-related variation in this part of the sequence. A small amount of the remainder of the variable sequence has been investigated similarly and has so far failed to reveal further differences, but in view of the amino acid analysis data it is probable that more will be found. It would therefore seem that the amino acid sequence differences responsible for the *a* group specificities are present in the variable region of the γ -chain.

Preliminary results on the analysis and *N*-terminal sequence of α -chains of A1 and A3 specificity (Wilkinson, 1969b) suggest that similar allotype-related variations may be found in the α -chains. If this is true also of the μ -chains it would seem that the heavy chains of immunoglobulins share a similar set of variable sequences and that the constant parts of the sequence are class-specific.

- Appella, E., Rejnek, J. & Reisfeld, R. A. (1969). *J. molec. Biol.* **41**, 473.
 Feinstein, A. (1963). *Nature, Lond.*, **199**, 1197.
 Frangione, B. (1969). *FEBS Lett.* **3**, 341.
 Fruchter, R. G. Jackson, S. A., Mole, L. E. & Porter, R. R. (1970). *Biochem. J.* **116**, 249.
 Koshland, M. E. (1967). *Cold Spring Harb. Symp. quant. Biol.* **32**, 119.
 Mandy, W. J. & Todd, C. W. (1968). *Vox Sang.* **14**, 264.
 Oudin, J. (1966). *J. cell. Physiol.* **67**, (Suppl. 1), 77.
 Prah, J. W., Mandy, W. J. & Todd, C. W. (1969). *Biochemistry, Easton* (in the Press).
 Prah, J. W. & Porter, R. R. (1968). *Biochem. J.* **107**, 753.
 Todd, C. W. (1963). *Biochem. biophys. Res. Commun.* **11**, 170.
 Wilkinson, J. M. (1969a). *Biochem. J.* **112**, 173.
 Wilkinson, J. M. (1969b). *Nature, Lond.*, **223**, 616.

LIPID AND NEUROCHEMICAL GROUPS JOINT COLLOQUIUM ON 'CATABOLISM OF COMPLEX LIPIDS OF THE BRAIN'

Enzymic Hydrolysis of Glycolipids and Phospholipids

By SHIMON GATT. (*Department of Biochemistry, Hebrew University-Hadassah Medical School, Jerusalem, Israel*)

The research effort in this laboratory in the recent few years has been directed mainly to the isolation and characterization of enzymes that catalyse the degradation of glycolipids and phospholipids. This resulted in partial purification of the enzymes that catalyse the complete degradation of sphingolipids of animal origin and of a plant β -galactosidase that hydrolyses monogalactosyl diglyceride. The pathways of total degradation of the sphingosine bases were elucidated and several enzymes that hydrolyse phospholipids were purified.

Degradation of glycosphingolipids. Five sphingolipid-hydrolysing enzymes were extracted from brain tissue, partially purified and characterized. These are: (1) neuraminidase; (2) β -*N*-acetylhexosaminidase; (3) β -galactosidase; (4) β -glucosidase; (5) ceramidase. The neuraminidase was tightly bound to particulate fractions and could not be solubilized; it was therefore employed as a protein dispersion in Triton X-100. It did not act on sialyl-

lactose or a sialic acid-rich glycoprotein, but utilized as substrates several gangliosides in which *N*-acetylneuraminic acid was linked to a terminal galactose or to a second sialic acid residue. *N*-Acetylneuraminic acid was not removed from gangliosides when located at a branch point (i.e. where both the sialic acid and *N*-acetylglucosamine were attached to galactose, as in G_{M1} and G_{M2} gangliosides). Three β -*N*-acetylhexosaminidases were isolated from brain. One, solubilized from particles, hydrolysed both *N*-acetylglucosaminides and *N*-acetylglactosaminides. The other two were purified from the 100000g supernatant; one hydrolysed *N*-acetylglucosaminides and the other hydrolysed *N*-acetylglactosaminides. Only the enzyme that was solubilized from the particles hydrolysed the *N*-acetylglactosaminidic linkages of the two glycolipids globoside and 'asialo Tay-Sachs ganglioside' (*N*-acetylglactosaminidogalactosidoglucosidoceramide). Tay-Sachs (or G_{M2}) ganglioside was hydrolysed at only about 2% of the rate of hydrolysis of the above two glycolipids. The brain β -galactosidase hydrolysed most glycosphingolipids in which the galactose residue was terminal. However, galactosylceramide and galactosylsphingosine, in which galactose is attached directly to the sphingosine base, were not attacked by this enzyme.