- Jovin, T., Chrambach, A. & Naughton, M. A. (1964). *Analyt. Biochem.* **9**, 351.
 Mainwaring, W. I. P. (1969). *J. Endocr.* **45**, 531.
 Unhjem, O., Tveter, K. J. & Aakvaag, A. (1969). *Acta endocr., Copenh.*, **62**, 153.

Separation of Δ^4 -3-Oxo Steroids and Δ^5 -3 β -Hydroxy Steroids by Chromatography on Digitonin-Silica Thin Layers

By T. TAYLOR. (*University Department of Child Health, Royal Hospital for Sick Children, Yorkhill, Glasgow C.3, U.K.*)

In an investigation of Δ^5 -3 β -hydroxy steroid dehydrogenase (EC 1.1.1.51) activity in foetal adrenal glands, a method was required for the complete separation of nanogram amounts of Δ^4 -3-oxo steroids from a mixture also containing Δ^5 -3 β -hydroxy steroids. Digitonin precipitation of Δ^5 -3 β -hydroxy steroids is well known (Dorfman, 1962), but the present author has found that near-quantitative precipitation requires relatively large amounts of Δ^5 -3 β -hydroxy steroids (e.g. quantities of 3 β ,17 α -dihydroxypregn-5-en-20-one greater than 100 μ g), and even after multiple precipitation traces of Δ^5 -3 β -hydroxy steroid may remain in the supernatant.

A chromatographic method has been developed that enables complete separation of the two steroid groups on thin layers of silica gel impregnated with digitonin.

A layer (0.3mm thick) is spread on t.l.c. plates from a slurry containing: silica gel (Chromalux; May and Baker Ltd., Dagenham, Essex, U.K.), 30g; digitonin (Koch-Light Laboratories Ltd., Colnbrook, Bucks., U.K.), 3g; water, 48ml; ethanol, 2ml. The spread plates are dried at room temperature and then heated for 1 h at 100°C. After application of the steroid mixture, the plates are developed in the solvent system 2-methylpropan-2-ol-water-ethyl acetate (5:2:5, by vol.) for 4.5 h.

Δ^4 -3-Oxo steroids run as a group (R_F 0.86) whereas 1 μ g amounts of Δ^5 -3 β -hydroxy steroids exhibit R_F less than 0.65. The precise R_F value of individual Δ^5 -3 β -hydroxy steroids depends both on structure and the amounts present (e.g. for 10 ng of cholesterol, R_F 0; for 0.1 μ g and 10.0 μ g of 3 β ,17 α -dihydroxypregn-5-en-20-one, R_F 0.5 and 0.05 respectively). A co-operative effect exists whereby the total mass of Δ^5 -3 β -hydroxy steroids present determines the R_F value of each component Δ^5 -3 β -hydroxy steroid and of the group as a whole (e.g. R_F of 0.1 μ g of 3 β ,17 α -dihydroxypregn-5-en-20-one in the presence of 10 μ g of 3 β -hydroxypregn-5-en-20-one is 0.05). With 1 μ g of Δ^5 -3 β -hydroxy steroid present it is normally possible to obtain uncontaminated re-

covery (94–98%) of Δ^4 -3-oxo steroids from a complex mixture.

Elution of the Δ^4 -3-oxo steroid zone of the plate with ethyl acetate yields (digitonin-free) steroids. Digitonides from the Δ^5 -3 β -hydroxy steroid zone are eluted with methanol, dried and subjected to pyridine hydrolysis (Frame, 1944), yielding the free steroids.

This work was supported by a grant to Dr W. Hamilton from the Scottish Home and Health Department Advisory Committee on Medical Research.

Dorfman, R. I. (1962). In *Methods in Hormone Research*, 1st ed., vol. 1, pp. 86–90. Ed. by Dorfman, R. I. New York: Academic Press Inc.

Frame, E. G. (1944). *Endocrinology*, **34**, 175.

Effect of Oestradiol 3-Benzate on the Bio-synthesis of Egg Proteins by Cod

By P. A. PLACK and N. W. FRASER. (*Natural Environment Research Council Fisheries Biochemical Research Unit, University of Aberdeen, Aberdeen AB1 3RA, U.K.*)

The major protein constituents of cod eggs were found to be two similar lipoproteins, of mol.wt. about 400 000 and containing 79% protein and 21% lipid. From its phosphorus content, the lipid consisted of about 65% phospholipid. These two lipoproteins were not separated by gel filtration on Sephadex G-200 but were partly resolved by chromatography on triethylaminoethyl-cellulose (Jared & Wallace, 1968). Antiserum to cod eggs was prepared in rabbits and both immunodiffusion and immunoelectrophoresis gave two precipitin arcs close together with extracts of cod eggs.

For a fish of 4 kg wet wt., containing about 800 g of dry matter, some 100 g of egg proteins are produced during maturation of the ovaries over a period of 3–4 months. Egg proteins were detected in the serum of maturing female cod by immunochemical methods and reached a concentration of 32 mg/ml, but they were not present in the serum of immature fish or of male fish with maturing testes.

Oestradiol-17 β 3-benzoate in oil was injected intramuscularly into immature cod to give a slow release of the hormone. Serum samples taken 10 days after injection were tested for egg proteins, and the minimum dose of oestradiol to give a positive result was 30 μ g/kg; with doses of 300 μ g or 1 mg/kg the concentration of egg proteins was 32 mg/ml. Similar results were obtained with both male and female fish. Egg proteins were also present in the livers of injected fish but not in their ovaries, whereas immature control fish had no egg proteins in their livers. These findings suggested that the egg

proteins might be synthesized by the liver under the control of oestradiol.

Liver slices from immature fish incorporated L-[¹⁴C]leucine into material precipitable with trichloroacetic acid, the incorporation being inhibited by puromycin. Addition of carrier egg protein to an extract of slices and precipitation with antiserum gave no radioactivity in the precipitate. Liver slices from immature fish that had received an injection of 1 mg of oestradiol benzoate/kg 5 days previously gave a rate of incorporation of L-[¹⁴C]leucine into the trichloroacetic acid precipitate greater than that of the controls and also gave incorporation of L-[¹⁴C]leucine into the egg protein immunoprecipitate, the rate being about 40% of that for the trichloroacetic acid precipitate.

Jared, D. W. & Wallace, R. A. (1968). *Comp. Biochem. Physiol.* **24**, 437.

Steroid Metabolism by Foetal Sheep Adrenals

By ANNE B. M. ANDERSON, C. G. PIERREPOINT, K. GRIFFITHS and A. C. TURNBULL. (*Tenovus Institute for Cancer Research and the Department of Obstetrics, The Welsh National School of Medicine, Heath, Cardiff CF4 4XX, U.K.*)

The development and control of steroid-metabolizing enzyme systems in the foetus has received little attention. It is of interest therefore that destruction of the foetal sheep pituitary or infusion of synthetic adrenocorticotrophic hormone into the foetus *in utero* (Liggins, 1968, 1969) delays or advances parturition respectively. It has been hypothesized consequently that the developing pituitary-adrenal axis is responsible for the timing of parturition.

This preliminary study is concerned with the transformation *in vitro* of pregnenolone and progesterone by foetal sheep adrenals. Steroid metabolism has been investigated in adrenals from fetuses of different gestational age, from a foetus whose birth was prematurely induced by the intraperitoneal infusion of Synacthen *in utero* (790 µg over 76 h) and from the adult animal. Isotopically labelled pregnenolone and progesterone were incubated for 2 h at 39.5°C in Krebs-Ringer bicarbonate-glucose medium with ovine adrenal tissue from (a) adult non-pregnant ewe, (b) foetus at 122 days gestation, (c) foetus at 143 days gestation and (d) foetus at 116 days gestation after Synacthen infusion *in utero*. Steroids were extracted, isolated after extensive chromatography and derivative formation, and measured by procedures previously described (Pierrepoint, Anderson, Griffiths & Turnbull, 1969).

Foetal and adult adrenal tissues extensively metabolized pregnenolone. In the foetal adrenal at

122 days progesterone was the main product from pregnenolone, in contrast with the Synacthen-stimulated and term foetal adrenals, and also the adult gland, where there was a large conversion of this substrate into corticosteroids. Cortisol and corticosterone were the major products from pregnenolone in the Synacthen-stimulated foetal adrenal and in the adult, whereas a relative inactivity of the 11β-hydroxylation enzyme system was demonstrated in the term adrenal, 11-deoxycorticosterone and 11-deoxycortisol being the chief metabolites.

The effect of infused Synacthen on progesterone metabolism was not only to produce a greater transformation of the steroid but also to increase the production of cortisol and corticosterone compared with the unstimulated gland of similar gestational age. In the foetal tissue at 122 and 143 days 11-deoxycorticosterone was the major metabolite of progesterone, the yields of corticosterone and cortisol being relatively small. Progesterone was converted mainly into corticosterone by adult adrenal tissue.

No 3β-hydroxy steroid sulphokinase-transferase activity was demonstrated in either foetal or adult sheep adrenal tissue.

We thank the Wellcome Trust for generous financial support.

Liggins, G. C. (1968). *J. Endocr.* **42**, 323.

Liggins, G. C. (1969). In *Ciba Found. Symp.: Foetal Autonomy*, p. 218. Ed. by Wolstenholme, G. E. W. & O'Connor, M. London: J. and A. Churchill Ltd.

Pierrepoint, C. G., Anderson, A. B. M., Griffiths, K. & Turnbull, A. C. (1969). *Res. vet. Sci.* **10**, 477.

The Distribution of ¹³¹I-Labelled Adrenocorticotrophin in the Normal Guinea Pig

By M. P. GOLDER, R. MAHLER and A. R. BOYNS. (*Tenovus Institute for Cancer Research, Department of Metabolic Medicine, The Welsh National School of Medicine, Heath, Cardiff CF4 4XX, U.K.*)

The organ distribution of injected adrenocorticotrophin has been investigated by using labelled and unlabelled preparations (Richards & Sayers, 1951; Sonenberg, Keston & Money, 1951). These were often of lower purity than those currently available and the doses were outside the physiological range. The present experiments were carried out with synthetic α¹⁻²⁴-adrenocorticotrophic hormone labelled at high specific radioactivity with ¹³¹I and the doses injected were about 20–30-fold less than those previously reported.

Normal unanaesthetized guinea pigs were injected intravenously with α¹⁻²⁴-adrenocorticotrophic hormone (CIBA, Basle, Switzerland)