dase and deacetylase activities cannot be associated with the same enzyme and certainly not with a single active site. Moreover, it is clear that indoxyl acetate cannot be used as a substrate for the assay of hyaluronidase activity. Other results obtained during Sephadex gel chromatography of hyaluronidase activity suggest that the enzyme exists as a monomer (mol. wt. 70 000) in dilute solution and as a dimer (mol. wt. 120 000) at higher concentrations.

The authors thank Dr S. Gottfried of Biorex Laboratories Ltd. for sponsoring the present study.

Diczfalussy, E., Fernö, O., Fex, H., Höberg, B., Linderot, T. & Rosenberg, T. (1953). *Acta chem. scand.* **7**, 913.
Guilbault, G. G., Kramer, D. N. & Hackley, E. (1967). *Analyt. Biochem.* **18**, 241.

Effect of Bivalent Cations on some Enzymes of Gluconeogenesis

By J. M. WIMHURST and K. L. MANCHESTER.
(Department of Biochemistry, University College London, Gower Street, London W.C.1, U.K.)

Three of the enzymes important in gluconeogenesis, pyruvate carboxylase, phosphoenolpyruvate carboxykinase and fructose 1,6-diphosphatase, are all activated by both Mn^{2+} and Mg^{2+} . Surprisingly little information is available on the behaviour of pyruvate carboxylase and fructose 1,6-diphosphatase of rat liver towards these ions or towards Ca^{2+} , whose presence variously in the mitochondrion or cytoplasm has been suggested (Gevers & Krebs, 1966) as a possible mechanism for control of glucose metabolism through alternate inhibition of pyruvate carboxylase and pyruvate kinase. The effects of these ions on the three rodent hepatic enzymes important in the reverse of glycolysis were investigated.

Mn^{2+} is an activator of all three enzymes at concentrations much lower than are found for Mg^{2+} , a phenomenon in common with several other enzymes (e.g. RNA polymerase; Roeder & Rutter, 1969). In each case inhibition by Mn^{2+} is seen at concentrations at which Mg^{2+} concentration is optimal. When Mn^{2+} is added in addition to optimum concentrations of Mg^{2+} differing responses are seen. With pyruvate carboxylase addition of up to 2 mM- Mn^{2+} in the presence of 8 mM- Mg^{2+} has no effect on activity (Harris, Berent & Wimbhurst, 1970). With fructose 1,6-diphosphatase, which Tate (1966) found to be activated by EDTA saturated with Mg^{2+} , addition of Mn^{2+} at concentrations in excess of 0.05 mM-EDTA produced inhibition of the enzyme despite the presence of excess of Mg^{2+} . By contrast, Mn^{2+} in the range 0.01–0.1 mM in the presence of 6 mM- Mg^{2+} produced a marked

increase in the activity of phosphoenolpyruvate carboxykinase over that with Mg^{2+} alone, results similar to those of Foster, Lardy, Ray & Johnston (1967).

Pyruvate carboxylase, phosphoenolpyruvate carboxykinase and fructose 1,6-diphosphatase were all sensitive to inhibition by Ca^{2+} ; with the first two enzymes the loss of activity in the Mn^{2+} -activated system was less than when Mg^{2+} -activated but for fructose 1,6-diphosphatase the inhibition produced by Ca^{2+} was similar at optimum concentration of either Mn^{2+} or Mg^{2+} . Ca^{2+} inhibition of pyruvate carboxylase was noted by Kimmich & Rasmussen (1969), who also observed inhibition of its activity in mitochondria, and similar observations were made for phosphoenolpyruvate carboxykinase by Bartley & Dean (1969). Since key enzymes in gluconeogenesis, both those in the mitochondrion and in the cytoplasm, are subject to inhibition by Ca^{2+} it seems unlikely that extrusion of the cation will stimulate glucose formation. On the other hand exit of Ca^{2+} in conjunction with citrate (Harris & Berent, 1969) will tend towards inhibition of phosphofructokinase, but evidence is against enhanced citrate concentrations in liver under conditions favouring gluconeogenesis (Start & Newsholme, 1968).

We are grateful to the Medical Research Council for support for this work.

Bartley, W. & Dean, B. (1969). *Biochem. J.* **115**, 903.
Foster, D. O., Lardy, H. A., Ray, P. D. & Johnson, J. B. (1967). *Biochemistry, Easton*, **6**, 2120.
Gevers, W. & Krebs, H. A. (1966). *Biochem. J.* **98**, 720.
Harris, E. J. & Berent, C. (1969). *Biochem. J.* **115**, 645.
Harris, E. J., Berent, C. & Wimbhurst, J. M. (1970). *FEBS Lett.* **6**, 93.
Kimmich, G. A. & Rasmussen, H. (1969). *J. biol. Chem.* **244**, 190.
Roeder, R. G. & Rutter, W. J. (1969). *Nature, Lond.*, **224**, 234.
Start, C. & Newsholme, E. A. (1968). *Biochem. J.* **107**, 411.
Tate, S. S. (1966). *Biochem. biophys. Res. Commun.* **24**, 662.

Acylation of a Cerebral Sodium-plus-Potassium Ion-Stimulated Adenosine Triphosphatase Preparation

By I. PULL. (Department of Biochemistry, Institute of Psychiatry, British Postgraduate Medical Federation, University of London, London S.E.5, U.K.)

Conversion of amino groups in a particulate Na,K-ATPase* preparation into acidic functions by reaction with 2-methyl- and 2,3-dimethylmaleic anhydride has been examined as a procedure

* Abbreviation: Na,K-ATPase, $Na^{+} + K^{+}$ -stimulated adenosine triphosphatase.

for allowing separation of its constituents. These reagents are likely to be specific for amino groups, and the aminoacyl bonds they form (Dixon & Perham, 1968) appeared sufficiently labile to allow their hydrolysis under conditions not deleterious to Na,K-ATPase activity.

Microsomal suspensions were prepared from frozen ox cerebral cortex essentially by the procedure of Schwartz, Bachelard & McIlwain (1962) and then treated with an NaI reagent (Nakao, Tashima, Nagano & Nakao, 1965). Determination of amino groups in this preparation with 2,4,6-trinitrobenzenesulphonic acid (Habeeb, 1966) showed that lipid amino groups comprise about 55% of the total. Reaction with methylmaleic anhydride at pH 8 and 20°C causes a loss of Na,K-ATPase activity that is 50% complete when about 5% of the amino groups have been blocked, partial protection against inactivation being afforded by ATP. An 11-fold molar excess of methylmaleic anhydride acylates 90% of the total amino groups, and centrifugation of this acylated material at 150000g for 1 h gives a clear supernatant containing 30% of the protein and 5% of the phospholipid. Dimethylmaleic anhydride at pH 8.5 acylates the enzyme preparation less effectively than does methylmaleic anhydride at pH 8, but loss of Na,K-ATPase activity correlates similarly with the extent of acylation. No more than 43% of the total amino groups are dimethylmaleylated even when the anhydride is in 20- or 30-fold molar excess; centrifugation of this material gives a clear supernatant containing 45% of the protein and less than 10% of the phospholipid.

Regeneration of amino groups from the methylmaleylated material takes place slowly at pH 6 and 20°C, but Na,K-ATPase activity of unmodified material is lost to an extent precluding restoration of enzyme activity. Dimethylmaleyl groups are readily removed at pH 7 and there is partial restoration of Na,K-ATPase activity to the deacylated material provided that not more than 20% of the amino groups were initially acylated.

This investigation was supported by the Research Fund of the Bethlem Royal Hospital and Maudsley Hospital.

Dixon, H. B. F. & Perham, R. N. (1968). *Biochem. J.* **109**, 312.

Habeeb, A. F. S. A. (1966). *Analyt. Biochem.* **14**, 328.

Nakao, T., Tashima, Y., Nagano, K. & Nakao, M. (1965). *Biochem. biophys. Res. Commun.* **19**, 755.

Schwartz, A., Bachelard, H. S. & McIlwain, H. (1962). *Biochem. J.* **84**, 626.

Sodium-plus-Potassium Ion-Activated Adenosine Triphosphatase in a Heavy-Membrane Fraction Isolated from Rat Skeletal Muscle

By A. S. F. ASH and A. SCHWARTZ. (*Department of Pharmacology, Royal Free Hospital School of Medicine, University of London, London W.C.1, U.K., and Division of Myocardial Biology, Baylor College of Medicine, Houston, Tex. 77025, U.S.A.*)

An Mg^{2+} -dependent $Na^+ + K^+$ -activated adenosine triphosphatase, participating in active cation transport, has been isolated from a variety of tissues including cardiac (Matsui & Schwartz, 1966) and skeletal muscle (Samaha & Gergely, 1965). The enzyme activity is generally associated with membrane fractions derived from the endoplasmic reticulum and the plasma membrane (Skou, 1965). Current techniques for obtaining this adenosine triphosphatase involve treatment with deoxycholate. However, the results obtained with rat skeletal muscle were variable and the activity of the enzyme was low (Rogus, Price & Zierler, 1969). A method is now described for obtaining this $Na^+ + K^+$ -activated enzyme with high and reproducible specific activity. The enzyme was isolated at 0–4°C from a heavy-membrane fraction, without deoxycholate treatment, after solubilization of actomyosin.

Rat skeletal muscle was homogenized in 0.25 M-sucrose–1 mM-tris–EDTA buffer, pH 7.8, with an Ultra-Turrax tissue processor ($3 \times 3s$, 20000 rev./min). A precipitate collected at 600g (10 min) was treated with 0.6 M-KCl in 20 mM-tris-maleate buffer, pH 7.0, to solubilize actomyosin, rehomogenized and centrifuged again at 650g (15 min). The precipitate was twice washed in the sucrose–EDTA medium, collected at 10000g (15 min) and suspended in 1 mM-tris–EDTA buffer, pH 7.0, to give a protein concentration of 10 mg/ml.

An increase in the Mg^{2+} -dependent adenosine triphosphatase activity from 19.7 ± 1.18 to $28.7 \pm 1.16 \mu\text{mol of } P_i/\text{h per mg of protein}$, produced by addition of 100 mM- Na^+ with 10 mM- K^+ , was inhibited by ouabain ($K_i \ 8 \times 10^{-5} \text{ M}$). Total adenosine triphosphatase activity declined with time, but the ouabain-sensitive release of P_i from ATP (optimum pH 7.4) was linear over 5 min. The $Na^+ + K^+$ -stimulated activity was not enhanced by NaI treatment. The preparation was stable for several hours at 0°C, but activity declined overnight at 0°C or –20°C.

Inhibition of the $Na^+ + K^+$ -stimulated enzyme by ouabain (0.5 mM) was increased by 10% on addition of a low concentration of oligomycin (0.5 $\mu\text{g}/\text{mg}$ of protein); this suggests that some of the enzyme activity was of mitochondrial origin. At a higher concentration (5 $\mu\text{g}/\text{mg}$ of protein) oligomycin alone