The four activities are explainable on the basis of a single thiol-dependent enzyme and a Ping Pong Bi Bi mechanism with an alternate hydrolytic step.

C.M. is grateful for a World Health Organisation predoctoral fellowship. The work was supported by the Wellcome Trust.

Demain, A. L. (1966). In *Biosynthesis of Antibiotics*, vol. 1, p. 29. Ed. by Snell, J. F. New York: Academic Press Inc.

Gatenbeck, S. & Brunsberg, V. (1968). *Acta chem. scand.* **22**, 1059.

Pruess, D. L. & Johnson, M. J. (1967). *J. Bact.* **94**, 1502.

Spencer, B. (1968). *Biochem. biophys. Res. Commun.* **31**, 170.

Vanderhaeghe, H., Claesen, M., Vlietinck, A. & Parmentier, G. (1968). *Appl. Microbiol.* **16**, 1557.

Changes in Polyribosome Metabolism in *Escherichia coli* K12 after Adsorption of Colicin E3 to Sensitive Cells

By B. W. SENIOR and I. B. HOLLAND. (*Department of Genetics, University of Leicester, Leicester LE1 7RH, U.K.*)

Fixation of colicin E3, a simple protein, to specific receptors in the surface of sensitive bacteria rapidly results in the complete inhibition of protein synthesis. RNA synthesis, including mRNA synthesis, continues without inhibition, but 70S ribosomes extracted from bacteria, after prolonged colicin E3 treatment, show greatly decreased activity *in vitro* deriving from some specific alteration of the 30S subunit (Konisky & Nomura, 1967).

We have now shown that inhibition of protein synthesis induced by colicin E3 is soon followed by the fairly rapid loss of polyribosomes to yield equal proportions of both 70S and 50S+30S ribosomal subunits. This effect cannot be ascribed to increased nuclease activity. Moreover, before the onset of polyribosome degradation, abnormal polyribosome complexes can already be detected, which, unlike normal polyribosomes, dissociate at low Mg^{2+} concentrations. Thus within 1 min of the complete cessation of protein synthesis at least 40% of polyribosomes are demonstrably unstable when the Mg^{2+} concentration is lowered to 1 mM.

'Pulse-chase' experiments *in vivo* indicate that the release of nascent polypeptide chains from polyribosomes is inhibited in colicin-treated cells. In contrast, other experiments indicate that protein synthesis is blocked as a result of inhibition of some initiation step. The changes in ribosome function provoked by the adsorption of colicin E3 may therefore affect a step in the protein-biosynthetic cycle, common to both polypeptide-chain extension

and initiation, as has been suggested for streptomycin (Modolell & Davis, 1969).

Colicin E3 has no effect on ribosomes *in vitro* (Konisky & Nomura, 1967) and like other colicins it does not apparently penetrate the bacterial cell surface. The colicin acts extracellularly, presumably by promoting a specific alteration of the cytoplasmic membrane. Inhibition of ribosome function is not, however, dependent on continued RNA or protein synthesis. The surface-bound colicin must therefore trigger the 'activation' (or inhibition) of some normal cell constituent, presumably membrane-bound, which then effects modification of the 30S ribosomal subunit.

B.W.S. is a Beit Memorial Fellow and the receipt of this Fellowship is gratefully acknowledged.

Konisky, J. & Nomura, M. (1967). *J. molec. Biol.* **26**, 181.
Modolell, J. & Davis, B. D. (1969). *Nature, Lond.*, **224**, 345.

Activity of Skeletal-Muscle Ribosomes from Intact and Castrated Rats treated with Triamcinolone Acetonide

By R. F. PETERS and A. M. WHITE. (*CIBA Laboratories Ltd., Horsham, Sussex, U.K.*)

Triamcinolone acetonide was administered at doses of 20, 5 and 0.2mg/kg by intraperitoneal injection into intact adult male rats (250g, Carworth CFHB Strain) or similar rats castrated 10 days previously. In each experiment ribosomes were isolated from five control animals (intact or castrated) and five drug-treated animals 12h after the administration of the drug. The amino acid-incorporating ability *in vitro* of the ribosome preparation from each animal was measured over a 20 min incubation period in the presence of diluted liver cell sap and energy (Bullock, White & Worthington, 1968). The method for the isolation of ribosomes was essentially as described previously (Peters, Richardson, Small & White, 1970) except that 0.1M-tris buffer was used in the homogenizing medium and treatment with amylase was omitted.

The means of the amino acid-incorporating activities of ribosomes from intact animals at triamcinolone doses of 20 and 5mg/kg were both 79% of control, and in castrated animals were 73 and 79% of control, respectively. However, at 0.2mg/kg the mean activity of ribosomes from intact animals was 93% of control, whereas that from castrated animals was 77%. When castrated animals were pretreated daily for 7 days with testosterone (2.0mg/kg per day) the mean ribosomal activity 12h after administration of triamcinolone acetonide (0.2mg/kg) was elevated to 90% of

control. In all results cited the differences between treated and control animals were statistically significant ($P < 0.05$). It is apparent therefore that the castrated animals were more sensitive to the action of the drug, when given at low doses than the intact animals.

In contrast with the 50% decrease in incorporating ability and the lower polyribosome content of ribosomes from skeletal muscle of castrated immature rats (Breuer & Florini, 1965), we could find only a slight decrease (92% of intact control) in the incorporating ability of ribosomes from castrated adult rats and virtually no differences between the sucrose density gradients of ribosomes from intact and castrated animals.

These results are consistent with the idea that the susceptibility of muscular protein synthesis to the antianabolic action of anti-inflammatory glucocorticoids is, in spite of previous findings (Bullock, Peters & White, 1969), to some extent moderated by androgens.

Breuer, C. B. & Florini, J. R. (1964). *Biochemistry, Easton*, 4, 1544.

Bullock, G. R., Peters, R. P. & White, A. M. (1969). *Biochem. J.* 115, 47P.

Bullock, G., White, A. M. & Worthington, J. (1968). *Biochem. J.* 108, 417.

Peters, R. F., Richardson, M. C., Small, M. & White, A. M. (1970). *Biochem. J.* 116, 349.

Effect of Cortisone on Transfer Ribonucleic Acid and Aminoacyl-Transfer-Ribonucleic Acid Synthetases in Rat Liver

By M. K. AGARWAL* and J. HANOUNE.† (*College of Physicians and Surgeons, Columbia University, New York, N.Y. 10032, U.S.A.*)

Prior RNA synthesis is consistently given as the explanation for the subsequent induction of selected enzymes by glucocorticoids in mammalian liver. Because of the central role of tRNA in translation of the genetic message and because of the known quantitative and/or qualitative changes in these parameters under a variety of pathophysiological syndromes (reviewed by Baliga, Borek, Weinstein & Srinivasan, 1969), including the changes brought about by two animal hormones [oestrogen on chick oviduct (O'Malley, Aronow, Peacock & Dingman, 1968) and thyroxine on rat liver (Yang & Sanadi, 1969)] and a plant hormone [6-benzyladenine (Anderson & Cherry, 1969)], it was important to determine whether the 20–30% increase in

tRNA synthesis (with orotic acid as the precursor), observed here as early as 30 min after cortisone administration, reflected general increase in all tRNA species or stimulation of selected tRNA species.

Co-chromatography of doubly labelled tRNA, prepared from saline- or cortisone-injected rats, in presence of [^{14}C]- and [^3H]-orotic acid respectively, on a benzoylated DEAE-cellulose column, where each type of tRNA is eluted as a sharp peak at a characteristic position, revealed a constant $^3\text{H}/^{14}\text{C}$ ratio over the entire range of u.v.-absorbing material, indicating that all tRNA species were uniformly labelled. Similarly, the amino acid-acceptance capacities were unaltered after cortisone administration when tested with synthetases both from normal and from hormone-injected rats. Possible alterations in selected isoaccepting species (see references) were next investigated. Co-chromatography on an RPC-Freon column of tRNA^{Leu} or tRNA^{Tyr}, obtained from saline- or cortisone-treated rats and aminoacylated *in vitro*, did not reveal qualitative differences in the isoaccepting species for these two tRNA species. Modification of tRNA function by cortisone through attachment or incorporation into tRNA was investigated by density-gradient centrifugation. No binding to tRNA of [^3H]cortisone administered *in vivo* or [^3H]cortisol *in vitro* was evident.

From these and other studies it seems that the general increase in liver tRNA synthesis effected by cortisone is likely to be a prelude to, and part of, the hepatomegaly that follows hormone administration, and is probably not related to the induction of specific enzymes by the glucocorticoid.

Anderson, M. B. & Cherry, J. H. (1969). *Proc. natn. Acad. Sci. U.S.A.* 62, 202.

Baliga, B. S., Borek, E., Weinstein, I. B. & Srinivasan, P. R. (1969). *Proc. natn. Acad. Sci. U.S.A.* 62, 899.

O'Malley, B. W., Aronow, A., Peacock, A. C. & Dingman, E. W. (1968). *Science, N.Y.*, 162, 567.

Yang, S. S. & Sanadi, D. R. (1969). *J. biol. Chem.* 244, 5081.

Incorporation of ^{14}C from [^{14}C]Glucose into Amino Acids by Rumen Bacteria

By ROSALIND J. ZALIN (introduced by M. A. CRAWFORD). (*Nuffield Institute of Comparative Medicine, Zoological Society of London, Regent's Park, London N.W.1, U.K.*)

The extent to which the rumen microbial population can synthesize amino acids from carbohydrate and simple nitrogen sources and thus supplement the low amounts of amino nitrogen in the ruminant diet is largely unknown. However, pure culture

* Present address: Department of Zoology, University of Sheffield, Sheffield S10 2TN, U.K.

† Present address: Institut de Pathologie Moléculaire, Paris, France.