quantities of pure DDE to rats (≤ 150 p.p.m.) and quail (≤ 100 p.p.m.) for 21 days. DDT was fed at a concentration of 150 p.p.m. to rats and 100 p.p.m. to quail.

All residue concentrations were considerably higher in quail than in rats fed at the same dose rate. The control values of all the other parameters were higher in rats than quail. DDE produced larger increases in hepatic microsomal protein, cytochrome *P*-450 and aniline oxidase activity in both species than DDT fed at the same dietary concentration. DDE also produced a larger increase in relative liver weight in rats, but no changes of relative liver weight were produced in quail on any diet. In general the increases produced by DDE in the rat parameters were larger relative to controls than those produced in quail given equivalent dietary concentrations and the increases become apparent at lower concentrations. It would appear that the microsomal oxidase system in quail is less sensitive to DDE than that in the rat.

It seems probable that DDE is a more potent inducer of microsomal enzymes than DDT, particularly at low dosages. Thus interspecies differences in the effects of DDT on hepatic enzyme systems may depend on both the rate of metabolism, and the amount of DDE it produces, rather than on the parent compound.

- Bailey, S., Bunyan, P. J., Rennison, B. D. & Taylor, A. (1969a). *Toxic. appl. Pharmac.* **14**, 13.
 Bailey, S., Bunyan, P. J., Rennison, B. D. & Taylor, A. (1969b). *Toxic. appl. Pharmac.* **14**, 23.
 Gillette, J. W., Chan, T. M. & Terriere, L. C. (1966). *J. Agric. Fd Chem.* **14**, 540.
 Hart, L. G. & Fouts, J. R. (1965). *Naunyn-Schmiedeberg's Arch. exp. Path. Pharmac.* **249**, 486.
 Kato, R. & Gillette, J. W. (1965). *J. Pharmac. exp. Ther.* **150**, 279.
 Lowry, O., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
 Omura, T. & Sato, R. (1964). *J. biol. Chem.* **239**, 2370.
 Peterson, J. E. & Robison, W. H. (1964). *Toxicol. appl. Pharmac.* **6**, 321.
 Taylor, A., Rea, R. E. & Kirby, D. R. (1964). *Analyst*, **89**, 497.

Microbial Degradation of 4-Carboxy-1-methylpyridinium Chloride, a Photolytic Product of Paraquat.

By K. A. WRIGHT and R. B. CAIN. (*Department of Botany, University of Newcastle upon Tyne, Newcastle upon Tyne NE1 7RU, U.K.*)

Washed suspensions of *Achromobacter* D utilizing MINA* released methylamine originating from the *N*-methyl group of the molecule (Wright & Cain,

* Abbreviation: MINA, 4-carboxy-1-methylpyridinium chloride (*N*-methylisonicotinic acid).

1969). Further work with cell-free extracts has now enabled us to identify the other end products of this degradation and their origin in the carbon atoms of the pyridine ring.

Hydroxylation does not seem to be involved in the metabolism of MINA because none of the possible mono- or di-hydroxylated derivatives was utilized by either washed suspensions or extracts that nevertheless released methylamine, nor was MINA itself hydroxylated in the model system of Udenfriend, Clark, Axelrod & Brodie (1954), a system that readily hydroxylated other non-quaternary pyridine compounds. Extracts utilized MINA only when supplemented with NAD^+ or NADH and then took up 1 mol of O_2 /mol of MINA degraded. NADH was first oxidized to NAD^+ by crude extracts and subsequently re-formed; these two stages could be separated by ammonium sulphate fractionation. Added NAD^+ was reduced to NADH only after a lag period.

With either NAD^+ or NADH supplementation, extracts catalysed the evolution of CO_2 , which was shown to arise from the carboxyl group on C-4 of the pyridinium ring by using [*carboxy*- ^{14}C]MINA. The same supplemented extracts released [^{14}C]methylamine from *N*-[*Me*- ^{14}C]MINA. The remaining five carbon atoms of the pyridinium nucleus were accounted for as formate and succinate, one of which accumulated in sufficient amounts for it to be isolated and identified chemically by mixed m.p., mass spectrometry, t.l.c. and g.l.c. of the appropriate esters. [2,3,5,6- ^{14}C]MINA gave rise to [^{14}C]formate and labelled succinate; the distribution of the radioactivity among the carbon atoms of succinate was consistent with its origin in C-3, C-5 and C-6 of the pyridine ring, and the formate arose from C-2.

The NADH and oxygen requirement, typical of that of an oxygenase, and the absence of hydroxylation indicate the possibility of direct oxidative fission of a partly reduced ring to form a dialdehyde (Sparrow *et al.* 1969), which is subsequently hydrolysed to release first formate and then methylamine, and generate succinic dialdehyde. Oxidation of the latter through succinic semialdehyde to succinate is supported by the observation that extracts prepared from MINA-grown cells have some 15-fold higher succinic semialdehyde activities than those prepared from succinate-grown cultures.

This study was supported by a grant from the Science Research Council. The ring-labelled MINA was kindly provided by Dr B. C. Baldwin.

- Sparrow, L. G., Ho, P. P. K., Sundaram, T. K., Zach, D., Nyns, E. J. & Snell, E. E. (1969). *J. biol. Chem.* **244**, 2590.
 Udenfriend, S., Clark, C. T., Axelrod, J. & Brodie, B. B. (1954). *J. biol. Chem.* **208**, 731.

Wright, K. A. & Cain, R. B. (1969). *Soil Biol. Biochem.* **1**, 5.

Benzoate Oxidation by Bacterium N.C.I.B. 8250

By C. A. FEWSON, A. LIVINGSTONE and P. J. ROACH.
(Department of Biochemistry, University of Glasgow,
Glasgow W.2, U.K.)

Although Ichihara, Adachi, Hosokawa & Takeda (1962) described a cell-free system for benzoate oxidation in *Pseudomonas aeruginosa* and *Micrococcus ureae*, several workers have experienced difficulties in measuring benzoate oxidase activity in bacteria and have had to resort to experiments with intact organisms (e.g. Mandelstam & Jacoby, 1965; Kemp & Hegeman, 1968; Kennedy & Fewson, 1968).

We have developed a specific, rapid and sensitive method for measuring benzoate consumption by whole cells or by extracts. This is based on u.v.-absorption measurements of residual benzoic acid after acidification of reaction mixtures and extraction through chloroform. It has several advantages over previous methods, which have generally depended on measurement of oxygen consumption with consequent uncertain corrections for oxygen uptake caused by endogenous metabolism, NADH oxidation or metabolism of the products of benzoate oxidation.

Whole cells of bacterium N.C.I.B. 8250 grown in a benzoate-salts medium oxidise benzoate at a rate of about 70nmol/min per mg of protein. After ultrasonic disruption and centrifugation at 10000g for 20min the supernatants show a low but reproducible benzoate oxidase activity (about 2nmol/min per mg of protein). Optimum activity is obtained by extraction and assay in tris buffer, pH 7.5. The enzymic activity has a specific requirement for Fe²⁺ ions. Preincubation with cupferron (30μM) or 2,2'-bipyridyl (100μM), but not sodium diethyldithiocarbamate (30μM), prevents benzoate oxidation. There is a requirement for FAD (5μM). Enzymic activity also depends on the continuous regeneration of NADH (ethanol and alcohol-NAD oxidoreductase, EC 1.1.1.1). NADPH is not effective. Benzoate does not disappear under anaerobic conditions. Various combinations of ATP, CoA, thiamin pyrophosphate, lipoic acid, Mg²⁺, mercaptoethanol and GSH have no beneficial effect on the rate of benzoate oxidation. A compound giving a positive Rothera reaction is formed as benzoate disappears from the reaction mixture. Presumably the active catechol oxygenase (EC 1.13.1.1) (Kennedy & Fewson, 1968) accounts for the absence of catechol accumulation.

The addition of extracts of bacterium N.C.I.B. 8250 inhibits benzoate oxidation by intact cells. This is superficially reminiscent of the situation in *Rhodopseudomonas palustris* (Dutton & Evans, 1968), but it is unlikely that the mechanism of inhibition is the same since the pathways of benzoate metabolism are quite different in the two organisms.

A.L. and P.J.R. are in receipt of Science Research Council Studentships.

Dutton, P. L. & Evans, W. C. (1968). *Biochem. J.* **107**, 28p.

Ichihara, A., Adachi, K., Hosokawa, K. & Takeda, Y. (1962). *J. biol. Chem.* **237**, 2296.

Kemp, M. B. & Hegeman, G. D. (1968). *J. Bact.* **96**, 1488.

Kennedy, S. I. T. & Fewson, C. A. (1968). *Biochem. J.* **107**, 497.

Mandelstam, J. & Jacoby, G. A. (1965). *Biochem. J.* **94**, 569.

Metabolism of Myricetin, Myricitrin and 3,4,5-Trihydroxyphenylacetic Acid

By G. E. SMITH and L. A. GRIFFITHS. (Department of Biochemistry, University of Birmingham, Birmingham 15, U.K.)

Previous studies on the metabolism of flavonoid compounds in mammals have largely been restricted to flavonoids showing hydroxylation in the 4'- or 3', 4'-positions only in the B ring, although flavonoids possessing a 3',4',5'-trihydroxyphenyl grouping are known to occur in plants of dietary importance.

Administration of myricetin (3,5,7,3',4',5'-hexahydroxyflavone) (200mg) to rats maintained on our standard diet gave rise to a major metabolite in the urine that has been identified as 3,5-dihydroxyphenylacetic acid by chromatographic and spectral methods. No glucuronide conjugates of myricetin were detected.

Since earlier studies have shown that the degradation of flavonoids is in part mediated by intestinal micro-organisms (Griffiths, 1964; Nakagawa, Shetlar & Wender, 1965; Das & Griffiths, 1968), the ability of intestinal micro-organisms to degrade myricetin *in vitro* was studied by incubating myricetin in a thioglycollate broth with mixed inocula of micro-organisms obtained by sterile section of the rat caecum. The incubation was carried out under anaerobic conditions at 37°C. The metabolites were recovered by a modification of a previously described ether extraction procedure (Griffiths, 1964) and shown to be 3,5-dihydroxyphenylacetic acid, 3,4,5-trihydroxyphenylacetic acid and unchanged myricetin.

Since it appeared probable that 3,4,5-trihydroxyphenylacetic acid was an intermediate in the