Partial Purification and Properties of Guinea-Pig Liver Polynucleotide Phosphorylase

By YEOW PHEW SEE AND P. S. FITT

Department of Biochemistry, University of Ottawa, Ottawa 2, Ont., Canada

(Received 5 May 1970)

1. Polynucleotide phosphorylase has been isolated and partially purified from crude preparations of guinea-pig liver nuclei. 2. The enzyme is particulate and associated with RNA and lipids characteristic of membranes. 3. It has phosphorolysis and exchange activities, but the latter may be due to a contaminating enzyme. 4. The phosphorolysis activity is dependent on bivalent cations, preferably Mg^{2+} , has a pH optimum between 8.6 and 9.2 and is inhibited by potassium chloride and sodium chloride. 5. The enzyme catalyses phosphorolysis of poly A, poly C, poly U, rRNA and tRNA. Poly G is only phosphorolysed to a very small extent and DNA is not a substrate. 6. The enzyme appears to lack nucleoside diphosphate polymerization activity.

The isolation and proof of specificity of a polynucleotide phosphorylase from guinea-pig liver was recently reported (Fitt & See, 1970). This was the first unequivocal evidence for the presence of polynucleotide phosphorylase in organisms other than bacteria.

The enzyme was isolated in a particulate form and catalysed phosphorolysis of polyribonucleotides and RNA to nucleoside diphosphates. NDP- P_i and NTP- P_i exchange activities were present in the partially purified preparation, but could have been due to a separate enzyme and no nucleoside diphosphate polymerization activity was detected. The enzyme appeared to be associated with the cell nucleus.

We now describe in detail the partial purification and the enzymic properties of guinea-pig liver polynucleotide phosphorylase.

EXPERIMENTAL

Materials. Animals, chemicals and intermediates were purchased from the following suppliers: [^{14}C]ADP and [^{32}P]P $_i$, Amersham/Searle Corp., Don Mills, Ont., Canada; sodium dodecyl sulphate (specially pure grade), B.D.H. (Canada) Ltd., Toronto, Ont., Canada; high-molecular-weight wheat-germ RNA (rRNA), Calbiochem, Los Angeles, Calif., U.S.A.; Merck silica gel G, Canlab, Ottawa, Canada; ApA, Gallard-Schlesinger Manufacturing Corp., Carle Place, N.Y., U.S.A.; poly A, poly C, poly G and poly U, Miles Chemical Co., Elkhart, Ind., U.S.A.; NMP, NDP and NTP, P-L Biochemicals Inc., Milwaukee, Wis., U.S.A.; Whatman DE23 DEAE-cellulose, Reeve Angel, Clifton, N.J., U.S.A.; guinea pigs,

Robidoux, St Constant, P.Q., Canada; yeast tRNA and calf thymus DNA, Sigma Chemical Co., St Louis, Mo., U.S.A.; pancreatic ribonuclease A, Worthington Biochemical Corp., Freehold, N.J., U.S.A.

Buffers. Except where stated, the pH of buffers refers to the value at 25°C.

Preparation of guinea-pig liver polynucleotide phosphorylase. The enzyme was prepared by a modification of the method described by Fitt & See (1970). The acetone precipitation step used in the earlier experiments had little effect and was omitted. Unless otherwise stated, all operations were carried out at 2–4°C.

Crude nuclei were isolated by the method of Chauveau, Moulé & Rouiller (1956). Six fed male guinea pigs (700–900 g each) were decapitated and bled. Their livers were removed and chilled in 100 ml of ice-cold 0.25 M-sucrose–1 mM- β -mercaptoethanol–10 mM-tris-HCl buffer, pH 7.4 ('sucrose-tris buffer'). The tissue was weighed, dried with absorbent paper, transferred to 2.2 M-sucrose–10 mM-tris-HCl buffer, pH 7.4, and cut into small pieces. The mixture was homogenized in a Lourdes model MM-1 homogenizer for 90 s at full speed and the homogenate was filtered through four layers of cheesecloth. The filtrate was left at 0°C for 20 min and was then centrifuged at 40 000 g for 1 h in a refrigerated centrifuge (International model B-20). The supernatants were discarded and the walls of the centrifuge tubes were wiped carefully with absorbent paper. The pellets were transferred to a Dounce glass homogenizer with sucrose-tris buffer (30 ml) and the nuclei were suspended in the buffer with three strokes of a loose-fitting pestle. The suspension was kept at 0°C for 20 min and then centrifuged at 40 000 g for 10 min. The supernatant was collected and the pellet was re-extracted twice with 30 ml portions of sucrose-tris buffer. The supernatants from the three extractions were combined to give the crude sucrose extract (Table 1). The pellet was extracted a further two times with 30 ml

portions of sucrose-tris buffer, and the supernatants were discarded. The pellet was then extracted three times, the same procedure being used, with 0.14M-NaCl-1mM- β -mercaptoethanol-2mM-EDTA, pH 7 (total volume 50ml) to give the crude saline extract (Table 1). Both the crude sucrose and saline extracts were dialysed separately overnight against 2l of 1mM- β -mercaptoethanol-2mM-EDTA whose pH was adjusted to 7 at 4°C ('BME-EDTA').

The dialysed sucrose extract was centrifuged at 40000g for 20 min to remove insoluble material. The pH of the stirred, cooled (0°C) supernatant was adjusted to 5.7 with 1M-sodium acetate buffer, pH 5.3, and the suspension was stirred for a further 3 min. The precipitate was collected by centrifugation and dissolved in 40 ml of 2mM-EDTA-1mM- β -mercaptoethanol-10mM-imidazole-HCl buffer, pH 7.2 ('imidazole buffer'). The cloudy solution was centrifuged and the pellet was discarded. The supernatant was applied the same day (at 30 ml/h) to a column (1.7 cm $\times$ 16 cm) of DEAE-cellulose (Whatman DE23) previously equilibrated with imidazole buffer. The column was washed with imidazole buffer until no more protein emerged, and the enzyme was then eluted at 60 ml/h with 0.3M-NaCl in imidazole buffer (30 ml; 3 ml fractions). The individual fractions were dialysed together overnight against BME-EDTA (2l) and assayed for enzymic activity. All the active enzyme was precipitated during the dialysis. The active precipitates were collected by centrifugation and washed twice by resuspension in BME-EDTA (5 ml) followed by centrifugation. The solid was collected and stirred with BME-EDTA (5 ml) until a fine suspension was obtained. The final product contained 3800 units, specific activity 680 units/mg (Table 1), and was used for all the experiments described in this paper unless otherwise stated. It is referred to as 'the sucrose fraction'.

A purer particulate enzyme (specific activity about 1100-1200 units/mg) could be obtained if the DEAE-cellulose column was eluted stepwise with 0.1M-, 0.2M- and 0.3M-NaCl in imidazole buffer, as previously described (Fitt & See, 1970). However, the yield was greatly decreased. The simplified procedure described above was used in this work since relatively large amounts of enzyme were needed and the only significant difference between the products obtained by the two methods appeared to be a slightly greater contamination of the less pure enzyme by phosphatases.

The dialysed saline extract was centrifuged at 40000g for 20 min to remove solid material. The pH of the clear supernatant was adjusted to 5.7 with 1M-sodium acetate buffer, pH 5.3, and the mixture was stirred for a further 3 min. The precipitate was collected by centrifugation, suspended in BME-EDTA (1 ml) and dialysed overnight against BME-EDTA (200 ml). The product contained 228 units, specific activity 253 units/mg (Table 1), and is referred to as the saline fraction.

Enzyme assays. Polynucleotide phosphorylase activity was determined by the phosphorolysis assay, which depends on the formation of [32 P]NDP during phosphorolysis of polyribonucleotides in the presence of [32 P] P_i (Fitt & See, 1970; Fitt & Fitt, 1967; Grunberg-Manago, Ortiz & Ochoa, 1956). The standard assay medium (0.1 ml) contained: 0.1M-tris-HCl buffer, pH 8.7; 4mM-MgCl₂; 0.6mM-EDTA; 1.3mM- β -mercaptoethanol; 10mM-K₂H₃₂PO₄ (30000-100000 c.p.m./ μ mol); poly A,

1.5mg/ml; enzyme. Incubation was at 37°C for 1 h in an Eberbach water bath-shaker at 300 strokes/min. The reaction was stopped with 1 ml of a mixture containing 1 part of an aqueous 10% (w/w) suspension of acid-washed Norit A and 10 parts of 2.5% (w/v) HClO₄. The charcoal suspension was left for 10 min at 0°C and was then collected on a 2.4 cm diam. Whatman GF/C glass-fibre filter. The filter was washed four times with 10 ml of water and was then inverted in a planchet, dried and its radioactivity determined in a Nuclear-Chicago low-background thin-window counter. The specific radioactivity of the K₂H₃₂PO₄ was determined by counting a dried portion of the solution over which a GF/C filter had been placed. Suitable controls were carried out in all cases and their radioactivities subtracted from the assay values. A unit was defined as the amount of enzyme catalysing incorporation of 1 nmol of [32 P] P_i into charcoal-adsorbable material/h.

NDP- P_i exchange activity was assayed by measuring the incorporation of [32 P] P_i into nucleoside diphosphates (Grunberg-Manago *et al.* 1956). The reaction mixture (0.1 ml) contained: 0.1M-imidazole-HCl buffer, pH 7.8; 10mM-MgCl₂; 0.6mM-EDTA; 1.3mM- β -mercaptoethanol; 10mM-K₂H₃₂PO₄ (30000-100000 c.p.m./ μ mol); 1mM-NDP; enzyme. The assay was performed as described for the phosphorolysis assay.

NTP- P_i exchange activity was assayed by measuring the incorporation of [32 P] P_i into nucleoside triphosphates. The conditions and procedure were the same as for the NDP- P_i exchange assay except that the pH was 7.4 and the substrate was NTP instead of NDP.

Ribonuclease activity in the particulate enzyme was measured by a modification of the procedure of Kalnitsky, Hummel & Dierks (1959). The reaction mixture (0.1 ml) contained: 0.1M-tris-HCl buffer, pH 8.7; 4mM-MgCl₂; 0.6mM-EDTA; 1.3mM- β -mercaptoethanol; poly A or wheat germ RNA, 150 μ g; enzyme. Incubation was at 37°C in a water bath-shaker for 1 h. The reaction was stopped with 0.5 ml of 0.75% (w/v) uranyl acetate in 25% (w/v) HClO₄. The mixture was kept in ice for 15 min and then centrifuged. A sample (50 μ l) of the supernatant was diluted to 1 ml with water and the E_{260} was determined by using as blank a similar, incubated sample that lacked enzyme.

Nucleic acid assay. RNA was determined by the orcinol reaction by using the procedure of Dische (1955). The difference in the extinction of the test solutions at 675 nm and 575 nm was used for calculation and the internal glucose standard was omitted. A standard curve was determined with wheat germ RNA. DNA was determined by the diphenylamine procedure of Burton (1956) with dAMP as standard. The particulate guinea-pig enzyme was dissolved in 5mM-sodium dodecyl sulphate and the detergent was present at the same concentration in blanks and standard samples.

Protein assay. Protein was determined by the method of Lowry, Rosebrough, Farr & Randall (1951). The particulate enzyme was first dissolved by addition of sodium dodecyl sulphate to a final concentration of 5 mM and the detergent was present in the blanks in the same concentration. A standard curve was determined with pancreatic ribonuclease A, the same concentrations of EDTA, β -mercaptoethanol and sodium dodecyl sulphate being used as were present in the enzyme samples.

Table 1. Purification of guinea-pig liver polynucleotide phosphorylase

The enzyme was prepared as described in the Experimental section. One unit is the amount of enzyme that catalyses the incorporation of 1nmol of [³²P]P_i into ADP/h in the conditions of the poly A-phosphorolysis assay.

Step	Sucrose fraction			Saline fraction		
	Units	Protein (mg)	Sp. activity (units/mg)	Units	Protein (mg)	Sp. activity (units/mg)
1. Crude extract	7800	194	40	482	6.5	74
2. pH5.7	11200	66	170	228	0.9	253
3. DEAE-cellulose, 0.3M-NaCl eluate	3800	5.6	680	—	—	—

RESULTS AND DISCUSSION

Preparation of guinea-pig liver polynucleotide phosphorylase. The results of a typical isolation of the guinea-pig liver enzyme are shown in Table 1. The final sucrose and saline fractions had very similar properties (see below), but the results in Fig. 1 show that they were cleanly separated by the extraction procedure. The latter did not rupture the nuclei since viscous nucleoprotein was not liberated. Further extraction of the nuclear pellet with more concentrated sodium chloride solutions failed to yield additional polynucleotide phosphorylase.

The purified particulate sucrose fraction contained RNA (about 10%, w/w) and lipids characteristic of membranes (Table 2), but no detectable DNA. To verify if the enzyme was in a particulate form in the crude extract, a dialysed sucrose extract was centrifuged at 40000*g* for 20min to remove visible solids and then at 100000*g*. After 1½ h, 53% of the activity had sedimented and this was raised to 69% after 3¼ h, suggesting that the enzyme was attached to membrane fragments of various sizes. This would explain the loss of about 65% of the activity during DEAE-cellulose chromatography (Table 1), when the largest particles would collect on the surface of the ion-exchanger. Further, since all the activity eluted from the column was recovered in particulate form after removal of the sodium chloride by dialysis, and the saline fraction was also solid, it seems certain that all the enzyme was membrane-bound. The nuclear pellet obtained by the 2.2*M*-sucrose method is very small, as would be expected after isolation of the nuclei in an extremely hyperosmotic medium. During extraction with 0.25*M*-sucrose, the pellet swells abruptly to several times its original volume and it seems probable that some part of the double-membrane system is broken away from the nuclei without destroying their overall integrity. (If the pellet is allowed to swell in water, the nuclei break and release their nucleoprotein to give a viscous gel.)

Fig. 1. Extraction of guinea-pig liver nuclei with iso-osmotic sucrose and NaCl solutions. The crude nuclei, prepared as described in the Experimental section, were extracted five times with 0.25*M*-sucrose-1 *mm*-β-mercaptoethanol-10*mm*-tris-HCl buffer, pH7.4, and then three times with 0.14*M*-NaCl-1 *mm*-β-mercaptoethanol-2*mm*-EDTA, pH7. The individual extracts were dialysed overnight against 1 *mm*-β-mercaptoethanol-2*mm*-EDTA, pH7 (20 vol.) and assayed for phosphorolytic activity. Sucrose extracts, ●; saline extracts, ○.

The isolation procedure of Chauveau *et al.* (1956) is considered to give nuclei that are relatively free from contamination (Roodyn, 1963; Siebert & Humphrey, 1965; Georgiev, 1967). Although improvements have been described (Maggio, Siekevitz & Palade, 1963; Blobel & Potter, 1966) that permit the isolation of extremely pure nuclei, the original method was used for the studies described in the present paper since it is quicker and more suited to the handling of large amounts of liver for preparative work. However, although our results suggest that guinea-pig liver polynucleotide phosphorylase is associated with the nuclear

Table 2. *Lipids associated with the particulate guinea-pig liver polynucleotide phosphorylase*

A suspension of the particulate enzyme (0.5 ml; 1.65 mg of protein/ml) was shaken for 20 min in a stoppered glass centrifuge tube with 6 ml of chloroform-methanol (2:1, v/v). The mixture was centrifuged and the clear chloroform-methanol layer removed and evaporated to dryness in a rotary evaporator. The residue was dissolved in chloroform and the solution applied to a t.l.c. plate coated with activated (110°C for 2 h) silica gel G. The chromatogram was developed with chloroform-methanol-water (65:25:4, by vol.) (Wagner, Hörhammer & Wolff, 1961). To locate the lipids, the plates were sprayed with 50% (v/v) H_2SO_4 and heated carefully over a Bunsen burner. The results of a typical separation are shown. The material that had the same R_F as cholesterol (spot 5) was examined further in a separate experiment. The lipids were separated as described above and located by staining with iodine vapour (Randerath, 1964). After the iodine had evaporated, spot 5 was scraped off the t.l.c. plate and the lipids were extracted from the silica gel by the method of Bligh & Dyer (1959). The resulting chloroform extract was concentrated, the solutions was applied to a second, similar t.l.c. plate and the chromatogram was developed with light petroleum (b.p. 30–60°C)-diethyl ether-formic acid (75:25:1.5, by vol.). The plate was exposed briefly to iodine vapour and the iodine was then allowed to evaporate. The chromatogram was then treated with the reagent of Lowry (1968), which is specific for cholesterol or cholesteryl esters. The two stains revealed three components, of which only one contained cholesterol or cholesteryl esters.

Spot no.	Standards		Extract	
	Lipid	R_F	R_F	Intensity
1	Lysophosphatidylcholine	0.096	0.097	Faint
2	Phosphatidylcholine	0.31	0.30	Dark
3	Phosphatidylethanolamine	0.62	0.61	Medium
4	—	—	0.72	Very faint
5	Cholesterol (plus other lipids)	0.93	0.94	Dark
6	Mixture of mono-, di- and tri-glycerides	0.98	0.98	Very dark

membrane, further detailed studies must be undertaken to localize it precisely in the cell.

The particulate enzyme could be dissolved by addition of sodium dodecyl sulphate to the suspension to a final concentration of 5 mM. Removal of the detergent by dialysis did not cause the precipitate to reappear, but the enzyme was inactivated. When magnesium chloride was added to the dialysed solution (final concn. 10 mM), an inactive precipitate similar in appearance to the original particles formed. Engelman (1968) has shown that the membrane of *Mycoplasma laidlawii* can be dissolved by 10 mM-sodium dodecyl sulphate and that it is reconstituted during dialysis against 20 mM-magnesium chloride, so these properties of the particles are also consistent with the enzyme being membrane-bound.

Most bacterial polynucleotide phosphorylases have been isolated in a soluble form or associated with ribosomes (Grunberg-Manago, 1963; Kimhi & Littauer, 1967). However, Abrams & McNamara (1962) found that virtually all the polynucleotide phosphorylase activity of *Streptococcus faecalis* was present in the cell membranes when the latter were isolated by a gentle procedure involving formation of protoplasts followed by osmotic lysis. Further, the polynucleotide phosphorylase of the extreme halophile *Halobacterium cutirubrum* is associated with the cell membrane (P. Peterkin & P. S. Fitt, unpublished work).

Attempts to recover active polynucleotide phos-

phorylase from the particles in a soluble form have been unsuccessful so far.

Enzymic properties of guinea-pig liver polynucleotide phosphorylase. The proof that nucleoside diphosphates are the primary product formed from polyribonucleotides in the conditions of the phosphorolysis reaction catalysed by both the sucrose and saline fractions has been published (Fitt & See, 1970). The enzyme is therefore a polynucleotide phosphorylase. It should be noted that even the crude sucrose and saline fractions show a high degree of specificity for nucleoside diphosphates as products of the reaction despite considerable contamination with degradative enzymes that are removed during the purification.

Fig. 2 shows the effect of divalent cations on the phosphorolytic activity of the sucrose and saline fractions. Both required Mg^{2+} , although Mn^{2+} could replace the latter to some extent. The sucrose fraction was slightly active in the presence of Co^{2+} , but Ca^{2+} and Zn^{2+} were ineffective (the last three cations were not tested with the saline fraction). These results are in general agreement with the known metal requirements of bacterial polynucleotide phosphorylases (Babinet, Roller, Dubert, Thang & Grunberg-Manago, 1965). The Mg^{2+} optimum of both the sucrose and saline fractions was 4 mM (Fig. 3).

Figs. 4 and 5 show respectively the enzyme saturation curve and the time-course of the reaction with the sucrose fraction in the standard phosphoro-

Fig. 2. Influence of divalent cations on the phosphorolytic activity of guinea-pig liver polynucleotide phosphorylase. The (a) sucrose and (b) saline fractions were assayed for poly A-phosphorolysis activity with the indicated concentrations of $MgCl_2$, ●; $MnCl_2$, ○; $CoCl_2$, △; $ZnCl_2$ and $CaCl_2$, □.

Fig. 3. Influence of Mg^{2+} concentration on poly A phosphorolysis by guinea-pig liver polynucleotide phosphorylase. Enzymic activity of the sucrose and saline fractions was determined in the standard phosphorolysis assay conditions at the indicated Mg^{2+} concentrations. Sucrose fraction, ●; saline fraction, ○.

Fig. 4. Enzyme-saturation curve for guinea-pig liver polynucleotide phosphorylase. Phosphorolysis activity was measured in the standard assay with the indicated volumes of enzyme solution (928 units/ml).

lysis assay conditions with poly A as substrate. If the reaction mixtures were not shaken during the incubation there was a marked lag phase in the reaction, undoubtedly because of inaccessibility of the particulate enzyme to the substrate, but in

the standard conditions this was not evident except at low enzyme concentrations or during the first few minutes of assay. It can be seen that the reaction showed the expected dependence on time and enzyme concentration.

The influence of pH on the phosphorolytic activity of the sucrose fraction was studied in the range pH 7–10 by using three different buffers (Fig. 6). The maximum activity was observed in the range

Fig. 5. Time-course of poly A phosphorolysis by guinea-pig liver polynucleotide phosphorylase measured in the standard assay conditions (with 23.2 units of enzyme activity).

Fig. 6. Dependence of phosphorolysis activity of guinea-pig liver polynucleotide phosphorylase on pH. Enzymic activity of the sucrose fractions was determined in the standard poly A phosphorolysis conditions with 0.1M buffers of the indicated pH and composition: imidazole-HCl buffer, ●; tris-HCl buffer, ○; glycine-NaOH buffer, ▲.

pH 8.6–9.2 in tris-HCl buffers. The Mg^{2+} concentration in these experiments was 4mM, and all the curves were shifted to lower pH values by approximately 0.2 unit when the bivalent cation concentration was 10mM. Similar results were obtained with the saline fraction. Our earlier studies (Fitt & See, 1970) were carried out at pH 7.2 with 10mM-magnesium chloride. The specificity of the reaction is unchanged in the optimum conditions, but the activity of the enzyme is increased threefold. In some preliminary experiments we tried phosphate buffer, pH 7.2, but the

Fig. 7. Effect of NaCl and KCl on the phosphorolysis activity of guinea-pig liver polynucleotide phosphorylase. Enzymic activity was measured by the standard assay at the indicated salt concentrations. NaCl, sucrose fraction, 28.5 units, ○---○; KCl, sucrose fraction, 28.5 units, △—△; NaCl, saline fraction, 6.25 units, ▲—▲; KCl, saline fraction, 6.25 units, ●---●.

assays were then irreproducible owing to high and variable blanks.

The effect of sodium chloride and potassium chloride on the phosphorolytic activity of the sucrose and saline fractions was studied (Fig. 7), because of their known ability to stimulate phosphorolysis by some bacterial polynucleotide phosphorylases (Hendley & Beers, 1961; Fitt, Dietz & Grunberg-Manago, 1968). Both fractions were inhibited markedly by these salts, and were inactive when their concentration in the reaction mixture was 0.2M or higher. Thus, all fractions from the DEAE-cellulose column step in the purification had to be dialysed before assay, making elution with a salt gradient impracticable. It is of course possible that the effect of the salts was on the particulate complex as a whole and not specifically on the enzyme.

Fig. 8 shows the substrate specificity of the enzyme. It catalysed the active phosphorolysis of poly U, poly C and poly A, was less active towards tRNA or high-molecular-weight RNA, and had little or no activity with poly G (although a definite effect on the latter was detected after prolonged incubation). DNA was unaffected under these reaction conditions. Similar results were obtained when the Mg^{2+} concentration was increased from 4mM to 10mM. These results are again in general agreement with the reported substrate specificity of bacterial polynucleotide phosphorylases (Grunberg-Manago, 1959, 1963), although the

activity with poly G was unusually low. The apparent preference of the enzyme for pyrimidine-containing polyribonucleotides may be due in part to contamination by a phosphatase specific for purine ribonucleoside diphosphates.

Gajda & Fitt (1969) have shown that *Azotobacter vinelandii* polynucleotide phosphorylase undergoes an aerial oxidation accompanied by loss of activity and that the oxidized enzyme is reactivated by β -mercaptoethanol. When partially purified guinea-pig liver polynucleotide phosphorylase was dialysed overnight against 2mM-EDTA, pH 7, to remove β -mercaptoethanol, and assayed with the latter omitted from the standard incubation mixture, only 28% of the phosphorolytic activity remained. In the presence of β -mercaptoethanol (10mM) in the assay medium, 76% of the original activity was detected, so it is probable that the guinea-pig enzyme also undergoes a reversible oxidation.

The purified sucrose fraction catalysed both ADP- P_i and ATP- P_i exchange. Table 3 shows

that in both cases the exchange activity was specific for the adenine nucleotides, in contrast to the phosphorolytic activity. Similarly, both exchange activities were stimulated 10–90% by low concentrations (5 or 10mM) of sodium chloride or potassium chloride, or a mixture of both, whereas the phosphorolytic activity was inhibited 20–30% in these conditions. The pH optima for the ADP- P_i and ATP- P_i exchange activities were 7.8 and 7.4, respectively, both considerably lower than that of the phosphorolytic activity and, unlike the latter, were unaffected by the nature of the buffer. Finally, both exchange activities were inhibited by ApA, unlike the NDP-exchange activity of bacterial polynucleotide phosphorylases. These observations, together with the fact that the ratio of both exchange activities to the phosphorolytic activity decreased tenfold during the purification, all suggest that they are due to some contaminating enzyme. However, the stability during storage of the ADP- P_i exchange activity was similar to that of the phosphorolytic activity, whereas the ATP- P_i exchange activity was much less stable, so it is still possible that the ADP- P_i exchange is a genuine activity of the guinea-pig liver polynucleotide phosphorylase.

We have so far been unable to detect any nucleoside diphosphate polymerization activity in our preparations of the guinea-pig liver polynucleotide phosphorylase over a wide range of pH and divalent cation concentrations, both with and without oligonucleotide primers, salts or polylysine (cf. Fitt *et al.* 1968; Fitt & Wille, 1969). Although the crude extracts contained nucleases which degraded polyribonucleotides to nucleoside 5'-phosphates, probably the enzymes described by Lazarus & Sporn (1967) and Sporn, Lazarus, Smith & Henderson (1969), these nucleases together with contaminating phosphatases were almost entirely removed during purification, so the lack of polymerization activity could not have been due to contamination by degradative enzymes. A bacterial polynucleotide phosphorylase with no polymerization activity has been described (Thang, Thang & Grunberg-Manago, 1967, 1969), but it was also devoid of NDP- P_i exchange activity. It is generally

Fig. 8. Substrate specificity of guinea-pig liver polynucleotide phosphorylase. The phosphorolysis activity was measured in the standard assay with the indicated concentration of polyribonucleotide. Poly U, ○; poly C, ▲; poly A, ●; poly G, △; wheat germ rRNA, ■; yeast tRNA, □.

Table 3. Comparison of the substrate specificity for exchange and phosphorolysis of the particulate guinea-pig liver enzyme

Reaction	Nucleotide or polymer base (N)			
	A	C	G	U
	(c.p.m. incorporated)			
NDP- P_i exchange	289	43	23	26
NTP- P_i exchange	284	40	99	19
Poly N phosphorolysis	2091	2641	0	3253

accepted (Grunberg-Manago, 1963) that exchange catalysed by bacterial polynucleotide phosphorylases is due to the overall polymerization-phosphorolysis reaction operating in near-equilibrium conditions, and not to the formation of an intermediate enzyme-NMP complex as suggested by Olmstead (1958). Since the guinea-pig liver polynucleotide phosphorylase appears unable to catalyse the polymerization of nucleoside diphosphates, it will be of interest to establish if the ADP-P_i exchange activity associated with the preparation obtained by the method described in the present paper is indeed a genuine activity of the enzyme.

We thank the Medical Research Council of Canada for an operating grant (to P.S.F.) and Dr P. P. Proulx and Mrs E. Gaertner for advice and assistance in the separation and identification of lipids.

REFERENCES

- Abrams, A. & McNamara, P. (1962). *J. biol. Chem.* **237**, 170.
- Babinet, C., Roller, A., Dubert, J. M., Thang, M. N. & Grunberg-Manago, M. (1965). *Biochem. biophys. Res. Commun.* **19**, 95.
- Bligh, E. G. & Dyer, W. J. (1959). *Can. J. Biochem. Physiol.* **37**, 911.
- Blobel, G. & Potter, V. R. (1966). *Science, N.Y.*, **154**, 1662.
- Burton, K. (1956). *Biochem. J.* **62**, 315.
- Chauveau, J., Moulé, Y. & Rouiller, C. (1956). *Expl Cell Res.* **11**, 317.
- Dische, Z. (1955). In *The Nucleic Acids*, vol. 1, p. 302. Ed. by Chargaff, E. & Davidson, J. N. New York: Academic Press Inc.
- Engelman, D. M. (1968). In *Membrane Models and the Formation of Biological Membranes*, p. 203. Ed. by Bolis, L. & Pethica, B. A. Amsterdam: J. Wiley and Sons Inc.
- Fitt, P. S., Dietz, G. W., jun. & Grunberg-Manago, M. (1968). *Biochim. biophys. Acta*, **151**, 99.
- Fitt, P. S. & Fitt, E. A. (1967). *Biochem. J.* **105**, 99.
- Fitt, P. S. & See, Y. P. (1970). *Biochem. J.* **116**, 309.
- Fitt, P. S. & Wille, H. (1969). *Biochem. J.* **112**, 497.
- Gajda, A. T. & Fitt, P. S. (1969). *Biochem. J.* **112**, 381.
- Georgiev, G. P. (1967). In *Enzyme Cytology*, p. 27. Ed. by Roodyn, D. B. New York: Academic Press Inc.
- Grunberg-Manago, M. (1959). *J. molec. Biol.* **1**, 240.
- Grunberg-Manago, M. (1963). *Prog. nucleic Acid Res.* **1**, 93.
- Grunberg-Manago, M., Ortiz, P. J. & Ochoa, S. (1956). *Biochim. biophys. Acta*, **20**, 269.
- Hendley, D. D. & Beers, R. F., jun. (1961). *J. biol. Chem.* **236**, 2050.
- Kalnitsky, G., Hummel, J. P. & Dierks, C. (1959). *J. biol. Chem.* **234**, 1512.
- Kimhi, Y. & Littauer, U. Z. (1967). *Biochemistry, Easton*, **6**, 2066.
- Lazarus, H. M. & Sporn, M. B. (1967). *Proc. natn. Acad. Sci. U.S.A.* **57**, 1386.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
- Lowry, R. R. (1968). *J. Lipid Res.* **9**, 397.
- Maggio, R., Siekevitz, P. & Palade, G. E. (1963). *J. Cell Biol.* **18**, 267.
- Olmstead, P. S. (1958). *Biochim. biophys. Acta*, **27**, 222.
- Randerath, K. (1964). *Thin-Layer Chromatography*, p. 126. New York: Academic Press Inc.
- Roodyn, D. B. (1963). *Biochem. Soc. Symp.* no. 23, p. 20.
- Siebert, G. & Humphrey, G. B. (1965). *Adv. Enzymol.* **27**, 239.
- Sporn, M. B., Lazarus, H. M., Smith, J. M. & Henderson, U. R. (1969). *Biochemistry, Easton*, **8**, 1698.
- Thang, M. N., Thang, D. C. & Grunberg-Manago, M. (1967). *Biochem. biophys. Res. Commun.* **28**, 374.
- Thang M. N., Thang, D. C. & Grunberg-Manago, M. (1969). *Eur. J. Biochem.* **8**, 577.
- Wagner, H., Hörhammer, L. & Wolff, P. (1961). *Biochem. Z.* **334**, 175.