acids and possibly a sterol ester. Similar classes of neutral lipids were found in the sporophore of *A. bisporus* (Hughes, 1962). The acetone eluate from the column contains the acylated form of the carbohydrates found in the aqueous fraction. The composition of this fraction will be described elsewhere.

Phospholipids were found only in the chloroform-methanol eluate. T.l.c. in chloroform-methanol-water (65:25:4, by vol.) and spraying with specific reagents showed principally phosphatidylinositol and phosphatidylethanolamine. Phosphatidylcholine was absent. These identifications were confirmed by acid hydrolysis and paper chromatography, which demonstrated glycerol, inositol, ethanolamine and traces of serine, but no choline. Mild alkaline hydrolysis and paper chromatography in propan-2-ol-aq. ammonia (sp.gr. 0.88)-water (7:1:2, by vol.) showed glycerylphosphorylinositol, glycerylphosphorylethanolamine and glycerylphosphorylserine.

This work is the early part of a study of the composition of *A. bisporus*, in the fruiting and vegetative stages, which may provide some insight into the metabolic differences underlying the initiation of sporophore development. It may also help to elucidate the role of lipids in stimulating spore germination (Lösel, 1967).

This work was supported by grants from the National Science Council and The Agricultural Research Institute, Irish Republic.

- Folch, J., Lees, M. & Sloane-Stanley, G. H. (1957). *J. biol. Chem.* **226**, 497.
 Hughes, D. H. (1962). *J. Mushroom Sci.* **5**, 540.
 Hughes, D. H., Lynch, D. L. & Somers, G. F. (1958). *J. agric. Fd Chem.* **6**, 850.
 Lösel, D. M. (1967). *Ann. Bot. N.S.* **31**, 417.
 Rast, D. (1965). *Planta*, **64**, 81.

Rennin Coagulation of Cow's Milk

By J. V. WHEELOCK and J. P. PENNEY. (*School of Biological Sciences, University of Bradford, Yorks. BD7 1DP, U.K.*)

There are wide variations between milks of individual cows in the time taken to form a clot with rennin (EC 3.4.4.3). These variations may be due to factors such as Ca^{2+} concentration (Pyne, 1953) or pH (White & Davies, 1958) of the milk, which affect the coagulation phase, or to the amount and composition of the κ -casein, which could affect the primary phase of rennin action.

The primary phase may be conveniently studied by following the release of glycopeptides from κ -casein as determined by the increase in nitrogen soluble in the 12%-(w/v)-trichloroacetic acid filtrate

of the milk after the addition of the enzyme (Alais, Mocquot, Nitschmann & Zahler, 1953). Using the milk samples from individual cows, which were in mid-lactation and free of bacterial infections of the udder, we have observed that the clotting time of the milk decreased with increasing initial rate of the primary phase. This finding demonstrates that the primary phase is a critical factor in determining differences in clotting time between milks of individual animals. Further, we have observed that the initial rate of release of glycopeptides increased with the amount of glycopeptides released by rennin.

It is possible that at the concentrations used the enzyme may be present in excess, so that the initial rate of release is determined by substrate concentration. However, this explanation is unlikely because when the rennin concentration is varied there is a corresponding change in the initial rate of release of the glycopeptides and in the clotting time of the milk (Castle & Wheelock, 1970).

It has been postulated that the micellar κ -casein stabilizes the casein micelles and that the serum or 'rennin-accessible' κ -casein is responsible for coagulation of the milk (Parry & Carroll, 1969). When the serum κ -casein is hydrolysed the insoluble para- κ -casein becomes bound to the surface of several micelles and so joins them together into a clot. It follows from this that the amount of serum κ -casein will be directly related to efficiency of coagulation. Assuming that the total release of glycopeptides by rennin action can be used as a measure of the amount of serum κ -casein in the milk, then our results are in complete agreement with the ideas presented by Parry & Carroll (1969).

J. P. P. thanks the University of Bradford for a Research Studentship.

- Alais, C., Mocquot, G., Nitschmann, H. & Zahler, P. (1953). *Helv. chim. Acta*, **36**, 1955.
 Castle, A. V. & Wheelock, J. V. (1970). *Biochem. J.* **119**, 12P.
 Parry, R. M., jun. & Carroll, R. J. (1969). *Biochim. biophys. Acta*, **194**, 138.
 Pyne, G. T. (1953). *Chem. Ind.* p. 302.
 White, J. C. D. & Davies, D. T. (1958). *J. Dairy Res.* **25**, 267.

Kinetics of the Primary Phase of Rennin Action on Whole Milk

By A. V. CASTLE and J. V. WHEELOCK. (*School of Biological Sciences, University of Bradford, Yorks. BD7 1DP, U.K.*)

The kinetics of the primary phase of rennin (EC 3.4.4.3) action on whole milk have been studied by using the approach outlined in the previous

communication (Wheelock & Penney, 1970). An integrated form of the Michaelis-Menten equation was used to calculate values for K_m and V from the results obtained. Values for milks from different cows varied from 6.70×10^{-6} to 5.42×10^{-5} M for K_m and from 7.01×10^{-11} to 4.52×10^{-10} mol of glycopeptide $\cdot$ s $^{-1} \cdot$ l $^{-1} \cdot$ μ g $^{-1}$ of rennin $\cdot$ l $^{-1}$ of milk for V .

There was a linear relationship between initial velocity of release of glycopeptides and enzyme concentration, which varied between milks. The proportionality constant was used as an estimate of specific activity and values obtained were between 0.60 and 1.15 mg of N $\cdot$ min $^{-1} \cdot$ l $^{-1} \cdot$ μ g $^{-1}$ of rennin $\cdot$ ml $^{-1}$ of milk.

Taken together these results suggest that there is a change in the affinity of the enzyme for the κ -casein substrate between different milk samples. This could be due to the ionic environment of the milk, the amount of κ -casein in the milk, which may determine the size of the casein micelles (Sullivan, Fitzpatrick & Stanton, 1959), or the composition of the κ -casein. Variation in κ -casein composition is largely due to differences in the carbohydrates attached to it (Mackinlay & Wake, 1965).

Recently Sinkinson & Wheelock (1970) have shown that the *N*-acetylneuraminic acid/*D*-galactose molar ratio in the glycopeptides increases as the glycopeptides are released by rennin action. This finding suggests that the glycopeptides with a lower proportion of *N*-acetylneuraminic acid are released at a faster rate than those with a high proportion. Therefore the kinetic parameters for the release of each glycopeptide may be partly determined by the carbohydrates that are attached to it. For an individual milk sample, the values of K_m and V would therefore be an average value, determined by the relative proportions of different κ -caseins present. As the proportions probably vary between milks, then there should be corresponding changes in K_m and V . The specific activity would be determined by the composition of the κ -casein that is initially attacked.

We are grateful to the Agricultural Research Council for a grant to finance this work.

- Mackinlay, A. G. & Wake, F. G. (1965). *Biochim. biophys. Acta*, **104**, 167.
 Sinkinson, G. & Wheelock, J. V. (1970). *Biochim. biophys. Acta*, (in the Press).
 Sullivan, R. A., Fitzpatrick, M. M. & Stanton, E. K. (1959). *Nature, Lond.*, **183**, 616.
 Wheelock, J. V. & Penney, J. P. (1970). *Biochem. J.* **119**, 12P.

Carbohydrates of Bovine κ -Casein Glycopeptides

By J. V. WHEELOCK and G. SINKINSON. (*School of Biological Sciences, University of Bradford, Yorks. BD7 1DP, U.K.*)

The carbohydrates *N*-acetylneuraminic acid, *D*-galactose and 2-acetamido-2-deoxy-*D*-galactose (Alais & Jollès, 1961; Wheelock & Sinkinson, 1969) are attached to the glycopeptides released from κ -casein by the action of rennin (EC 3.4.4.3). Quantitative variations in these carbohydrates are largely responsible for the heterogeneity of κ -casein (Mackinlay & Wake, 1967). For an individual milk sample the average value for the carbohydrates attached to all the glycopeptides may be obtained by estimating the increase in carbohydrates attached to glycopeptides soluble in the 12% (w/v)-trichloroacetic acid filtrate after rennin action on the milk. In a previous investigation (Sinkinson & Wheelock, 1970a) we obtained molar proportions 3:5:2 and 2:2:1 for *N*-acetylneuraminic acid:*D*-galactose:2-acetamido-2-deoxy-*D*-galactose in the κ -casein glycopeptides. Values of 1:1:1 (Jollès, Alais, Adam, Delfour & Jollès, 1964) and 1:2:1 (Kuwata, Niki & Arima, 1969) have also been reported. There is, however, little information on the carbohydrates attached to the individual glycopeptides that contribute to the average value.

The glycopeptides present in the trichloroacetic acid filtrate of milk after rennin action were separated by ion-exchange chromatography. Then 30–40 mg of glycopeptides was dissolved in 2.5 ml of buffer (49 mM-KH₂PO₄–1 mM-Na₂HPO₄–0.1 M-NaCl adjusted to pH 5.1). This solution was eluted on a DEAE-cellulose column with 600 ml of buffer by using a linear 0.1–0.2 M-NaCl concentration gradient. Five major fractions and one minor fraction were obtained, which were each analysed for carbohydrates by g.l.c. (Sinkinson & Wheelock, 1970b).

Each glycopeptide contained *D*-galactose and 2-acetamido-2-deoxy-*D*-galactose. The first glycopeptide eluted was the only one to contain *D*-mannose, but was devoid of *N*-acetylneuraminic acid. The second glycopeptide contained *N*-acetylneuraminic acid, but the proportion was less than 1 mol/mol of glycopeptide and may have been due to contamination by the glycopeptides eluted later. These all contained approx. 1 mol of *N*-acetylneuraminic acid, 2 mol of *D*-galactose and 1 mol of 2-acetamido-2-deoxy-*D*-galactose/mol of glycopeptide.

The results show that the carbohydrate composition of the greater proportion of the κ -casein glycopeptides is similar and therefore will largely