size of these components, the extent of their separation and their equal labelling *in vivo* is consistent with the interpretation that they represent the mRNA species for the α - and β -globin chains. The 12S RNA also has a unique labelling pattern with [³²P]P_i *in vivo*, distinct from both 9S RNA and rRNA, and thus is unlikely to be an rRNA breakdown product.

The molecular weight of the 9S mRNA, as calculated from its sedimentation coefficient and its migration in polyacrylamide gels, is slightly greater than expected from the known size of the globin amino acid chain. DNA-RNA hybridization results with the mRNA, described in the following communication (Williamson, Morrison & Paul, 1970), may indicate a non-messenger function for a portion of the mRNA polynucleotide chain.

This work was supported by grants from the Medical Research Council and the British Empire Cancer Campaign for Research.

- Chantrenne, H., Burny, A. & Marbaix, G. (1967). *Prog. nucleic Acid Res. molec. Biol.* **7**, 173.
 Labrie, F. (1969). *Nature, Lond.*, **221**, 1217.
 Lanyon, G., Paul, J. & Williamson, R. (1968). *FEBS Lett.* **1**, 279.
 Laycock, D. G. & Hunt, J. A. (1969). *Nature, Lond.*, **221**, 1118.
 Lockard, R. E. & Lingrel, J. B. (1969). *Biochem. biophys. Res. Commun.* **37**, 204.
 Schapira, G., Dreyfus, J. & Maleknia, N. (1968). *Biochem. biophys. Res. Commun.* **32**, 549.
 Williamson, R., Lanyon, G. & Paul, J. (1969). *Nature, Lond.*, **223**, 628.
 Williamson, R., Morrison, M. & Paul, J. (1970). *Biochem. J.* **119**, 60P.

Hybridization with Deoxyribonucleic Acid of Messenger Ribonucleic Acid for Mouse Globin

By R. WILLIAMSON, M. MORRISON and J. PAUL.
 (Beatson Institute for Cancer Research, Glasgow C.3, U.K.)

The isolation of large amounts of purified mRNA for globin (Morrison, Williamson, Lanyon & Paul, 1970) has made the study of its hybridization to DNA possible. Because of the difficulty of obtaining sufficient amounts of high-specific-radioactivity RNA after labelling *in vivo*, the RNA was isolated unlabelled and then tritiated with [³H]dimethyl sulphate (Smith, Armstrong & McCarthy, 1967). Such RNA methylated *in vitro* shows the same specificity in DNA-RNA hybridization as RNA labelled *in vivo*. Mouse embryo DNA used in the hybridizations was purified by centrifugation through caesium chloride.

Hybridization was in formamide at 37°C, by the Gillespie-Spiegelman technique as modified by McConaughy, Laird & McCarthy (1969). The conditions are calculated to be equivalent to hybridization at 67°C in 4×SSC. A saturation value of 0.5% of the genome was found, corresponding to approx. 50000 gene copies for the mRNA, many more than anticipated. The hybridization reaction is competed out to 100% with unlabelled 9S mRNA, but only slightly competed by a mixture of 18S and 28S rRNA. Saturation occurs only at fairly high RNA/DNA ratios, but is very fast, being substantially complete in the first 10 min of reaction.

Crude thermal-denaturation profiles were obtained by heating the filters after hybridization in 1ml of 1×SSC at increasing temperatures. A low *T_m* for the hybrid was obtained, indicating a very extensive mismatching of base pairs.

The simplest explanation of these results is that inherent similarities between stretches of the amino acid sequences of proteins allow cross-hybridization to occur with extensive mismatching between one mRNA chain and the DNA genes coding for other polypeptide chains. This would certainly be expected for similar proteins such as the globins of haemoglobin and myoglobin, but the extent of hybridization seen here seems rather too large. It is also possible that the large extent of hybridization is due to the pairing, not only of the coding portion of the messenger RNA, but also to hybridization of the portion 'extra' to the expected messenger length (see Morrison *et al.* 1970). Such an extra sequence might be involved in initiation (Labrie, 1969) or in control of translation by 'ticketing' (Sussman, 1970), or be a transcribed portion of a control sequence for transcription (Britten & Kohne, 1968; Georgiev, 1969). In any of these cases such a sequence might be expected to be repeated at many points through the genome, not only at the site of globin mRNA.

This work was supported by grants from the Medical Research Council and the British Empire Cancer Campaign for Research.

- Britten, R. J. & Kohne, D. E. (1968). *Science, N.Y.*, **161**, 529.
 Georgiev, G. P. (1969). *J. theor. Biol.* **25**, 473.
 Labrie, F. (1969). *Nature, Lond.*, **221**, 1217.
 McConaughy, B. L., Laird, C. D. & McCarthy, B. J. (1969). *Biochemistry, Easton*, **8**, 3289.
 Morrison, M., Williamson, R., Lanyon, G. & Paul, J. (1970). *Biochem. J.* (This Meeting).
 Smith, K. D., Armstrong, J. L. & McCarthy, B. J. (1967). *Biochim. biophys. Acta*, **142**, 323.
 Sussman, M. (1970). *Nature, Lond.*, **225**, 1245.