Myoglobin Variants

By F. E. BOULTON, R. G. HUNTSMAN, H. LEHMANN, P. LORKIN and A. E. ROMERO-HERRERA. (*Department of Pathology, Lambeth Hospital, London S.E.11, U.K., and Medical Research Council Abnormal Haemoglobin Unit, University of Cambridge, Cambridge CB2 1QW, U.K.*)

The study of human haemoglobin variants has led to some insight into the variation of human genes. Some of the mutations are known to cause disease and others appear to be clinically benign. Some mutations are known to alter the physiological properties of the haemoglobin molecule. All known substitutions can be explained on the basis of single-point mutation in the DNA codon, but some of these are more frequent than others. Thus there are more changes involving the replacement of guanine by adenine in the mRNA than expected (Lehmann & Carrell, 1969).

It is therefore of interest to assess the frequency and type of other human protein variants. We have investigated nearly 6000 samples of human myoglobin by paper and acrylamide-gel electrophoresis. Three variants have been observed, but of necessity these methods will only detect substitutions producing a change of molecular charge. One would expect many neutral substitutions to remain undetected. Of the three variants it has been possible to identify two: 53 (D4) Glu $\rightarrow$ Lys (Boulton, Huntsman, Lorkin & Lehmann, 1969) and 138 (H16) Arg $\rightarrow$ Gln; both being guanine-to-adenine mutations of the mRNA.

During the survey a new haemoglobin was found in an English family, namely I-High Wycombe, with a mutation of residue $\beta 59$ (E3) Lys $\rightarrow$ Glu. This haemoglobin was easily identified because the tryptic hydrolysis between the tryptic peptides βV ($\beta 41-59$), which contains methionine, and $\beta TpV1$ ($\beta 60-61$) does not occur in the absence of the lysine residue at position $\beta 59$, and a new methionine-containing peptide $\beta 41-61$ was observed which was considerably more negatively charged than either peptide βTpV or peptide $\beta TpV1$ in normal haemoglobin. Amino acid analysis of this peptide showed the expected Lys $\rightarrow$ Glu replacement.

Boulton, F. E., Huntsman, R. G., Lorkin, P. A. & Lehmann, H. (1969). *Nature, Lond.*, **223**, 823.

Lehmann, H. & Carrell, R. W. (1969). *Br. med. Bull.* **25**, 14.

The Mechanism of Action of Rat Liver Glutamyl-Transfer-Ribonucleic Acid Synthetase

By MARY BURLEIGH and M. J. H. SMITH. (*Department of Biochemical Pharmacology, King's College Hospital Medical School, London S.E.5, U.K.*)

It has been suggested that the biosynthesis of aminoacyl-tRNA occurs in two stages, activation of the amino acids and their transfer to the tRNA (Hoagland, 1955; Hoagland, Zamecnik & Stephenson, 1957). The ATP-pyrophosphate exchange reaction has been considered to be a measure of the first stage. The participation of enzyme-bound aminoacyl adenylate as an obligatory intermediate (Wong & Moldave, 1960) was questioned by Loftfield & Eigner (1969) who studied the formation of aminoacyl hydroxamates. Salicylate has been shown to inhibit differentially the synthesis of aminoacyl-tRNA but to inhibit generally the ATP-pyrophosphate exchange reaction (Burleigh & Smith, 1970). Further, it has been shown that glutamate will not promote any ATP-pyrophosphate exchange in the absence of tRNA (Deutscher, 1967).

We have therefore studied the effects of salicylate on the glutamyl-tRNA synthetase activity present in a rat liver 105000g supernatant that has been passed through Sephadex G-25 to remove endogenous amino acids and ATP. The mechanism of inhibition was investigated by using fixed concentrations of two of the substrates and varying the third in the presence of different concentrations of salicylate. The drug showed a simple competitive inhibition with respect to glutamate and mixed competitive-non-competitive inhibitions with ATP and tRNA. These results are consistent with a mechanism for the action of the enzyme involving a compulsory addition of glutamate followed by the random addition of the other two substrates leading to the formation of a single quaternary complex (Fromm, 1967).

These observations strongly suggest that the esterification of glutamyl-tRNA involves the concerted interaction of the glutamate, ATP and tRNA, and similar considerations may apply to other aminoacyl-tRNA synthetases (Boyer, 1960). The formation of enzyme-bound aminoacyladenylate from certain amino acids, reflected by the ATP-pyrophosphate exchange, may occur as a side reaction.

This work was supported by a grant from the King's College Hospital and Medical School Research Committee.

Boyer, P. D. (1960). *A. Rev. Biochem.* **29**, 15.

Burleigh, M. & Smith, M. J. H. (1970). *Biochem. J.* **117**, 68P.