the smallest acceptor to give an incorporation product. The linkage of the incorporated radioactive glucose to the maltosaccharide was shown to be (1→6) by partial acid hydrolysis and methylation studies on a branched heptasaccharide formed from maltohexaose and radioactive glucose. The incorporation products were resistant to pullulanase, indicating that the radioactive glucose residue is present as a single stub. The action of β -amylase on the branched oligosaccharides gave further information on their structures.

The variation of the extent of incorporation with pH was studied with different acceptor substrates. With glycogen or Schardinger dextrans as acceptor substrates optimum incorporation occurred at pH 7.0 and 5.6, whereas with glycogen β -limit dextrin as acceptor a single optimum value at pH 7.0 was produced. With maltosaccharides of degree of polymerization 7 and 13 maximal incorporation occurred at pH 5.6. These results indicate that two related types of incorporation reaction that differ in pH optima are taking place.

We thank Professor D. J. Manners for his advice and encouragement during the course of this work.

Hers, H. G. (1964). *Adv. metab. Disorders*, 1, 2.

Manners, D. J. & Stark, J. R. (1967). *Abstr. FEBS 4th Meet., Oslo*, p. 14.

Nelson, T. E. & Larner, J. (1970). *Biochim. biophys. Acta*, 198, 538.

Properties of an α -Mannosidase from *Streptomyces* WL 6

By J. K. BAIRD and W. L. CUNNINGHAM. (*Department of Brewing and Biochemistry, Heriot-Watt University, Edinburgh EH1 1HX, U.K.*)

The purification and properties of α -mannosidases from several sources have been described recently. The enzyme from jack-bean meal (Li, 1967; Snaith & Levvy, 1968) hydrolyses α -mannosides containing aromatic aglycones, whereas the enzyme from an *Arthobacter* species shows little activity against such compounds but will degrade yeast mannan extensively (Jones & Ballou, 1969), a property that is not always associated with the other α -mannosidases.

Tanaka & Phaff (1965) observed that glucose and mannose were liberated from yeast cell walls by *Streptomyces* WL 6, an organism isolated by them from soil. They concluded that the streptomycete produced both glucanase and mannanase. During a study on the production of protoplasts from yeast cells by using the culture filtrate of this organism, we observed that an active α -mannosidase was also present. This α -mannosidase could be induced, in

quantity, by substituting yeast mannan as the carbon source, as found with *Streptomyces griseus* (Hockenhull, 1960).

Streptomyces WL 6 was grown for 6 days at 25°C in medium of pH 6.5, containing yeast mannan (0.5%), yeast extract (0.05%), tryptone (0.05%) KH_2PO_4 (1.36%) and salts. Ammonium sulphate was added to the culture fluid after removal of the cells. The fraction precipitated by 10–70% saturation was dialysed against 10 mM-phosphate buffer, pH 6.5, containing 0.1 mM- Ca^{2+} . Non-diffusible material was further purified by chromatography on DEAE-Sephadex.

A fraction was obtained that showed a 36-fold increase in purification of α -mannosidase activity measured with *p*-nitrophenyl α -mannoside as substrate. This preparation also hydrolysed manno-biose, mannotriose and mannotetraose derived from yeast mannan, methyl α -mannoside to a limited extent, and *p*-nitrophenyl α -glucoside, methyl α -glucoside, isomaltose, maltose and laminaribiose not at all. K_m values obtained were: for *p*-nitrophenyl α -mannoside, 3.3×10^{-3} M; for manno-biose, 2.0×10^{-3} M; for mannotriose, 7.5×10^{-3} M. Yeast mannan was slowly hydrolysed to mannose to a limited extent, indicating that the enzyme has exomannanase activity. The activity of the enzyme against *p*-nitrophenyl α -mannoside was stimulated by Ca^{2+} but inhibited by Cu^{2+} , Mg^{2+} and Zn^{2+} . There are therefore some differences between this enzyme and the α -phenylmannosidase present in *Streptomyces griseus* (Hockenhull, Ashton, Fantes & Whitehead, 1954).

We are indebted to Dr H. J. Phaff for a culture of *Streptomyces* WL 6. We thank the Science Research Council for a research studentship to J. K. B.

Hockenhull, D. J. D. (1960). *Prog. ind. Microbiol.* 2, 133.
Hockenhull, D. J. D., Ashton, G. C., Fantes, K. H. & Whitehead, B. K. (1954). *Biochem. J.* 47, 93.

Jones, G. H. & Ballou, C. E. (1969). *J. biol. Chem.* 244, 1043.

Li, Y.-T. (1967). *J. biol. Chem.* 242, 5474.

Snaith, S. M. & Levvy, G. A. (1968). *Biochem. J.* 110, 663.

Tanaka, H. & Phaff, H. J. (1965). *J. Bact.* 89, 1570.

An Effect of Amino Acid Analogues on Extension Growth in Pea Root Segments under Aseptic Conditions

By D. VAUGHAN and EVELYN CUSENS. (*Department of Biochemistry, The Macaulay Institute for Soil Research, Aberdeen AB9 2QJ, U.K.*)

Segments cut 2–4 mm from the root tip of 2-day-old seedlings of *Pisum sativum* (var. Meteor) elongate by 30% when cultured aseptically in water for 18 h. Elongation is enhanced by the

presence of sugars: sucrose > fructose > glucose. As there is no cell division, increases in length and fresh weight are due to growth of existing cells.

During extension growth the invertase (β -fructofuranosidase) activity follows the growth rate (Heyes & Vaughan, 1967). Since Brown (1963) postulated that extension growth is an expression of protein synthesis, the effect on growth and changes in invertase activity of several amino acids and some of their analogues was studied over 18 h in the presence of 2% (w/v) sucrose.

Methionine, phenylalanine and proline did not affect extension growth compared with controls. Two analogues (ethionine and *p*-fluorophenylalanine) inhibited extension growth some 60% at 3 mM; inhibition was measurable after 6 h. The inhibitions could be prevented by adding 3 mM-methionine and -phenylalanine to the culture medium.

In sucrose media, with or without amino acids, changes in invertase activity closely follow the growth rate, being maximal after 10–12 h and thereafter declining. Ethionine and *p*-fluorophenylalanine prevented increases in invertase activity, an effect abolished by adding the natural amino acid. The increase of acid phosphatase activity was similarly affected, but there were no changes in the total protein content of segments cultured under any of the conditions described.

The cell walls contain about 20% of the invertase activity and 90–95% of the 4-hydroxy-L-proline of the cell. The hydroxyproline content of the wall trebles during incubation in sucrose. Hence segments were cultured in the *cis* or *trans* isomers of this imino acid. Extension growth was enhanced by 15–20%, but invertase activity was depressed by 20–25% compared with the sucrose controls. The effects are abolished by adding proline.

The results support the idea that protein synthesis is required for cell growth. But results with hydroxyproline raise the question as to whether this imino acid is an analogue of proline or exerts its effect on growth because of its special position in cell-wall proteins.

We thank Professor L. Fowden for the *cis*-4-hydroxy-L-proline preparation.

Brown, R. (1963). *Brookhaven Symp. Biol.* **16**, 157.

Heyes, J. K. & Vaughan, D. (1967). *Proc. R. Soc. B*, **169**, 89.

Transfer of Adsorbed Bilirubin to Specific Binding Proteins

By J. A. T. P. MEUWISSEN and K. P. M. HEIRWEGH.
(Liver Physiopathology Laboratory, Rega Institute,
University of Leuven, Leuven, Belgium)

In a screening study of the interaction of alkaline bilirubin solutions with water-insoluble compounds

in bead form, it was observed that cross-linked dextran behaves as a relatively strong adsorbent (more than 95% adsorbed on Sephadex G-10 at pH 9.0). Alkaline destruction of bilirubin was drastically decreased by the presence of Sephadex in the suspension. A saturation limit of the solid matrix for the pigment could be demonstrated and measured quantitatively.

After stepwise neutralization of the suspension with phosphate buffers (final pH 7.4) combined with a washing at each step, followed by a final washing with water to remove ions, a bilirubin-Sephadex powder was prepared by freeze-drying and stored *in vacuo* over P_2O_5 .

In agitated suspension of the bilirubin-Sephadex complex (in buffers pH 4.0–10.0), the pigment remained adsorbed below pH 8.0. This is probably due to the extremely low solubility of bilirubin at these pH values (Brodersen & Theilgaard, 1969). In contrast, when the suspension contained specific bilirubin-binding proteins, rapid solubilization of the pigment took place. The adsorbed bilirubin seems to be present in a kind of activated state as, in otherwise identical conditions, solubilization of crystalline bilirubin did not occur.

The fast desorption of the pigment from the bilirubin-Sephadex complex was exploited to construct a sensitive assay of bilirubin-binding capacity. Bilirubin-Sephadex complex and either bovine serum albumin, lysozyme, β -lactoglobulin, rat serum, human serum or rat liver cytosol were incubated with continuous agitation. Lysozyme and β -lactoglobulin did not desorb the pigment. In the other preparations the amount of solubilized bilirubin was related hyperbolically to protein concentration. Equilibrium was attained after 16 h at 4°C or after 4 h at 25°C. On the basis of total protein concentration the amount of bilirubin solubilized by rat liver cytosol was about one-tenth that found with rat serum.

If during the preparation of the bilirubin-Sephadex complex the saturation limit of the solid matrix was exceeded, very high bilirubin/protein ratios were found in the subsequent binding assays (up to 33 mol of bilirubin/mol of albumin). This is probably due to the presence on the solid matrix of loosely bound bilirubin. It is likely that the desorbed pigment in this case is largely in colloidal form (R. Brodersen, personal communication), as indicated by spectrophotometry and restoration of the original bilirubin spectrum by alkalization (Brodersen & Theilgaard, 1969).

Brodersen, R. & Theilgaard, J. (1969). *Scand. J. clin. Lab. Invest.* **24**, 395.